




Digitized by the Internet Archive
in 2010 with funding from
University of Toronto

<http://www.archive.org/details/memoirs08harv>



51853
6/6/98

MEMOIRS

OF THE

MUSEUM OF COMPARATIVE ZOOLOGY,

AT

HARVARD COLLEGE.

VOL. VIII.

41853
6/6/98.

UNIVERSITY PRESS, CAMBRIDGE.

JOHN WILSON AND SON.

1881—1884.

OL
-
H²E
10

CONTENTS.



- No. 1. THE IMMATURE STATE OF THE ODONATA. PART II. Subfamily *Eschmia*. By LOUIS CABOT. pp. 39. With 5 Plates. August, 1881.
- No. 2. EXPLORATION OF THE SURFACE FAUNA OF THE GULF STREAM UNDER THE AUSPICES OF THE COAST SURVEY. III. PART I. The *Porpitiidae* and *Velellidae*. By ALEXANDER AGASSIZ. pp. 16. With 12 Plates. July, 1883.
- No. 3. THE REPTILES AND BATRACHIANS OF NORTH AMERICA. By SAMUEL GARMAN. Published by permission of the Kentucky Geological Survey. pp. 185. With 10 Plates. June, 1883.



Memoirs of the Museum of Comparative Zoölogy

AT HARVARD COLLEGE.

VOL. VIII. No. 1.

THE

IMMATURE STATE OF THE ODONATA.

PART II.—SUBFAMILY ÆSCHNINA.

BY LOUIS CABOT.


WITH FIVE PLATES.

CAMBRIDGE:
Printed for the Museum.
1881.



THE
IMMATURE STATE OF THE ODONATA.

PART II.—SUBFAMILY ÆSCHININA.



P R E F A C E.

UP to this time seven species only have been described by H. Hagen. One from Brazil, *Gynacantha* spec.; the others from Europe, *Anax formosus*; *Æschna rufescens*, *grandis*, *cyanea*, *affinis*; *Brachytron pratense*.

There have been figured before,—

Anax formosus; Muralto; Drury? Evans; Dufour.

Æschna cyanea; Swamerd? Frisch? Reaumur; Donovan; Dufour.

Æschna mixta; Reaum?

Æschna grandis; Roesel; De Geer; M. Harris.

Æschna rufescens; Dufour?

Æschna constricta; Samborn.

In all six species, three only, however, are recognizable, and none of the figures are accurate enough for specific determination. Of the twenty-four species now described and figured, containing the types of the seven species described by H. Hagen and others, there are of

Europe, nine.

America, nine; seven from U. S. A.

Asia, four.

Africa, one.

Half of the species described were raised, viz., the nine European species and three United States species. The actual number of described species of *Æschnidae* has not been exactly ascertained, but the twenty-four immature states given are about one tenth. In the United States there are twenty-one species known; of these, seven in the immature state are given. In Europe, thirteen are known; of these, nine in immature states are given. To make any general remarks on the classification of *Gomphina* and *Æschnina* has been found impossible, until the *Libellulina* and *Cordulina* have been described.

The following grouping is the best that can be made at present :—

Mask flat (agreeing with Agrionina), — all *Æschmina* and *Gomphina* (except *Cordulegaster*).

Mask, spoon-shaped (agreeing with *Libellulina*), — *Cordulegaster*.

Antennæ, four-jointed and flattened, — *Gomphus* and *Allies*.

Antennæ filiform and seven-jointed, — *Cordulegaster* and *Æschmina*, excepting two aberrant genera having six and five joints.

All legs with tarsus three-jointed, — *Æschmina* and *Cordulegaster*.

All legs with tarsus two-jointed, — *Ictinus*.

The two anterior pairs of legs with tarsus two-jointed, — *Gomphina*.

These general characters are sufficient to determine the place of any nymph.

AUGUST, 1881.

BIBLIOGRAPHY.

1634. Th. Moufet, *Insectorum Theatrum*, p. 321 and p. 322, figured a nympha without agreeing description, as *Scolopendra marina*. The figure is recognizable only for the genus.
1684. F. Muralto, *Ephem. Nature Curios.* Dec. II. Ann. II. p. 194. Figured and described as *Phryganeon Perlac*, the nympha of *Anax formosus*.
1706. Anonym. (probably W. Homberg). *Observations sur l'insecte poisson, qui se transforme en demoiselle.* *Histoire Acad. des Sc. Paris*, p. 9, described shortly the nympha of an *Æschna* and the transformation.
1730. G. L. Frisch, *Beschreibung von allerlei Insecten in Teutschland*, T. VIII. pl. x. p. 20, figured the full-grown nympha of a female (perhaps *Æ. grandis* or *Æ. cyanea*), with a good description.
1732. L. Roberg, *De Libella lacustri*, Upsalæ 4^{to}, p. 9, copies the figure given by Frisch without description.
1737. J. Swammerdam, *Biblia naturæ*, pl. xii, f. 4, figured the nympha of probably *Æ. cyanea*.
1742. De Reaumur, *Mém.* Vol. VI. pl. 36, f. 3, 4, figured the nympha of *Æ. mixta* or *Æ. affinis*; and pl. 39, f. 1-4, the nympha of *Æ. cyanea* in the act of transformation. The general description and the observations are excellent, but not sufficient for specific characters.
1743. E. Hebenstreit, *de insectorum natalibus*, f. cc, figured the nympha of an *Æschna*. Species not determinable.
1749. J. Roesel, *Insectenbelustigung*, Vol. II. p. ii. pl. 3, f. 1-9, figured very well the nympha of *Æ. grandis* in all stages; and pl. 4, f. 10-12, in the act of transformation. The prolix description is not sufficient for specific characters.
1752. J. Bartram, *Observations on the Dragonfly, or Libellula, from Pennsylvania*, *Philos. Transact. London*, Vol. XLVI. pp. 323 and 400, described shortly the nympha of an *Æschna* and the transformation.
1770. D. Drury, *Illustr. of Nat. Hist.*, Vol. I. pl. 47, f. 3, figured the nympha of perhaps *Anax formosus*. The description, p. 115, is not sufficient.
1771. Ch. De Geer, *Mémoires*, Vol. II. pl. 19, f. 12-19, figured the nympha of *Æ. grandis*, with a good description.

1772. M. Harris, Exposition of Engl. Insects, pl. xii, f. 3, figured the nympha of *Æ. grandis*.
1793. E. Donovan, the Natural History of British Insects, Vol. II. pl. 44, figured as *Libellula depressa*, the nympha of *Æschna* (perhaps *cyanea* or *grandis*). The figure is good, except some minor errors.
1826. Kirby and Spence, Introduction, etc., Vol. III. pl. 16, f. 5, figured as *Libellula*, the nympha of an *Æschna*. Described in letter xxx.
1830. (Rennie), Insect Transformations, p. 137, has a copy of Roesel's transformation of *Æ. grandis*.
1845. W. F. Evans, British Libellulina, pl. 1, f. 20, figured the nympha of *Anax formosus* as *Æschna*.
1852. Leon Dufour, Etudes anatom. and physiolog. sur les larves des Libellules. Ann. sc. natur. ser. 3. Vol. XVII, pp. 65-110, pl. 3. He is the first to attempt a systematical arrangement of the nymphæ. He described and figured *Æ. grandis* (= *A. formosus*), *Æ. De Geerii* (*Æ. rufescens*), *Æ. inominata* (*Æ. cyanea*).
1853. H. A. Hagen, Stettin. Entom. Zeit. Vol. XIV., has given a review of the literature, and described seven species,—*A. formosus*; *Æ. cyanea*, *affinis*, *rufescens*, *grandis*; *B. pratense*; *Gynacantha spec.*
1857. F. Brauer, Neuroptera Austr. pp. xiv.-xvi., gives an excellent review of the nymphæ. He had raised himself a large number of species, and to him is due most of our knowledge about the species. The nymphæ described by Hagen were mostly communicated to him by F. Brauer.
1862. Fr. Sanborn, Tenth Ann. Rep. Mass. Board of Agric., p. 149, figured the nympha of *Æ. clepsydra* (= *Æ. constricta*). The figure is reproduced in A. S. Packard's Guide, p. 602.
1879. Miss Olga Poljetajewa, Mémoires Russian Entom. Soc., Vol. XI. (in Russian language) described the Odonata from St. Petersburg and the nymphæ of *Æ. juncea*, *viridis*, both shortly, and of *Æ. grandis* very detailed and complete.

General descriptions and figures are to be found in the works of Vander Linden, Charpentier, Burmeister, Latreille, Westwood, Lacordaire, Rambur, Stephens, Newport, Newman, and in a large number of popular works; but they are not detailed enough for species or genera. Fossil nymphæ are figured and described. The literature is given by Heer and Hagen.

THE
IMMATURE STATE OF THE ODONATA.

PART II. — SUBFAMILY ÆSCHININA.

ÆSCHININA.

BODY elongated. Length from three to six times the breadth. Gills in the end of abdomen. Head large, breadth greater than length, except in *Gynacantha*, in which the length is the greatest. The form of head, mostly square, depends upon situation and form of eyes. Eyes always large and produced at inner hind angles in triangular lobe, separated from each other by advanced portion of occiput, which is generally raised and cut square at front border. The eyes vary in prominence, being least prominent in *Gynacantha*, more so in *Anax*, and most prominent in *Æschna*; — the line behind the eyes is sometimes straight, sometimes oblique, and sometimes curved. Vertex divided into two parts more or less well defined; the posterior portion is generally more nearly square than the anterior, and has generally indications of ocelli. The anterior portion is cut straight in front, and is generally somewhat semicircular; both portions are about on the level of the eyes. On each side of the vertex is an elongated polished spot, in front of which are inserted the antennæ. Antennæ short, slender, and with two exceptions seven-jointed. The two basal joints are globular, stouter than the others; the third, the longest, generally longer than the two basals taken together; fourth usually shortest; following joints longer. In *Gynacantha* the sixth joint is the longest. *E. Heros* has only six-jointed antennæ; the last joint the longest. In the *La Guayra* specimen the antennæ are abnormal, being only five-jointed. Occiput rather flat, shorter in *Brachytron* than in the other described species; sides more or less oblique, sometimes nearly straight; hind angles more or less rounded; in *Gynacantha* forming a sharp angle; hind border more or less notched, approaching in some species a straight line. That part of the head which lies before the eyes forms a

somewhat semicircular space, and consists of the following parts: a transverse short part, representing the front of the imago, separated by a well-marked border from a similar space, representing the epistoma and rhinarium joined in front to the large transverse upper lip, the front part of which is larger than the base and has oblique sides. On each side next to the upper lip, the base of the mandibles is visible. The mandibles, maxillæ, and tongue offer as far as observed no special characters. Mask flat, long, covering the mouth parts beneath, as far as upper lip; extending to between middle legs, except in *Gynacantha*, in which it extends to hind legs. The fore border is always broader than base, never more than twice as large, and the whole mask is more or less gradually enlarged forward. Side bent up and narrowly margined. The middle third of fore border is produced and more or less rounded, sometimes forms an obtuse angle, and is always cleft, generally not deeply, — in *Gynacantha* more deeply, surmounted by a comb of small hairs; on each side the cleft there is sometimes a small tooth, which in *Gynacantha* is strongly developed. Palpus consists of a narrow, nearly straight lobe, either straight at tip or slightly rounded, produced at inferior angle in more or less developed tooth, and more closely meeting the opposite one. The inner edge is either smooth or finely denticulated; movable hook, strong, rounded, very sharp, somewhat bent towards the tip and generally reaching the base of opposite one, and exceptionally it either extends beyond the base of opposite or does not quite reach it. Prothorax small, rather convex, rounded behind, more or less produced at the sides; the stigmata are large, open, and transverse, placed behind the prothorax and not covered by it; except in the *La Guayra* specimen, in which they are completely covered. The sides of the prothorax are produced above the fore legs in two processes, more or less divided and of varying proportion to each other. The form and size of these processes, being mostly rather distinct in form and size in the different species, seem to afford good specific characters. Analogous but less developed processes are found above the middle legs, and indications of them above the hind legs. Legs equally distant at base, or so nearly as not to give any character of importance, except in the *La Guayra* species, in which the hind legs are markedly more distant at base. Legs slender, not reaching tip of abdomen; femora and tibiæ of nearly equal length; tarsi half length of tibiæ, three-jointed; — basal joint very short; claws strong, bent, sharp. Thorax comparatively small. Wing cases large, reaching fourth segment or beyond.

Abdomen large, long, more or less tapering, more than half length of body, rounded above, flat beneath. Segments of equal length, tenth always, and occasionally the ninth shorter, no dorsal spines except in the La Guayra species. Lateral spines always on segments seven to nine, generally on six to nine, and exceptionally on five to nine and four to nine. Appendages as long or longer than the two last segments; inferiors long, sharp, triangular; middle appendage generally a little shorter, notched at tip, — sometimes of equal length, with undivided or split tip. Lateral superiors cylindrical, generally shorter, exceptionally as long and sharp as inferiors. Male has on the basal portion of middle appendage a triangular projection; short, sometimes cut at tip. Genitals not very distinctly marked; on ventral segments two, three, and nine. Female valve is visible on segment nine, sometimes reaching end of segment.

The whole body is either smooth, or has a granulated appearance, which is due to microscopical spines and small holes, sometimes with flattened yellow or pale colored hairs of differing size, never however such as to give a hairy look. The La Guayra species is distinguished by its strongly sculptured appearance. Generally, the occiput has on each side the median line a polished depression, and on the hind angles a series of parallel polished bands, separated by narrow rough lines. The abdomen has on each segment, less distinct on the first and apical segments, eight polished impressions; two near the median line and two on the edge of abdomen on each side. Color pale gray, sometimes darker or brownish black.

There are frequently black spots on occiput, and the sides near the eyes are sometimes bordered with black. Legs have frequently darker rings on femora and tibiæ, sometimes on tarsi. Abdomen has sometimes a broad dorsal band, pale in color, divided in centre by a band which is darker, except along the median line, which is again pale. Appendages have the tip sometimes darker than the basal part.

GYNACANTHA.

Rambur, *Hist. des Neuroptères*, p. 209.

Body elongated, very slender, flattened. Length of head greater than breadth. Occiput deeply notched at base; sides straight, forming right angle with base. Eyes narrow in proportion, two thirds length of head, not prominent, broadest in middle, not much produced at inner angles. Antennæ seven-jointed; sixth joint longest. Mask extending between hind legs, very

long, narrow, middle third of front border somewhat produced in two triangular lobes, rather deeply cleft; on each side of cleft highly developed spinous processes. Legs very slender. Processes enclosing right angle; posterior longer, anterior thicker. Abdomen long; lateral spines on segments six to nine. Appendages sharp, middle one as long as the inferiors, not deeply notched; laterals somewhat shorter, male projection conical.

I. GYNACANTHA? SPECIES.

PLATE III. FIG. 2.

Hagen, Stettin Zeit. XIV. p. 268.

Locality unknown, probably Brazil. Nymphæ two, male and female not fully grown, types described by Hagen. Length, 42 mm.; breadth, 7 mm. A smaller nympha from Charleston, S. C., and one from Florida, and some 25 mm. length from Rio San Francisco, Brazil, Thayer Exp., are identical.

Body exceptionally slender and elongated, flat. Head very flat, and longer than broad. Eyes elongated, two thirds the length of head, narrow, moderately prominent, at the hind inner angles somewhat produced; separated from occiput by straight line. Ocelli hardly discernible, space between the eyes in general shape almost square. The two parts of the vertex are not very distinctly marked. The lobe at the sides large and polished; part behind the eyes one third the length of head; sides straight; occiput long and deeply notched; hind angles acute, and with polished bands separated by roughened lines on upper side. Antennæ very slender, little longer than head, exceptional in having the sixth joint longest. Mask extending to between hind legs, very long and narrow, enlarged abruptly at apical third to twice the width of basal portion; middle third of front border deeply cleft, into two very prominent teeth, placed one on each side on the inner slope of the cleft. Palpus finely denticulated, cut nearly straight at tip; hooks of palpus prolonged so as to more than meet; movable hooks extending beyond the base of each other, cylindrical. Prothorax nearly as broad as back part of head, rounded behind. Processes enclosing right angle, posterior one longer, anterior thicker. Legs slender, hind legs reaching middle of seventh segment. Thorax small; wing cases in the largest specimen, extending only a little beyond first segment. Abdomen long, slender, small at base and enlarged as far as seventh segment; thence tapering, smooth; the eight impressions not very distinct. The four middle ones, transverse; dorsal band

darker on posterior segments, not well marked on the anterior ones. Side border somewhat projecting. Segments six to nine have lateral spines. Appendages nearly equal in length to segments nine and ten, sharp of nearly equal length, lateral superiors a little shorter; lateral inferiors longest; middle appendage very slightly notched. Male projection triangular, one third of length of middle appendage. Female valve distinguishable in young specimen. The determination is by supposition; no species living both in South Carolina and Brazil being known; indeed no species of *Gynacantha* is known in the United States. This is, however, not of so much importance, as Baron de Selys has described a species from Mexico. The nympha plainly belongs to *Æschnina*, and its abnormal shape strongly points to *Gynacantha*; at least no other genus so abnormal is known. The different species of *Gynacantha* are very similar and are widely spread; and as a species is described from Mexico and three from Cuba, one may also be found in South Carolina; this is even more probable by one full-grown female nympha, 40 mm. length, received from Haskinsville, St. John's River, Florida.

ANAX.

Leach, Edinboro' Encyclopædia, 1817. XI. p. 1, 137.

The largest of the *Æschnina*. Head large, flat, breadth greater than length. Eyes large, forming two thirds of head, prominent, broadest at posterior portion, much produced at inner angle. Antennæ have third joint longest. Occiput rounded at hind angles. Mask extending to middle legs, produced at middle third of front border in short, rounded, cleft lobe. Legs very slender. Abdomen large, lateral spines on segments seven to nine. Appendages long, sharp; middle one notched at tip. Male projection short, cut straight at tip.

2. ANAX FORMOSUS.

PLATE I. FIG. 1.

Selys, Revue des Odonates, p. 110.

Nympha, male and female, in alcohol and dry; one male just half transformed. Length, 52 mm.; breadth, 10 mm. Locality, Vienna, Austria, raised by Dr. F. Brauer; Silesia, Schneider. Coll. Mus. Comp. Zoöl.

Head very flat, rather broader than long. Eyes large, more prominent

than in *A. Junius*, separated from occiput by nearly straight line. Posterior part of vertex has indication of ocelli. Elongated elevations outside of vertex, large and marked. Antennæ small, similar to *Æschna*. Occiput short, hind angle rounded with indistinct bands on upper side, notch of hind border rather shallow. Mask long, extending to between middle legs, narrow, gradually enlarged forward. Middle third produced in rounded lobe, cleft, with comb of hairs. Palpus meeting opposite one; narrow, cut straight at end, with lower angle produced in short tooth, finely denticulated; movable hook, sharp bent, reaching base of opposite one. Prothorax as broad as occiput, rather short. Stigmata behind the prothorax uncovered. Processes nearly equal in length, short, blunt, enclosing right angle, posteriors largest. Legs long, slender, nearly cylindrical, hind legs longest, reaching end of segment eight. Femora longer than tibiæ; tarsi about one half length of tibiæ; apical joint making one half of tarsus; claws strong, sharp, bent; thorax comparatively small. Wing cases reaching nearly to segment five. Abdomen rather small at base, gradually enlarged to segment seven; thence tapering, rounded above, each segment with eight impressions. Dorsal band dark, interrupted throughout its length by paler line; lateral paler bands on each side; margin of abdomen darker, indistinctly marked with paler linear spots. Strong lateral spines on segments seven to nine; that on ninth as long as segment ten. Segments of equal length, tenth shorter. Inferior appendages long, sharp, as long as two last segments; middle one shorter than inferiors, notched; lateral superiors cylindrical, sharply pointed, half as long as middle one; male projection very small, cut square at tip, less than one half the length of laterals: female valve two thirds of segment nine, small. Nymphæ described are types of *H. Hagen*, *Stett. Zeit.* XIV. p. 268, and *F. Brauer*, *Neur. Austr.* p. xvi. *Dr. Hagen* considers this species to be the "*Æschna grandis*" described by *L. Dufour*, *An. Nat. Ser. 3, XVII.* p. 69, pl. 3, f. 1. The younger nymphæ from Vienna, mentioned by *H. Hagen*, *Stett. Zeit.* XIV. p. 267, as belonging perhaps to *Anax Parthenope*, prove to be young specimens of *Anax formosus*.

3. ANAX MAURICIANUS.

Rambur, *Neuroptères*, p. 184.

Nympha, male and female, dry, full-grown, and young. Length, 46 mm.; breadth, 10 mm.: the smallest young, length, 26 mm. Locality, Mauritius

Island, Mr. Pike, one full-grown male and two younger ones in alcohol, together with some imagos. Zanzibar, Mr. C. Cooke, ten specimens, male and female, in alcohol; length, 23 to 26 mm.

Very similar to *A. formosus*; differing by the two processes above the first legs, more blunt, enclosing an obtuse angle; mask narrower, the apical third more suddenly enlarged; middle appendage longer, nearly as long as the inferiors.

I had at first the specimens from Zanzibar, considered to belong to *A. formosus*, and the young ones are indeed similar to this species. One male, 34 mm. long, does not show the male projection; the smallest, 26 mm. long, has the wing cases just covering the first segment. Apparently, the characters are not so well expressed in the young ones; and I consider them probably belonging to *A. Mauricianus*, though this species is not yet received from Zanzibar.

4. *ANAX JUNIUS*.

PLATE I. FIG. 2.

Drury; Hagen, Synops. N. Amer. Neur. p. 118.

Nympha, male and female, dry and in alcohol. Length, 40 to 52 mm.; breadth, 10 mm. Locality, Springfield, Mass.; Boston and Cambridge, Mass.; Amherst, N. H.; Caledonia Creek and Crown Point, N. Y., Mr. J. A. Lintner; New Jersey; Detroit, Mich., Mr. H. J. Hubbard, a large number, young and full-grown; Charleston, S. C.; Pilichody, Mobile, Ala.; St. Louis, Mo., Mr. Ch. V. Riley, Kentucky; N. Mexico; Port de France, Martinique. Coll. Mus. Comp. Zool.

Very similar to *Anax formosus*. Differs in having the palpus not so straight at tip, with stronger end hooks; processes more obtuse, and enclosing an obtuse angle, of same size and shape; lateral appendages having the lower end more abruptly pointed than *A. formosus*, in which it is tapering. Male projection notched at tip. Very young specimens, 16 mm. long, from Springfield and from Detroit, have the palpus cut straight at tip; wing cases covering first segment. Besides the numerous specimens in alcohol, there are three raised by Mr. Ch. V. Riley and Mr. S. Henshaw.

Several specimens from San Diego, Cal., differ in having two black teeth in the middle of the comb of the front border of the mask, and the abdomen more bulky and broader. From the same locality and from the same collection *A. validus* was received. Nevertheless, it seems very doubtful

that nymphæ very similar to those of *A. Junius* should belong to the gigantic and very different *A. validus*.

A specimen in alcohol from New Mexico, Dr. Yarrow, differs by a shorter and broader mask. Perhaps it belongs to a new, still unknown species.

A very young specimen in alcohol from Key West Island, Florida, Mr. Garman, differs by a large head and very long legs. Perhaps it belongs to *A. longipes*.

5. *ANAX JULIUS*.

Brauer, Voyage of the "Novara," p. 63.

Seven nymphæ, male and female, in alcohol and dry. Length, from 28 to 50 mm; breadth, from 6 to 10 mm. Coll. Mus. Comp. Zoöl. Locality, Kanagava, Japan.

Very similar to the other forms of *Anax*. Processes as in *A. Junius*. Lateral appendages as in *A. formosus*. Male projection short, blunt; indication of notch at tip. In full-grown female, wing cases reaching segment five; palpus somewhat larger at tip, straighter, superior angle very little rounded; female valve, two thirds of ninth segment.

The nymphæ were received, together with the imago. No other species is known from Japan.

6. *ANAX GUTTATUS*.

Burm. Hdb. T. II. p. 340.

Twelve nymphæ, male and female, nearly full-grown, and young, in alcohol. Length, 60 to 40 mm.; breadth, 6 to 10 mm. Coll. Mus. Comp. Zoöl. Locality, Ebon, Marshall Island, Pacific Ocean, Rev. B. G. Snow.

Longer and less broad than the other species. Mask reaching the base of hind legs, narrow, more than four times longer than the breadth of base, gradually enlarged forward; front margin rounded, cleft; palpus narrowed to tip, superior angle obtuse, rounded; lower angle forming a very strong tooth; processes unequal, upper part blunt, lower part smaller, angular behind; enclosing a very obtuse angle, with a smaller median notch. Legs slender, long. Inferior appendages longer than the two last segments; lateral appendages sharply pointed; male projection narrow at tip, cut square; female valve small, half as long as the segment.

The nymphæ were received, together with the imago. The supposed identity is probable.

7. ANAX AMAZILI.

Burm. Hdb. II. p. 841.

Nympha, full-grown, male and female, dry; one in alcohol. Length, 53 to 58 mm.; breadth, 10 mm. Locality, Jamaica, Mr. H. J. Hubbard; Central America.

Similar to *A. Junius*, but considerably longer, though not broader; the mask four times longer than broad at the base (little more than three times longer in *A. Junius*); palpus cut straight at tip, with strong hook; processes above the first legs of equal size and length, enclosing a very obtuse angle; lateral appendages sharply pointed (in *A. Junius* dilated internally to the abruptly pointed tip); basal projection of the middle appendage of the male narrowed on tip and notched, about one third as long as the lateral appendages (broad on tip and emarginated, half as long as the lateral appendages in *A. Junius*).

A. Amazili is known from Cuba and Barbados, not yet from Jamaica; but there is no other species of *Anax* known from the Antilles.

8. ANAX SPECIES.

Nymphæ, full-grown and young, male and female in alcohol. Length, 20 to 42 mm.; breadth, 9 mm. Locality, Holy tank at Ibanca, East India, Rev. M. Carleton.

Similar to *A. Junius*. Processes more blunt; posterior smaller, enclosing an obtuse angle. Mask longer, the front margin less prominent, more rounded. Palpus narrowed at tip, upper angle rounded, lower one produced in a strong oblique end hook, below finely denticulated. Appendages, as in *A. Junius*, abruptly pointed, more so at the male; projection of male slightly notched at tip.

The determination of the species is not possible. One, *A. immaculifrons*, was received by Rev. M. Carleton, though not from the same locality. *A. Bacchus*, perhaps identical with *A. Parthenope*, lives in the sub-Himalaya; and *A. perplexus* Hagen, from Kooloo Valley, Himalaya, was received by Rev. M. Carleton; both sexes. Perhaps the nymphæ belong to the latter undescribed species.

ÆSCHNA.

Fabricius, Syst. Entom.

Body stout in proportion. Head, breadth twice the length, flat. Eyes large, very prominent, rounded forward, prolonged at inner angles in enlarged lobe. Antennæ have third joint longest. Mask extends to middle legs, large, broad; front border produced in short rounded lobe, cleft at middle. Legs longer and stouter than *Anax*. Abdomen broader; lateral spines on segment five or six to nine. Appendages long, sharp; middle one notched at tip; laterals shorter than middle one. Male projection conical; female valve reaching tip of segment.

9. ÆSCHNA RUFESCENS.

PLATE V. FIG. 4.

Vander Linden; Selys, Revue des Odonates, p. 129.

Nympha skins, male and female. Length, 40 to 44 mm.; breadth, 9 mm. Locality, Vienna, Austria. Raised by Dr. Brauer; Lueneburg, Mr. Heyer; Königsberg, H. Hagen.

Stout. Head large, flat, broader than long. Eyes large, very prominent, separated from occiput by nearly straight line. Occiput short, hind angles rounded and with polished bands on upper side; a polished spot on each side of the median line; hind border notched. Ocelli indicated. Anterior portion of vertex semicircular. Antennæ short, not extending beyond the eyes; seven-jointed, third joint longest. Mask extending between middle legs, stout; length greater than breadth; enlarged gradually forward; middle third somewhat produced; cleft in middle, with comb of hairs. Palpi broad, straight, cut square at end and produced only slightly in end hooks, finely denticulated. Movable hooks sharp and bent, thin, just reaching the base of their opposite. Prothorax not as broad as occiput, hind border rounded. Processes enclosing an acute angle, anterior one half length of posterior, both sharp. Legs rather slender, hind legs longest, extending to eighth segment; femora somewhat arcuated; two faintly-marked pale rings on lower portion of femora. Wing cases extending to fourth segment. Abdomen stout, very convex above, tapering from the middle to the tip, smooth; two parallel dark dorsal bands best defined at beginning of segments. Lateral spines on segments six to nine, strong, last shorter than segment ten. Appendages sharp, long as the last two segments; lateral

inferiors longest, lateral superiors two thirds the length of middle, middle nearly or quite straight at tip. Male projection stout, blunt, triangular, angular, half length of lateral superior appendages. Female ventral valve reaches end of segment. The described nymphæ are the types of Hagen Stettin Zeit. XIV. p. 267. Brauer, Neuropt. Austr. p. xvi.

Æ. De Geerii Dufour, Ann. se. nat. ser. 3, XVII. p. 69, pl. 3, fig. 3, is apparently this species. L. Dufour has overlooked the small lateral spine on segment 6.

10. *ÆSCHNA GRANDIS*.

PLATE II. FIG. 1.

Linnæus. Selys, Revue des Odonates, p. 131.

Nympha skins. Length, 44 mm; breadth, 9 mm. Locality, Lueneburg, Mr. Heyer; Kœnigsberg, Hagen.

Dr. Hagen observed the transformation of this species more than thirty years ago. The specimens are, however, no longer in his collection. A large number of nympha skins sent him by M. Heyer, unassorted, as *Æ. grandis* and *Æ. viridis*, were easily distinguishable as two species, but it could not be determined which were *Æ. grandis* and which *Æ. viridis*. Two specimens, both labelled *Æ. viridis*, pinned during transformation, were sent by M. Heyer; but of these one has only the head developed, and the other the parts only as far as the metathorax; and, as the color and shape are not fully shown, they can neither of them be considered as fully determined. These latter specimens, seeming probably *Æ. viridis*, Dr. Hagen refers the other nympha skins, which were sent before and were similar, to *Æ. viridis*; those dissimilar to *Æ. grandis*.

Very similar to *Æ. rufescens*; hind angles of occiput more sloping, line back of eyes more curved, palpus more convex and narrower. Processes different, of equal length, more widely separated, anterior thinner, somewhat bent outward at tip. Appendages different; middle shorter in proportion to inferiors than in *Æ. rufescens*. Lateral superiors about one half length of middle. Male projection shorter, tip sharper. This species is more brightly colored than *Æ. rufescens*. Abdomen brownish, dorsal band broad and dark, divided by a light median line. The nymphæ described are the types of Hagen, Stettin Zeit. XIV. p. 268, and F. Brauer. Neur. Austr. p. xvi.

Miss Olga Poljetajewa has raised this species, and fully described the nymphæ in Horæ Ross, St. Petersburg. 1879, 5, p. 15. The description confirms the identity with the above-described nymphæ.

11. *ÆSCHNA CYANEA*.

PLATE IV. FIG. 3.

Mueller. Selys, *Revue des Odonates*, p. 115.

Nymphæ, in alcohol and skins, male and female. Length, 48 mm.; breadth, 9 mm. Locality, Vienna. Raised by F. Brauer; Lueneburg, raised by M. Heyer; raised by Prof. Rosenhauer, Erlangen, Bavaria.

Body elongated and slender. Head large, flat, narrow behind. Eyes large, prominent, globular, occupying half the head; inner hind angles much produced. Line between occiput and eyes curved. Sides of head oblique, hind angles rounded. Occiput notched; on each side of median line a polished spot, on upper side of hind angles polished bands separated by roughened lines. Vertex, antennæ, and mouth parts as in *Æ. rufescens*. Head grayish, with paler band at sides; a pale spot on each side the median line of occiput. Mask extending between the middle legs; narrow at base, one half width of front border. Middle third of front border somewhat produced, at very obtuse angle, cleft with a comb of hairs. Palpi meeting broad, cut straight at ends, and produced very slightly into end hooks. Movable hooks sharp and thin, reaching base of the opposite one. Prothorax small, as broad as back part of occiput, ash gray at each side, pale at border. Processes short, scabrous, enclosing a little more than a right angle; anterior shortest; legs slender, hind legs reaching eighth segment. Femora and tibiæ ringed with two darker bands. Thorax not very large; wing cases reaching fourth segment. Abdomen long, slender, enlarged gradually to beyond the middle, thence tapering, grayish; dorsal band dark, divided by paler band; a pale band on each side of dorsal band. Segments of about equal length, except tenth, which is shorter. Lateral spines on segment six to nine. Inferior appendages as long as two last segments, triangular, sharp; lateral superiors one half length of inferiors. Middle appendage notched; male projection conical, rather enlarged at base, shorter than lateral appendages. The whole body finely granulated with what appears, when highly magnified, to be spines pointing backward, and with short yellow hairs inserted behind them. Genital parts of the male indicated, those of the female extending somewhat beyond the ninth segment.

The nymphæ described are types of Hagen, *Stettin Zeit.* XIV. p. 266; and F. Brauer, *Neur. Austr.* p. xvi. *Æschna innominata*, Dufour, *Ann. se. nat. ser. 3*, XVII. p. 69, pl. 3, f. 5, is apparently *Æ. cyanea*.

12. *ÆSCHNA JUNCEA*.

PLATE IV. FIG. 1.

Linnæus; Selys, *Revue des Odonates*, p. 116.

Three nymphæ, male and female, skins, by M. Heyer; five from J. Boll. Length, 44 m.; breadth, 8 mm. Locality, Lueneburg, Zurich. Five nymphæ raised by Mr. J. Boll, Switzerland.

Very similar to *Æ. cyanea*, a little shorter. Processes strikingly different. Anterior only half as large as posterior and much smaller. Granulation, color of legs, body, and appendages as in *Æ. cyanea*. The lateral appendages are more slender and more pointed, and the hind angles of occiput more rounded than in *Æ. grandis*.

The imago has been raised, and the nymphæ shortly described, by Miss Olga Poljetajewa, l. c. p. 15.

13. *ÆSCHNA VIRIDIS*.

PLATE V. FIG. 5.

Eversmann. Selys, *Revue des Odonates*, p. 127.

Nymphæ skins, male and female; and full-grown nymphæ, dry. Locality, Lueneburg, M. Heyer; Kœingsberg, H. Hagen. Length, 42 mm.; breadth, 9 mm.

Similar to *Æ. rufescens*, more slender. Head narrower behind; hind angles of occiput more rounded. Sides of head not so straight. Mask somewhat shorter and broader. Palpus narrower, more convex. Processes nearly equal length, enclosing nearly a right angle; tips bent outward. Abdomen more slender, uncolored; lateral spines on segments six to nine. Middle appendage shorter and more notched than in *Æ. rufescens*; lateral superiors short, one half length of middle one; male projection nearly as long as the lateral superiors, conical, tip rather sharp.

The species was raised by Miss Olga Poljetajewa, l. c. p. 15. She says the nymphæ is very similar to *Æ. juncea*, except that the lateral spines of abdomen are developed as well as in *Æ. grandis*.

14. *ÆSCHNA MIXTA*.

PLATE V. FIG. 2.

Latreille; Selys, *Revue des Odonates*, p. 122.

Nymphæ skins, male and female, raised by M. Heyer. Length, 33 mm.; breadth, 7 mm. Locality, Lueneburg.

Similar to *Æ. cyanea*. Body shorter, and stouter in proportion. Head flatter. Eyes larger and more prominent, separated from occiput by nearly straight line. Vertex more in the same plane with the eyes. Occiput shorter and straighter behind. Bands on hind angles less marked; color grayish; a pale elongated spot on each side of median line, and a narrow pale band on hind angle, often scarcely discernible. Mask long, extending through middle legs, gradually enlarged; front border a little more than twice width of base; middle third of front border somewhat produced and rather widely cleft, with comb of hairs. Palpus broad, rounded at upper angle, and produced in very short black teeth at lower angles; denticulation beneath discernible; movable hook, reaching the base of opposite one. Prothorax as broad as back part of occiput, rounded behind. Processes stout and blunt, not much separated, anterior more contracted. Legs very slender; hind legs extending to ninth segment, ash-gray; femora with two pale rings. Wing cases extending to middle of fifth segment. Abdomen short, stout, tapering from seventh segment; dorsum very convex; finely granulated as in *Æ. cyanea*, but with whitish hairs; color grayish; each segment with eight darker impressed spots, four on middle and two on each side near margin. Segments of nearly equal length, except tenth, which is half as long; lateral spines on segments six to nine; that of ninth nearly as long as tenth segment. Inferior appendages as long as two last segments, triangular, sharp; middle appendage notched, not so long as inferiors; lateral superiors rather more than half length of middle one, cylindrical, slender, and sharp. Male projection conical, rather sharp, nearly as long as lateral superiors. Male genital parts slightly marked; female valve extending somewhat beyond ninth segment.

15. *ÆSCHNA AFFINIS*.

PLATE V. FIG. 3.

Vander Linden; Selys, *Revue des Odonates*, p. 124.

Nympha, male, raised by F. Brauer. Length, 33 mm.; breadth, 7 mm. Locality, Vienna.

Similar to *Æ. mixta*. More slender in proportion. Eyes not so large or prominent. Occiput more rounded. Mask shorter and broader in proportion, extending through middle legs, enlarged gradually forward, so that front border is twice as broad as base; middle third not so produced as in *Æ. mixta*, and finely cleft, with a comb of hairs. Palpus broad and straight

at ends, and more nearly meeting than in *Æ. mixta*. Movable hooks as in *Æ. mixta*. Prothorax similar to *Æ. mixta*. Processes blunt, depression between them very shallow, of nearly equal size. Abdomen more slender, iron gray; the whole insect of this color; dorsal band darker, interrupted by lighter spots. Lateral spines on segments five to nine, that on fifth very small. Spine of ninth shorter than tenth segment. Finely granulated; hairs pale, and when present finer and shorter than in *mixta*. Appendages similar, except lateral superiors, only half as long as middle one, somewhat longer than male projection. The nympha described is the type of Hagen, *Stettin Zeit.* XIV. p. 267; and F. Brauer, *Neur. Austr.* p. xvi.

16. *ÆSCHNA EREMITICA.*

PLATE II. FIG. 2.

Scudder, *Proc. Boston N. H. Soc.* XI. p. 213.

Nymphæ, male and female, skins, and some not full-grown, in alcohol. Length, 39 to 47 mm.; breadth, 10 mm. Locality, mouth of Red River of the North; Hermite Lake, White Mountains, New Hampshire, Mr. Sanborn; Minnesota. *Coll. Mus. Comp. Zool.*

Similar in general shape to *Æ. rufescens*. Head much larger, remarkably large in proportion. Eyes large and prominent. Occiput not very deeply notched behind; sides oblique, and rounded at hind angles; three polished spots on middle; polished bands on hind angles. Antennæ, third joint nearly as long as first and second; fourth, about one half third; other, somewhat longer. Mask broad, passing through middle legs, gradually enlarged forward; base more than half width of front border; middle third produced in small rounded lobe, cleft and with comb of hairs. Palpus cut straight at end; rather flat, nearly or quite smooth. End hooks short; movable hook bent, sharp, reaching base of opposite one. Prothorax as broad as hind part of occiput. Legs slender, hind legs reaching ninth segment. Femora with dark ring on lower part. Processes long, of equal length, sharp, enclosing somewhat less than right angle, a little bent outward. Wing cases reaching segment five. Abdomen rather broad; enlarged gradually to seventh segment, thence tapering rather abruptly. Segments with four dorsal and four lateral darker spots, and two polished impressions on each side; tenth segment shortest. Lateral spines on segments five to nine, that on ninth nearly as long as tenth segment. All segments finely

granulated. Appendages longer than two last segments; sharp, middle one a little shorter than inferiors, deeply notched; lateral superiors cylindrical, sharp, nearly two thirds as long as middle one.- Female valve not quite reaching end of segment nine. Male projection conical, rather sharp, a little shorter than lateral superiors. The two male specimens are not fully developed in these parts, being only 30 and 33 mm. long. A full-grown female specimen from Minnesota, 47 mm. long, belongs apparently to this species. The determination seems to be justified, as Mr. Sanborn found nymphæ and imago at the same place, Hermit Lake, in middle of July.

17. *ÆSCHNA CONSTRICTA*.

PLATE III. FIG. I.

Say. Hagen, Synopsis N. American Neur, p. 123.

Nymphæ, two full-grown, in alcohol. Length, 41 mm.; breadth, 7 mm. Locality, Shooley's Mountain, Pa.; Boston, Mass., raised by Mr. S. Henshaw. Coll. Mus. Comp. Zool.

Similar to *Æ. eremita*. Longer and more slender in proportion. Head very similar. Mask narrower at base, more abruptly enlarged at apical third. Palpus finely denticulated, tip cut straight. Processes similar to *Æ. eremita*, not so long nor so sharp, the anterior less bent, the posterior a little larger. Legs similar in shape, shorter in proportion; hind legs not reaching segment eight. Granulation on abdomen very fine. Lateral spines on segments six to nine, an indication on fifth. Ninth segment half length of tenth. Inferior appendages as long as two last segments, very sharp; middle one not so long as inferiors; deeply notched; lateral superiors more than half length of middle one, cylindrical and abruptly sharpened. Male projection conical, somewhat shorter than lateral superiors, sharp. Hairs among granulation, small, yellowish.

Two full-grown female nymphæ, one from Maryland, the other without locality, and a small male seem to belong to this species. Male, in alcohol, from Lake Champlain, is sufficiently like to be placed in this species. Somewhat less slender, darker in color; two black rings on femora and tibiæ and on end of tarsal joints.

A very small nympha from Bethel, Me., and some younger ones from Cambridge, Mass., are colored in the same manner. Abdomen in many specimens dark, varied with short linear spaces of paler color and with large

black rings on femora and tibiæ. This nympha is figured by Mr. F. Sanborn, Report Mass. Board of Agricult., 1862, p. 149, and copied in A. S. Packard's Guide, p. 602. Mr. Sanborn stated it to be the nympha of one of the most common species, and Mr. Packard considered that it belonged to *Æ. constricta* or *Æ. clepsydra*. The type of Mr. F. Sanborn's figure is in his collection of nymphæ.

A full-grown nympha from Sitka, Northwest America, belongs perhaps to *Æ. Sitkensis*. It is in bad condition, and is not figured, but is very similar to the above described species.

Three nymphæ, supposed to belong to *Æ. clepsydra*, multicolor, and verticalis were not described. None of them is full-grown.

18. *ÆSCHNA* SPECIES.

PLATE IV. FIG. 2.

Nymphæ, male and female, very young to full-grown. Length, 20 to 57 mm.; breadth, 9 mm. Locality, Himalaya, Kooloo, and Sutledge River. Rev. M. Carleton.

Similar to *Anax*; very long, but narrow; third joint of antennæ about as long as the following together. Processes blunt; posterior more prominent, enclosing an obtuse angle. Mask cleft wider, with a small black tooth on each side. Palpus cut straight at tip, upper angle sharp and lower one produced in a strong end hook; below finely denticulated. Lateral appendages half length of middle one, abruptly pointed. Male projection shorter, conical. Femora with dark rings before the lower end. Abdomen not very thickly covered with spinous tubercles.

No imago arrived with the nymphæ. Perhaps they belong to *Æ. ampla*.

19. *ÆSCHNA* SPECIES.

PLATE I. FIG. 4.

Nine nymphæ, male and female, full-grown and young, in alcohol. Length, 45 mm.; breadth, 9 mm. Locality, Rio San Francisco and Carundahi, Rio do Macacos; one male nympha from Reajinha River, near Corpo de Pompo, from Barbacena, Brazil. Thayer Expedition. Coll. Mus. Comp. Zool.

Full-grown and very young nymphæ, 24 mm. long, wing cases not reaching first segment, from Rio do Macacos have appendices similar.

Similar to *Æschna grandis*, stout. Eyes rather large and more prominent. Ocelli indicated. Antennæ very small; third joint length of first and second and twice as long as fourth. Occiput more deeply notched than in *Æschna grandis*, sides oblique and rounded; a number of polished bands on upper side. Mask extending to between middle legs, broad, enlarged gradually forward, middle third produced in short rounded lobe, cleft, sides of cleft rounded and on each side a small black tooth. Palpus denticulated, cut straight at tip, movable hook extending somewhat beyond base of opposite one. Prothorax rounded behind; processes sharp, enclosing rather less than right angle; posterior nearly twice as long as anterior, and bent backward. Legs rather slender; femora arcuated; hind legs reaching middle of seventh segment. Abdomen broad and stout, rather densely covered with short yellow hairs; dorsal band dark; lateral spines on segments six to nine, spine on ninth shorter than tenth segment. Appendages rather longer than two last segments, very sharp; lateral superiors nearly as long as inferiors, very sharp; middle appendage somewhat shorter and deeply notched. Male projection conical, rather sharp, one half length of middle appendage. Female valve reaching somewhat beyond ninth segment. This species cannot be determined. It may perhaps belong to a peculiar group named *Ura-cautha* by De Selys. To this group belong *Æschna Januariæ*, Hagen's *Synopsis*, *Æschna castor*, Brauer, Voyage of Novara, and two or three species not yet described.

Two nymphæ from Rio Janeiro, Brazil, are similar, but belong probably to a related but different species.

BRACHYTRON.

Evans, British Libellulina, p. 22.

Body elongated, lateral spines on segments six to nine, slender. Head rather small and convex; base of occiput narrow. Eyes small in proportion. Front border of mask produced in oblique angle, cleft; no teeth. Legs rather short. Appendages short; middle one somewhat shorter than inferiors, cut at tip; laterals short. Male projection conical; female valve not reaching tip of segment.

20. BRACHYTRON PRATENSE.

PLATE V. FIG. 1.

Mueller. Selys, Revue Odonates, p. 115.

Nympha skins, male and female. Length, 39 to 42 mm.; breadth, 7 mm. Locality, Vienna, raised by F. Brauer; raised by Mr. Heyer, Lueneburg; Danzig, by Mr. Brischke; Silesia from Charpentier.

Long and slender. Head smaller in proportion than in the other European species, narrower behind, more convex on top. Eyes rather small, placed at forward angles of head, prominent, and separated from occiput by very curved line. The inner hind angles much produced and rounded, with no appearance of facets, which occur in other species, with exception of *A. affinis*, which wants facets in inner part of produced angle. Occiput longer, hind angle sharper and notched behind at almost a right angle; a spot on each side of median line, which forms a ridge; polished bands on hind angles. Antennæ differ in having third joint only half length of fourth. Mask reaching middle legs, less gradually enlarged, enlargement being chiefly of apical third; more produced at middle third of front border, cleft narrower, and comb of hairs longer. Palpus shorter, more denticulated; movable hook not meeting the base of opposite one, nearly as thick at base as palpus. Prothorax small, a little wider than hind border of occiput, more rounded behind. Stigmata uncovered. Processes not much separated; anterior nearly twice as long as posterior. Legs shorter and stronger; hind legs reaching middle seventh segment; femora somewhat arcuated. Wing cases reaching fourth segment. Abdomen long, slender, broadest at middle, tapering gradually, smooth; four dark spots in middle of each segment, two on each side, and near margin two polished impressions. Segments nearly equal in length, except tenth, which is somewhat shorter. Lateral spines small on segments six to nine; only indicated on sixth sometimes; that on ninth only one third length of tenth segment. Appendages short and blunt, a little longer than segment ten. Inferiors cut obliquely, middle one nearly as long, cut straight at tip, not notched. Lateral superiors nearly as long as middle one, bluntly pointed. Male projection large, conical, rounded, more than half length of lateral superiors. Male genitals slightly marked; female valve not reaching end of ninth segment.

The nymphæ described are the types of Hagen, Stett. Zeit. XIV. p. 268, and Brauer, Neur. Austr. p. xvi.

GOMPHÆSCHNA.

Body long, slender. Eyes rather small. Abdomen with lateral spines on segments five to nine. Appendages as long as segments nine and ten. Inferiors sharp, bent inward at tip; middle one a little shorter, sharp, bifid at tip; laterals very short.

21. GOMPHÆSCHNA FURCILLATA.

PLATE II. FIG. 4.

Say. Hagen, Synopsis N. Amer. Neur. p. 131.

Two nymphæ, in alcohol, young. Length, 26 mm.; breadth, 7 mm. (figured). Locality, Berkshire, Tioga County, N. Y. Three male and female, length, 32 mm., Brookline, Mass. Two young, Cambridge, Mass.

Head large, very broad; length less than half the breadth, much narrower behind. Eyes very prominent, placed at forward angles, separated from occiput by very curved line; space between the eyes wide, convex. Posterior portion formed of three orbicular elevations, showing the indications of ocelli more distinctly than in the other species. Antennæ small, third joint somewhat longer than fourth. Occiput short, rough, with yellowish scales, sides oblique: hind angles somewhat more than a right angle, with parallel polished bands; elliptical polished spot on each side of median line; hind border not very deeply notched. Mask extending to between middle legs gradually enlarged forward. Middle third of front border rather strongly produced; cleft and with a well defined tooth on each side of cleft. Palpus meeting the opposite one, straight on tip, somewhat arcuated; inner border rounded and finely denticulated; end hook small; movable hook bent, reaching base of opposite one. Prothorax narrow, as broad as occiput, somewhat bent up at sides. Processes conical, sharp, enclosing rather less than right angle; posterior somewhat shortest and not so sharp. Legs rather strong, arcuated; hind legs reaching segment eight. Femora and tibiæ ringed twice with darker color. Wing cases reaching somewhat beyond second segment. Abdomen stout in proportion; lateral spines on segments five to nine, strong; that on ninth nearly as long as tenth segment; tenth segment as long as ninth. Appendages stout, sharp, hairy, as long as two last segments. Inferiors pyramidal, sharp, and somewhat curved inward; middle one nearly as long as inferiors, and split at tip; lateral superiors only one fourth as long as middle one, conical. Male projection conical, as long as

the lateral superiors. Male genitals indicated; female valve, in young specimen, — only 19 mm. long, — only half length of segment, hairy. There is a smaller specimen in the coll., Lee Co., Virginia, length, 12 mm., similar; differs only in having the middle appendix more widely split and the inferiors more curved.

This determination is only conjectural, as *G. furcillata* has in imago the inferior appendage of the male bifid. Perhaps the young nympha from Virginia may belong to *G. antilope*.

NEURÆSCENA.

Similar to *Gomphæschna*; movable hook, extending somewhat beyond the base of opposite one. Lateral spines on segments four to nine. Appendages long, middle one as long as inferiors, pointed; laterals very short.

22. NEURÆSCHNA VINOSA (QUADRIGUTTATA).

PLATE II. FIG. 3.

Say. Hagen, Synopsis N. Amer. Neur. p. 130.

Nine nymphæ, male and female, full-grown and young, in alcohol. Length, 35 mm.; breadth, 7 mm. Coll. Mus. Comp. Zool. Locality, Berkshire, Tioga County, N. Y., from M. A. Mayer; Kentucky; Vermont; Lake Superior.

Long, rather stout. Head as in *G. furcillata*. Eyes broader at inner hind angles. Vertex shorter. Ocelli hardly indicated; antennæ, third joint, as long as first and second; fourth more than half length of third; following as long as fourth. Occiput notched behind. Sides oblique; parallel bands on hind angles less defined. Mask extending to between middle legs, rather narrow in proportion; middle third of front border produced in obtuse lobe, cleft, and with a black tooth on each side of cleft. Palpus similar to *G. furcillata*. Movable hook reaching somewhat beyond base of opposite one. Prothorax similar. Processes more prominent, sharp, of nearly equal length, and enclosing rather less than right angle. Anterior more compressed. Legs rather stouter in proportion, short; hind legs reaching seventh segment. Femora and tibiæ with two paler rings. Tarsal joint darker at end. Wing cases reaching somewhat beyond middle of fourth segment. Abdomen straighter at side than *Gomphæschna*; rough, marked with eight polished impressions. Lateral spines on segments four to nine, those on ninth nearly

as long as tenth segment. Appendages a little shorter than segments nine and ten, stout, sharp, hairy. Inferiors triangular, sharp; middle appendage as long as inferiors, sharp-pointed at tip; lateral superiors short, less than one fourth length of middle one, blunt. Male projection as long as laterals, triangular, rounded at tip; female valve half length of ninth segment, hairy at edges. There is a young specimen in the collection, 17 mm., similar.

The above determination is conjectural. There is no other described species to which this nympha seems so likely to belong.

EPIÆSCHNA.

Hagen Ms's.

Body stout; antennæ six-jointed, sixth longest, pointed. Prothorax bluntly pointed at outside angles. Legs short in proportion. Abdomen large and broad. Segments four to nine have lateral spines; inferior and middle appendages of equal length.

23. EPIÆSCHNA HEROS.

PLATE I. FIG. 3.

Æschna Heros. Fab. Ent. Pyst. Suppl. p. 285.

Hagen. Synopsis, N. Amer. Neur. p. 128.

Nymph. female, full-grown, alcoholic. Length, 50 mm.; breadth, 12 mm. Locality unknown; Portland, Me., raised by Mr. Jones. Coll. Comp. Zool.

Stout, head large, rather convex, slopes rather abruptly at sides. Eyes rather small in proportion, prominent, egg-shaped, prolonged obliquely at inner angles in a lobe which is divided by a linear impression, in front of which the lobe has indication of facets, behind rough and like occiput, except just along border of occiput, where it is obscurely faceted. Vertex rather above the level of the eyes; indications of ocelli on posterior portion; anterior semicircular, small. Antennæ six-jointed, small; third longer than fourth, somewhat shorter than the two basals; sixth longer than fourth and fifth together, pointed. Occiput longer than fore part of head, roughened by small black tubercles; sides oblique, rounded at hind angles; rather deeply notched behind; polished bands on hind angles. Mask extending to middle legs, enlarged at apical half rather abruptly, and rounded at sides; middle third of foreborder not much produced, cleft rather shallow, on each side of cleft a small tooth. Palpus meeting opposite one, cut straight at end,

lower angle produced in tooth, inner edge sharply denticulated. Movable hook, sharp, bent, not reaching base of opposite one. Prothorax small, nearly as broad as occiput, notched in front; median line impressed; front angles produced and sharp. Processes long, not very sharp, not much separated; posterior half as long as anterior. Legs short, of nearly equal length, middle legs reaching seventh segment; femora somewhat bent. Wing cases nearly reaching segment five. Abdomen broad, rounded above, rounded off rather abruptly beyond seventh segment. Segments of nearly equal length, tenth shortest, almost smooth; eight dark points on each segment, as in *B. pratense*. Stout lateral spines on segments four to nine, smaller on four and five. Lateral spines on segment nine only one third length of tenth segment. Appendages stout, blunt, one half longer than segment ten. Lateral inferiors cut obliquely at tip; middle a little notched on tip, as long as inferiors. Lateral superiors cylindrical, strong, pointed, somewhat more than half the length of middle one. Female valve reaching end of segment nine. There are in the collection two very young male nymphæ, 20 and 22 mm. long; one from Medford, Mass., very similar to the above. Antennæ six-jointed, male projection just indicated, conical, flat, shorter than the lateral superiors.

The peculiar features, unusual number of joints of antennæ, etc., were strong reasons for referring it, even before it was raised, to *Epiæschna Heros*, which is the largest species living in this country, and differs so considerably from other species as to form a new genus.

ÆSCHNA??

Body stout, short, broad in proportion. Head, length half breadth. Eyes in anterior half of head, orbicular, rather small comparatively, prolonged at inner hind angles in narrow and extended lobe. Antennæ five (?), jointed, third joint longest. Mask extending to middle legs, small comparatively; middle third of foreborder produced in cleft lobe, having a small tooth on each side of cleft. Legs strong, long, and with femora notched at upper edge; hind legs more separated at base than fore and middle legs. Abdomen rather rounded behind than tapering; tenth segment enclosed in ninth. Dorsal spines on all segments; lateral spines on all segments. Appendages short, stout, sharp on tip; middle one blunt. Male projection conical.

This curious nympha was referred to *Gomphina* at first, with which it

agrees more in general appearance than with *Æschmina*. The three-jointed tarsi and slender antennae are like *Cordulegaster*, from which, however, it is excluded by the different shape of mask, which again refers it to *Æschmina*.

A nearer determination is now not possible.

24. *ÆSCHINA* (?) SPECIES.

PLATE III. FIG. 3.


Male nympha, dried. Length, 37 mm.; breadth, 13 mm. Locality, La Guayra, Venezuela.

Stout, short, broad, somewhat depressed, in general looking like some forms of *Gomphus*. Head, body, and legs covered with cylindrical or pyramidal spines, dark at the base and light-yellow at upper portion, which has a slight depression, in which there may be an orifice. The spines point backward, and vary in size. Head large; breadth, twice length. Eyes prominent, prolonged at the inner angle into a triangular lobe, which in the specimen is wrinkled so as to have appearance of vanes of a feather. Situated in front of this is a polished ridge lighter in color. Upper lip very large. Vertex rather wide and slightly convex; posterior portion showing indications of ocelli. Antennae rather short, five-jointed; two basal joints globular; third as long as both basals, cylindrical; fourth shorter than fifth, which is also shorter than third and pointed. It is possible there may be two more rudimentary joints; but from the single specimen, which has only one of the antennae, it is impossible to decide. Occiput rather deeply notched, sides oblique, hind angles rounded. Just back of eyes is a small conical projection near each side. Mask small in proportion, extending between middle legs, narrow and gradually enlarged; middle third of foreborder somewhat produced, and cleft surmounted by comb of light hairs, and with a small tooth on each side of cleft. Palpus nearly if not quite smooth. Movable hooks, slender, and bent near tip. Prothorax as broad as occiput; cut rather square at sides, and projected in rather sharp angle at forward part. Stigmata entirely covered. Thorax rather long in proportion. Legs strong, formed for running; fore and middle legs of nearly equal length; hind legs longer, reaching ninth segment. The femora of all the legs are stouter and shorter than the tibiae, and have a notched hook on the upper edge; tarsi three-jointed. Processes above fore legs of equal length; posterior one triangular, sharp; anterior large, square, notched at tip. Abdomen stout,

broader than the thorax, and rather rounded behind than tapering. The segments are produced at the sides in modified spines, giving the general appearance of the abdomen's being notched and slightly turned up at the edges. The segments have dorsal spines more developed on the third and fourth segments than on the following. Segments of nearly equal length; ninth longest at sides, and enclosing the tenth, which is much shorter. Appendages short. Middle appendage pyramidal, two thirds as long as inferiors, blunt. Male projection one half length of middle appendage. Inferiors three-cornered, length somewhat greater than breadth. Lateral superiors less than half as long as middle one, very small, blunt, pyramidal. Genital parts are visible on second and third ventral segments and valves on ninth.

It is very difficult to place this nympha. The mask is like *Æschna* in the produced and cleft middle third of foreborder. The antennæ have two joints less than *Æschna* (?) as far as can be seen. The legs have three-jointed tarsi, as in *Æschna*, but are not equally distant at base, as is the case in all known forms of *Æschna*, while the notched appearance of upper edge of femora is wholly peculiar. The processes above front legs are *Æschna*-like, there being no such formation in *Gomphina*. The form of abdomen is more like *Gomphus* than *Æschna*, as are the dorsal hooks; the sides of abdomen are more like *Gomphus* and *Hagenius*, as is also the enclosed tenth segment. The appendages are peculiar, but more like *Gomphus* than *Æschna*. The entirely closed stigmata are again unlike *Æschnina*. The imago is perhaps still unknown. It may belong to *Staurophlebia*, but is certainly not *Æschna* or *Anax*. The large size of upper lip would suggest *Zonophora* among *Gomphina*, but *Cordulegaster* is the only known genus of *Gomphina* having three-jointed tarsi, and the form of the mask is quite unlike that of *Cordulegaster*. The marked and peculiar sculpture is unlike *Æschnina* and *Gomphina*, unless in some species of North American *Æschnina*, which show a somewhat similar sculpture. Altogether, the nympha is the most peculiar observed.

A second specimen of this nympha, in the collection of Professor Rosenhauer, in Erlangen, Bavaria, is stated to be received from Chili. Perhaps the aberrant nympha may belong to a species of the aberrant genus *Petalia*.



SYNOPSIS OF SPECIES DESCRIBED.

SUBFAMILY ÆSCHNINA.

DIVISION I. — ANTENNÆ SEVEN-JOINTED.

A. *Head longer than broad; mask extending through hind legs; front border of mask in the middle with two sharp triangular lobes.*

Gynacantha Ramb. (Supposition.)

Body very slender, flattened; head long; eyes small, not very prominent, slightly produced at inner hind angles; occiput sharply prolonged behind on each side; mask long, narrow; lateral spines on segments six to nine.

1. **G. spec.**

Nympha, full-grown. Brazil, South Carolina, and Florida.

Characters of the genus: appendages sharp; the middle one as long as the inferiors, cut at tip, slightly notched; laterals a little shorter; male projection conical.

B. *Head broader than long.*

Anax Leach.

The giants of the family. Body long, stout; eyes large, more prominent in the middle of the flat head; mask extending beyond middle legs; front border produced in a short rounded lobe, cleft in middle; legs slender; abdomen large; lateral spines on segments seven to nine; appendages strong, sharp, the middle one shorter, notched; laterals half as long; male projection short, cut at tip; female valves shorter than the segment.

2. **A. Formosus Vander Linden.** (Raised.)

Europe; full-grown nymphæ.

The two processes above first legs short, broad at tip; in right angle, the posterior larger; both enclosing a right angle; lateral appendages sharply pointed; male projection cut straight.

3. A. Mauricianus Rambur. (Supposition.)

Nympha, full-grown, male and female, Mauritius Isl.; Zanzibar.

Similar to *A. formosus*; processes blunt, enclosing an obtuse angle; mask more suddenly enlarged; middle appendage nearly as long as inferiors.

4. A. Junius Drury. (Raised.)

Nymphae, full-grown and young, male and female. United States.

Processes similar to *A. formosus*, more obtuse at tip, equal in size; lateral appendages cylindrical, pointed suddenly at tip; male projection notched at tip.

5. A. Julius Brauer. (Supposition.)

Nympha, male and female. Japan.

Processes and lateral appendages similar to *A. formosus*; male projection as in *A. Junius*.

6. A. Guttatus Burm. (Supposition.)

Nymphae, full-grown, male and female. Marshall Isl., Pacific Ocean.

Body more elongated; processes unequal, anterior blunt, posterior smaller, angular behind; enclosing a very obtuse angle, with small median notch; male projection narrow at tip, cut square.

7. A. Amazili Burm. (Supposition.)

Nymphae, full-grown, male and female. Jamaica; Central America.

Similar to *A. Junius*, but more elongated; processes of equal length and size, enclosing a very obtuse angle; lateral appendages sharply pointed; male projection shorter; notched at tip.

8. A. spec.

Nymphae, male and female. East India.

Similar to *A. Junius*; processes more blunt, posterior smaller, enclosing an obtuse angle; lateral appendages abruptly pointed; male projection notched at tip.

Æschna Fabr.

Body, long, stout; head large; eyes very prominent at the forecorner; mask extending to middle legs, broad, front border slightly produced in round or oblique lobe, cleft; legs longer, stouter; abdomen broader; lateral spines on segments six to nine, or five to nine, or four to nine; appendages long, sharp; male projection conical; female valves reaching tip of segment.

A. Lateral spines on sixth to ninth segments; no teeth on front border of mask; lateral appendages shorter than the inferiors; middle one notched at tip.

9. Æ. Rufescens Vander Lind. (Raised.)

Nymphae, full-grown, male and female. Europe.

Hind angles of head rounded; mask large, gradually enlarged; processes enclosing an

acute angle, triangular, sharp, the anterior half length of the posterior; lateral appendages only one fourth shorter than middle one.

10. *Æ. Grandis* Linn. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Hind angles of head more oblique; mask similar to *Æ. rufescens*; processes longer, more separated, of equal length, sharp, bent a little outwards, the anterior rather slender; lateral appendages about half the length of the middle one.

11. *Æ. Cyanea* Muell. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Hind angles of head oblique; basal half of mask, narrower; processes short, about of equal length, enclosing a right angle; posterior larger, rounded in sides; lateral appendages half length of middle one.

12. *Æ. Juncea* Linn. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Similar to *Æ. cyanea*; anterior process half length of posterior, much smaller.

13. *Æ. Viridis* Eversm. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Similar to *Æ. rufescens*; hind angles of head more oblique; processes of equal length, sharp tips a little bent outwards, enclosing right angle; lateral appendages half length of middle one.

14. *Æ. Mixta* Latr. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Similar to *Æ. cyanea*; hind angles of head rounded; mask gradually enlarged at apical third; processes blunt, stout, equal in length, anterior narrower; not much separated; lateral appendages about half length of middle one.

15. *Æ. Affinis* Vander Lind. (Raised.)

Nymphæ, full-grown, male. Europe.

Similar to *Æ. mixta*, more slender; processes blunt, very slightly separated, of equal length and size; lateral appendages shorter.

B. Lateral spines on fifth to ninth segments; no teeth on front border of mask; lateral appendages shorter than the inferiors; middle one notched at tip.

16. *Æ. Eremitica* Scudder. (Supposition.)

Nymphæ, full-grown, male and female. Arctic America and New Hampshire.

Similar to *Æ. rufescens*; hind angles of head oblique; processes long, equal, sharp, tips bent a little outwards, enclosing less than right angle; lateral appendages two thirds length of middle one; female valves not quite reaching tip of segment.

17. *Æ. Constricta* Say. (Raised.)

Nymphæ, full-grown, male and female. United States.

Similar to *Æ. eremitica*, more slender; apical half of mask less enlarged; processes similar, less sharp, the anterior less bent outwards, the posterior a little larger; lateral spine of segment fifth generally only indicated; lateral appendages longer than half the middle one.

C. Lateral spines on sixth to ninth segments; front border of mask with two small teeth; tip of middle appendage notched.

18. *Æschna* spec.

Nymphæ full-grown, male and female. Himalaya.

Similar to *Anax*, very long; processes blunt, posterior more prominent, enclosing obtuse angle; lateral appendages half length of middle one, abruptly pointed.

19. *Æschna* spec.

Nymphæ, full-grown, male and female. Brazil.

Stout, shape of *Æ. grandis*; processes prominent, sharp, the posterior longer, bent outwards; enclosing right angle; lateral appendages very sharp, length of the inferiors.

***Brachytron* Evans.**

Body long, slender; head smaller, rather convex above, much narrower behind; front border of mask produced in oblique angle, cleft; legs shorter; abdomen long, narrow; appendages short, middle one a little shorter than inferiors, cut at tip; laterals shorter; male projection conical; female valves not reaching tip of segment.

20. *B. Pratense* Mueller. (Raised.)

Nymphæ, full-grown, male and female. Europe.

Processes slightly separated, sharp, the posterior half length of the anterior; lateral spines on sixth to ninth segments.

***Gomphæschna* Selys.**

Body long, slender; eyes small; lateral spines on segments five to nine; middle appendage a little shorter than the inferiors, bifid at tip.

21. *G. Furcillata* Say. (Supposition.)

Nymphæ, male and female, nearly full-grown. United States.

Characters of genus; processes conical, sharp, posterior shorter, enclosing less than right angle.

***Neuræschna* Selys.**

Similar to *Gomphæschna*; lateral spines on segments four to nine; appendages long, the middle one as long as the inferiors; all sharply pointed.

22. N. Vinosa Say. (Supposition.)

Nymphæ, full-grown, male and female. United States.

Similar to *G. furellata*. Characters of genus: processes prominent, sharp, of equal length, enclosing less than right angle; body rough.

DIVISION II. — ANTENNÆ SIX-JOINTED.

Epiæschna Selys.

Similar to *Æschna*; antennæ six-jointed; lateral spines on segments four to nine; inferior and middle appendages of equal length.

23. E. Heros Fabr. (Raised.)

Nymphæ, full-grown, male and female. United States.

Characters of genus: processes long, the posterior half as long as anterior, not much separated; middle appendage a little notched at tip.

DIVISION III. — ANTENNÆ FIVE-JOINTED.

Æschna??**24. Species Nova.**

Nympha, male. La Guayra, Venezuela, and Chili.

Genus and species very abnormal. Cf. the description. Perhaps belonging to *Gomphina* and to genus *Petalia*.

EXPLANATION OF THE PLATES.



[*a*, sideview ; *b*, labium ; *c*, appendages ; *d*, processes above the base of first pair of legs ;
e, front view of the head.]

PLATE I.

- FIG. 1. *Anax formosus*.
" 2. *Anax Junius*.
" 3. *Epiæschna heros*.
" 4. *Æschna* species. No. 19.

PLATE II.

- FIG. 1. *Æschna grandis*.
" 2. *Æschna eremitica*.
" 3. *Neuræschna vinosa*.
" 4. *Gomphæschna furcillata*.

PLATE III.


- FIG. 1. *Æschna constricta*.
" 2. *Gynacantha* species. No. 1.
" 3. *Species*. No. 24.

PLATE IV.

- FIG. 1. *Æschna juncea*.
" 2. *Æschna* species. No. 18.
" 3. *Æschna cyanea*.


PLATE V.

- FIG. 1. *Brachytron pratense*.
" 2. *Æschna mixta*.
" 3. *Æschna affinis*.
" 4. *Æschna rufescens*.
" 5. *Æschna viridis*.



L. Cass. del.

1811. 17





Family Acanthina










PLATE 100





Family Aculeata





Memoirs of the Museum of Comparative Zoölogy

AT HARVARD COLLEGE.

VOL. VIII. No. 2.

EXPLORATION

OF THE

SURFACE FAUNA OF THE GULF STREAM

UNDER THE AUSPICES OF THE COAST SURVEY.

By ALEXANDER AGASSIZ.

III. PART I.

The Porpitidae and Vevellidae. By ALEXANDER AGASSIZ.

PUBLISHED BY PERMISSION OF CARLILE P. PATTERSON AND J. E. HILGARD,
SUPERINTENDENTS U. S. COAST AND GEODETIC SURVEY.

WITH TWELVE PLATES.

CAMBRIDGE:

Printed for the Museum.

JULY, 1883.



EXPLORATION
OF THE
SURFACE FAUNA OF THE GULF STREAM

UNDER THE AUSPICES OF THE COAST SURVEY.

By ALEXANDER AGASSIZ.

III. PART I.*

The Porpitiidae and Velellidae. By ALEXANDER AGASSIZ.

(Published by permission of CARLILE P. PATTERSON and J. E. HILGARD, Supts. U. S.
Coast and Geodetic Survey.)

WHILE at the Tortugas † examining the structure of the coral reefs, I took advantage of my opportunities to study the surface Fauna of the Gulf Stream, and when not otherwise occupied devoted the time I could spare to complete the notes and drawings I had accumulated regarding *Porpita* and *Velella* under less favorable circumstances at other points of Florida, at Newport, and on board of the "Blake." These notes are now published, as giving the principal points on the Natural History of a small and limited group of Oceanic Hydroids, interesting from their affinities, on the one side, to the Tubularians, with which Vogt, Kölliker, and Agassiz were inclined to associate them, and, on the other, with the Siphonophoræ proper, with which, as

* MR. C. O. WHITMAN was sent to Key West this spring in hopes of obtaining the material necessary to complete this memoir, and at the same time to investigate anew the whole subject of the structure and functions of the so-called yellow cells. Although Mr. Whitman spent six weeks at Key West, he was unable to accomplish the object of his trip, not a single *Velella* appearing in the harbor of Key West during the whole of his visit. I have therefore thought it advisable not to delay the publication of the descriptive part of this memoir any longer, and to complete it when the necessary preparations could be finished.

† See Letter No. 5, ALEXANDER AGASSIZ to CARLILE P. PATTERSON, on the explorations in the vicinity of the Tortugas in 1881. Bull. M. C. Z., VIII., No. 3, p. 145. I spent the months of March and April, 1881, at Key West and at the Tortugas, under the auspices of the United States Coast Survey, the late Mr. Patterson, the Superintendent, having kindly placed at my command a steam launch while engaged in examining the distribution of corals and studying the surface fauna of the Gulf Stream. The Hon. Secretary of the Navy kindly allowed the commanding officer at Key West, Lieutenant Winn, to give me permission to occupy the loft of the Navy storehouse as a laboratory.

will be seen, they have much less in common. This group of Hydrozoa is eminently characteristic of the Gulf Stream, and wherever its influence extends there Porpita, Verella, and Physalia have been found. In fact these surface animals are excellent guides to the course of the current of the Gulf Stream, — natural current bottles, as it were. They are thrown up along the whole length of the Atlantic Coast of the United States, from the Straits of Florida to the south shores of Cape Cod and of Nantucket. Physalia, Verella, and Porpita are occasionally driven into Narragansett Bay; the former is an annual visitant, the latter has only been found once, in 1875, and Verella has come into Newport harbor during three summers. It is undoubtedly also to the action of the Gulf Stream that we must ascribe the presence of the few species of Siphonophoræ which appear on the southern coast of New England towards the middle and last of September, such as Eudoxia, Epibulia, and Dyplophysa, which are all found at the Tortugas. On the contrary, Agalma and Nanomya are northern visitants at Newport, brought down by the arctic shore current from the northern side of Cape Cod, Agalma being common at Eastport. Other species of our southern New England free Hydroids, such as Eutima, Trachynema, Eucheilota, Liriope, Zanelea, and many other species which have been described by McCrady, from Charleston, S. C., are also brought north every year along the course of the Gulf Stream, and during the summer are blown to the westward towards the New England coast and the Atlantic coast of the Middle States by the prevailing south-westerly winds.

Verella mutica Bosc.

The Florida species of *Verella* occasionally finds its way north as far as Newport and Nantucket; it is found in great numbers in the Straits of Florida, between Cuba and the Florida reefs. Thousands of them are brought in by favorable winds and tides into Key West harbor, and are carried by the same agencies between the Tortugas channels. They are usually seen in large schools, and, although capable of considerable independent movement, by means of their tentacles, in a smooth sea, yet they are practically at the mercy of the winds and currents. They are destroyed in great numbers by even moderate waves, which, upsetting them, drive them ashore, or kill them, if they are kept keel downward for any length of time. They apparently need a considerable amount of movement, for when kept in confinement they do not thrive, soon die, and are rapidly decomposed. The dead

floats are thrown upon the beach behind Fort Jefferson at the Tortugas in great numbers, forming regular windrows, and, when dry, are blown by the winds to the highest parts of the beach.

The Florida species is much larger than the Mediterranean *V. spirans*. Specimens measuring nearly four inches in length are not uncommon. On Plate I is figured in profile, from above and below, a huge *Velella*, nearly five inches in length. This is a somewhat unusual size. The outline of the mantle, seen from above, is less elliptical than in *V. spirans*, it is somewhat rectangular, with rounded corners (Pl. I, Fig. 2), and is also proportionally broader than in the Mediterranean species. Seen from above (Pl. I, Fig. 2), the color of the mantle is of a metallic bluish green, with a deep cobalt blue edge surrounding the outer edge of the float, and a similar band, forming an irregular ellipse with re-entering sides, placed somewhat diagonally across the float. Between these bands the color of the mantle passes rapidly from a yellowish green to the dark-blue inner and outer bands. Through the outer edge of the mantle the base of the outer blue tentacles of the lower side of the float can be indistinctly seen. The whole of the mantle is dotted with the patches of the so-called liver-cells, of a brownish color. The extreme edge of the dark outer part of the mantle is fringed with a light cobalt blue band, in which are placed the glandular organs of the free edge of the mantle. The free outer edge of the mantle is usually turned down so as to form slight indentations, or apparently sharp incisions in the general outline (Pl. I, Figs. 2, 3). The figure from below (Fig. 3) shows how the edge of the mantle is carried when folded under to produce the incisions seen from the upper side. The mantle, where it covers the central part of the float, is of a light greenish blue, with a metallic lustre, with a few patches of liver-cells, diminishing in number towards the base of the keel. The greenish lines of color form concentric lines parallel with the chambers of the float, crossed by triangular radiating rays extending from the fixed edge of the mantle towards the base of the keel, dividing the float into irregular alternating sections of light and colored triangular spaces. The keel is of a delicate steel color, with a thickened edge of the mantle (Pl. I, Fig. 2) running round it. In this the patches of liver-cells are closely packed together, and form dark-violet triangular patches, extending at right angles from the edge of the keel to the edge of the mantle. Seen from above the float is divided longitudinally by a long triangular band of liver-cells, which are seen through the float, so as to divide the float into two

nearly equal parts (Pl. I, Fig. 1). Seen from below (Pl. I, Fig. 3), the mantle is of a lighter bluish green color, with a light blue edge (the marginal glands), followed by a somewhat darker belt passing into the greenish color of the mantle.

The tentacles (the closed prehensile polypites) are long, slender, of a bluish color, forming a double row round the outer edge of the float. The longest only of these tentacles extend beyond the free edge of the mantle so as to be seen projecting beyond it, when the *Velella* is viewed from above. These tentacular polypites taper very gradually; they seem capable of but slight expansion and contraction, and are quite sluggish compared with the smaller, active, feeding and reproductive polypites. These are arranged in five or six rows between the rows of tentacular polypites and the large central polypite (Pl. I, Fig. 3); the large, blue, prehensile, closed tentacular polypites are covered at the base (Pl. VI, Fig. 17) by elliptical or circular patches of lasso-cells (Pl. VI, Fig. 16), which about half-way towards the extremity become more crowded, and unite so as to form a band of lasso-cells on each side of the polypite (Pl. VI, Figs. 14, 15). In some young polypites the bands alone exist; while in others the elliptical patches alone are found (Pl. VI, Fig. 18). The smaller, the feeding and reproductive polypites are most active, and capable of great expansion and contraction. They are covered towards the upper extremity with elliptical knobs of lasso-cells (Pl. II, Figs. 5, 6, 7), the edge of the open extremity of the polypite forming ten to twelve indistinct lobes. At the base of the polypite there are, according to its size, from five to eight clusters of *Medusæ* buds, in different stages of development (Pl. II, Figs. 1, 5, 7). While the large central polypite is the main feeding mouth, the smaller lateral ones also perform, to a limited extent, the functions of feeding polypites; but, being all connected at their bases with the general vascular system, the fluids they take in enter at once into the general circulation. Both the central polypite, as well as the smaller lateral polypites, eject the digested substances which have gone through the general circulation.

As has been shown by Weissmann, the circulation of the fluids in the *cenosare* in fixed *Hydroids* is kept up mainly by the muscular contraction of the walls, or by the action of the *ciliæ* lining the cavities. A similar condition exists in the canals forming the vascular system of the float of *Porpita* and of *Velella*, and in the polypites, where the fluids are rapidly propelled by the action of *ciliæ* lining the inner walls.

The large, exterior, marginal, prehensile tentacles (Pl. VI, Fig. 14) are edged along the extremity with a band of lasso-cells, composed of large circular cells, closely packed together (Pl. VI, Fig. 15). Towards the base these bands of lasso-cells become disconnected (Pl. VI, Fig. 17), forming irregularly shaped disconnected circular patches (Pl. VI, Fig. 16). In young tentacles these bands of lasso-cells are not clearly defined, and when they first make their appearance they appear as patches near the base, gradually extending towards the extremity, there to form the connected bands of the older tentacles. The large prehensile tentacles, the feeding and reproductive polypites, are all attached to the lower side of the space occupied by the float, and to the part of the mantle immediately adjoining it. Huxley considers the tentacles of *Veleva*, as well as those of the *Porpitiidæ*, as identical with those of the *Hydridæ Sertularidæ*. The latter undoubtedly are to be so considered; but the structure of the tentacles of *Veleva* clearly shows that they are embryonic tentacles, analogous to the "Fangfäden" of *Physalia*, and of other Siphonophores, in which the lasso-cells are arranged in the most simple form, as bands along the edge; while in *Physalia* they form the peculiar well-known reniform appendages paved with large lasso-cells. It is difficult in *Veleva* and *Porpita* to distinguish the young polypite from the hydrocytes of other Siphonophores, and they do not appear to be present in those genera.

The small *Medusæ* buds already contain the peculiar yellow cells so characteristic of the free *Medusæ*. Those which I have raised from *Veleva* (Pl. II, Figs. 15, 16) differ somewhat from that figured by Professor Agassiz (*Contributions to the Nat. Hist. of the U. S.*, Vol. III, p. 53). On becoming free, the young *Medusa* (Pl. II, Fig. 10) has two rudimentary tentacles, one of which (t') is somewhat longer than the other (t), which is in this stage a mere tentacular knob. The close resemblance of the *Medusa* at this stage with such Tubularian *Medusæ* as *Esuphysa*, and *Ectopleura* is very striking. It has, like them, a row of large lasso-cells extending from the base of the tentacles (Pl. II, Fig. 14) to the abactinal pole (Pl. II, Figs. 10, 11, 13). The yellow cells are arranged in clusters along the sides of the four broad chymiferous tubes (Pl. II, Figs. 10, 11), as well as on the surface of the short, rounded, conical, rudimentary proboscis. The *Medusa* of *Veleva* figured by Gegenbaur* has eight chymiferous tubes and one large tentacle; that figured by Vogt,† on the contrary, has only four chymiferous tubes, but

* *Zeits. f. Wiss. Zool.* VIII., Pl. VII, Fig. 10. 1856.

† *Mém. Inst. Nat. Gènevois*, I. 1853, Pl. II.

eight rows of yellow cells, two on each side of the broad tubes, which may have misled Gegenbaur. The Medusæ buds figured by Kölliker* agree with the younger stages as figured by Vogt. The Medusa, when it first becomes free, is elongate, somewhat conical at the abactinal pole. After a couple of days the outline becomes flattened and more hemispherical (Pl. II, Fig. 11). The young Medusæ move with considerable activity by sudden jerks, like some of the Tubularian Medusæ. The tentacles did not increase in length during the time they were kept in confinement (ten days); nor did I fish up any others more advanced than those here figured (Pl. II, Fig. 11) during my stay at the Tortugas.

Kölliker has given an excellent account of the course of the so-called liver system in the Mediterranean species. I have been able to trace, as he has done, its ramifications through the mantle, over its free surface (Pl. IV, Fig. 14; Pl. V, Fig. 8), extending beyond the float (Pl. IV, Figs. 8, 11), to the edge, as well as the ramifications extending over the float and the surface of the keel (Pl. V, Figs. 1, 2, 4, 5). In addition to the two main branches of the system extending round the edge of the keel from the base of the float (Pl. V, Figs. 4, 5, *v.*), there are two other large branches, which run across the float in the deep groove (Pl. III, Figs. 3, 14, 15, 17. *f.*), running obliquely across it. These two branches run up on each face of the keel (Pl. IV, Fig. 15. *v.*; Pl. V, Figs. 1, 2. *v.*) from the float, and then anastomose with the main branches described by Kölliker. All along their course, from the fixed edge of the mantle to the main branch running about parallel to the edge of the keel, the main branches give off a system of meshes and branches which cover the whole of the float and keel, and anastomose (Pl. IV, Fig. 5; Pl. V, Figs. 1, 2, 3, 4) with those extending over the mantle from the fixed edge of the mantle to its periphery. The secondary branches, forming the free edge of the mantle of the keel, which are given off from the main marginal branch, send off short simple secondary branches at right angles to the primaries, thus forming a sort of frill. The extremity of these tubes, again, is connected by a small marginal canal (Pl. V, Figs. 4, 5). The so-called liver † is suspended from the lower side of the float, running up into its conical portion (Pl. V, Fig. 11). The main longitudinal branches (Pl. V, Figs. 11, 12, 13. *l.*) give rise to all the finer ramifications which extend through the whole thickness of the mantle. Fluids circulate with great rapidity

* Die Schwimmpolypen von Messina. 1853.

† In *Velella* and *Porpita* the hepatic organ, as has been pointed out by Huxley, occupies the same position with regard to the pneumatocyst which it occupies in *Rhizophysa*.

through the vascular system; and the terminal pouches of the liver are filled with the brown granular mass usually considered to be a true liver. The main tubes send off an endless number of fine ramifications (Pl. IV, Fig. 14), which assume all possible shapes from that of a flat, angular pouch (Pl. IV, Figs. 11, 12; Pl. V, Fig. 10), to an elongate, many-pointed star (Pl. IV, Fig. 15), or to a tube bristling with fine projections (Pl. IV, Fig. 14; Pl. V, Fig. 8), which become lost in the thickness of the mantle. The ramifications on the lower side of the mantle communicate with the reproductive individuals, as has already been seen by Vogt.

The central polypite is large, whitish (Pl. VI, Figs. 10, 11, 12, 13), with strong interior longitudinal muscular bands, capable of great expansion and contraction. It communicates at its base with the vascular system. Near the aperture of the central polypite we find the extremity covered with patches of small lasso-cells (Pl. VI, Fig. 22), forming near its opening irregular lips. The central opening is specially mobile.

There are but few air-tubes (tubules) starting from the lower surface of the float, and forcing their way through the liver to the base of the reproductive polypites. The majority terminate as a single tube, and they rarely ramify, as is stated to be the case by Krohn, in the Mediterranean species. The air tubes vary greatly in number in different specimens. They take their origin from the lower side of the float, in the five or six chambers nearest the centre (Pl. III, Figs. 18, 19, 20). They generally occur two or three together, sometimes in tufts of four starting close together. They sometimes branch, as has been described by Krohn,* but apparently not as commonly as is the case in the Mediterranean species. The air tubes extend through the liver in a more or less winding course (Pl. III, Figs. 20, 21,) (but much more directly than in *Porpita*), and find their way to the base of a few of the small feeding and reproductive polypites (Pl. VI, Fig. 19). In the only case where I have succeeded in tracing the termination of the air sac, it ended in a blind tube. The air tubes are arranged much like those of *Porpita*, so that their course can be traced on the upper side of the liver when the float is removed and we examine the liver from the upper side. The outer partitions of the interior chambers of the float, where the air tubes take their origin, extend in a series of prongs and processes (Pl. III, Fig. 19) beyond the general surface of the float, so that the rough walls of these inner chambers are in marked contrast to the smooth outer walls of the other chambers of the float.

* Archiv f. Naturg., 1848, I. p. 30.

The mantle covering the float extends, as is well known, not only over the horizontal surface of the float, but also over the sail. It projects beyond that, forming a sort of flap (Pl. V, Fig. 4), much as the mantle projects beyond the horizontal part of the float. From the two extremities of the float, at the base of the keel or sail, there runs along the free edge a large tube of the vascular system (described by Kölliker and Vogt), from which branch the dendritic processes forming the triangular patches (Pl. V., Figs. 4, 5) of the free edge of the mantle of the keel. This free sail mantle is of a light claret color, with a blue edge, and with bluish branching tubes forming the ramifications of the vascular system. These tubes anastomose again at the outer edge, forming an irregular marginal canal. There are no glands to the free edge of the keel mantle (Pl. V, Fig. 5), like those found on the free edge of the horizontal mantle of the float. The yellow cells of the sail mantle are packed principally in patches at the extremities of short tubes opening into the main canal, fringing the keel at the base of the free sail mantle (Pl. IV, Fig. 5, Pl. V, Figs. 2, 10). The dendritic tubes are a series of flattened elliptical pouches, opening into one another, and joined together by frill-like folds of the main tubes (Pl. V, Fig. 7). The two surfaces of the mantle join at the edge of the float, so that the part of the mantle which covers the sail and extends to the outer edge of the float, unites there with that part of the mantle which protects the inner side of the float, and to which the appendages of the lower surface are attached. These two surfaces, thus soldered together, extend some distance beyond the float, forming the free edge of the mantle of the Velella. The mantle itself is slightly contractile, and whenever the Velella is thrown over into any unnatural attitude, or forced on its side, it makes violent attempts by the movement of its prehensile tentacles, aided by movements of the free margin of the mantle, to recover its normal attitude.

Rhizophysa and other Siphonophores are capable of sinking below the surface and swimming back to the surface, but neither Velella nor Porpita appear capable of such movements, a very young Physalia, collected at the Tortugas, intermediate between the stages figured by Huxley (Oceanic Hydrozoa, Pl. X. Figs. 1, 2), was found to swim at various levels in the jar in which it was kept.

All the Vellellæ floats I have examined are left-handed, that is, the sail runs northwest to southeast, the longitudinal axis of the float being placed north and south. I have counted over twenty-five hundred dead floats, thrown on the beaches at the Tortugas, in all of which the position of the float was as stated above.

The young stages of *Velella* differ very materially in appearance from the full-grown *Velella*. In the youngest stage I have had the opportunity to examine, the sail forms a flat elliptical arch (Pl. VI, Fig. 8), extending well beyond the extremities of the conical float, the mantle forming an umbrella-shaped projection, below which extend, when expanded, the eight marginal tentacles and the large central polypite. The rudimentary small feeding polypites are about as numerous as the tentacles, and form an inner ring at their base, much as in the older stages figured here (Pl. VI, Figs. 2, 3). The vascular system extends in straight vessels radiating from the float through the mantle to the outer edge. In a somewhat older stage, seen from the lower side (Pl. VI, Fig. 3), the marginal tentacles and feeding polypites are more numerous, the vessels of the liver system more clearly defined, the sail has become somewhat conical, and the whole float and mantle somewhat flattened. In a still more advanced stage (Pl. VI, Fig. 1), the sail has become more conical, the float greatly flattened, and the canals of the liver system clearly defined, as is well seen in a view from the lower side (Pl. VI, Fig. 2). The central polypite (Pl. VI, Figs. 1, 2 *cp.*) at this stage is most prominent, capable of great expansion; the feeding polypites have greatly increased in number as well as size; there are as yet no signs of the reproductive polypites.

Stuart* has given a detailed development of the Medusa of *Velella*, and traced it directly to a free Medusa with four chymiferous tubes with large masses of yellow cells along the tubes. These Medusæ he kept alive for a few days, but could trace no further stages of development. The oldest Medusa he observed showed as yet no sign of any tentacular appendages. Stuart, however, seemed satisfied that the *Chrysomitra* of Gegenbauer, having sixteen chymiferous tubes and two distinct tentacles with tentacular knobs at the base of the tubes, was really a more advanced stage of the Medusa of *Velella*, though he did not observe the intermediate stages of development between it and the Medusæ he raised directly from *Velella*.

The youngest *Velella* figured by Huxley (*Oceanic Hydrozoa*, Pl. XI, Fig. 9) measured about one tenth of an inch in length. He noticed that the pneumato-cyst did not extend into the crest, in which he also saw the rudimentary canals extending from the limb to the crest. On examining the pneumato-cyst he found a single central vesicle with the first trace of a concentric line, the rudiment of the first concentric chamber. This structure agrees well

* Archiv f. Anat. Phys. u. Wiss. Med. 1870, p. 366.

with that of a young *Verella*, and differs radically from that given by Pagenstecher for *Rataria* (as young *Verella*?), in which the pneumatocyst clearly shows the eight primary divisions so characteristic of *Porpita* with the central opening of the disk. Huxley also observed that in young *Verellae* the hepatic mass does not exist, and is only gradually developed under the pneumatocyst, and that the canals which cross it are mere subdivisions of the somatic cavity produced by the lobes of this organ and their mutual anastomoses. My observations on *Verella* lead me to agree with Kölliker regarding the mode of junction of the liver canals and of the canals on the free edge of the mantle and on the upper part of the float. The dendritic-like structure of the canals of the crest have also been noticed by Huxley in a young *Verella* measuring somewhat less than half an inch in length. The circulation within the canals was most active, and wholly due to ciliary action. The figures given by Huxley (page 126, *Oceanic Hydrozoa*) of the peculiar corrugations and lobes of the lower surface of the float, being taken from alcoholic specimens, are not quite satisfactory representations of their appearance in fresh specimens.

As has already been suggested by Kölliker, Agassiz, and McCrady, the relationship of the *Verellidae* and *Porpitidae*, as well as the *Physalidae*,* to the Tubularian Hydroids is very great. If we compare this group as a whole to the other Siphonophores, the absence of "Deckstücke" and of swimming bells seems to distinguish them from all the other Siphonophores except *Rhizophyza*, which perhaps is only a representative of the embryonic stage of the *Physalidae*, and does not belong into the close association with the other Siphonophores, with which it has usually been placed. What is known of these different families seems to indicate a far closer relationship with the Tubularian Hydroids, such as *Hydractinia*, which may perhaps be the closest ally of the genera named above, and in which the chitinous exten-

* The chambers of the crest of *Physalia* can be considered as a sort of ginder which stiffens the whole float, and to a certain extent takes the place of the chitinous crest of *Verella*. The structure of the crest is seen in section to be a broadly rectangular triangular cell, subdivided by horizontal bars to form the smaller trapezoidal cells of a second, third, fourth, and fifth story, adjoining triangles being again connected longitudinally by similar bars. The float of *Physalia* remains in fact in the embryonic stage in which we find the sail or crest of *Verella* and young *Porpita*. In the latter the crest gradually disappears to cover the upper part of the float; in the former it continues through to the mature stage, being supported by a chitinous vertical projection from the float, which is absent in younger stages, while in *Physalia* we have only the mantle, if I may so call it, of the crest left, the pneumatophore not secreting a chitinous float or any structure homologous to the circular chitinous float of *Verella* or the chitinous disk of *Porpita*. The presence of a net-work of canals at the base of the float of *Physalia*, similar to that of the upper part of the float of *Verella* *Porpita*, has been made probable by some observations of Quatrefages. (*Ann. Scienc. Nat.* 1853).

sion of the base of the cœnosarc may perhaps be considered as the first indication of the formation of a float. In this case this rudimentary pneumatophore is attached to the ground or to a movable body-like *Fusus*, etc., and forms papillæ, between which arise the two different kinds of polypites, the sterile tentacular polypite and the reproductive polypite, giving rise to either male or female colonies. This combination is exactly similar to that of *Porpita*, in which we have a chitinous float and tentacular and proliferous polypites arranged in addition round a central sterile polypite.

The homologies we have attempted to trace between *Porpita* and the Tubularian Hydroids might perhaps be still further extended. It is well known that in nearly all Tubularians the base of the cœnosarc, by which they are attached to the ground, extends either as filaments or rootlets over a considerable space; these filaments, or expansions of the chitinous tubes, forming either a connected series of canals, more or less complicated, as in *Clava*, *Cordylophora*, *Coryne*, etc., or a net-work of canals as in *Dicoryne*, *Bougainvillia*, *Tubularia*, *Hydractinia*, *Podocoryne*, etc.; or else such filamentary processes as those of *Corymorpha*, in all of which there is a more or less active circulation connected with that of the cavity of the Tubularian. In *Corymorpha* we find a series of longitudinal canals, more or less branching and anastomosing, extending along the cœnosarc. Let us now imagine this long *Corymorpha* cœnosarc reduced in length and at the same time flattened so as to form a disk somewhat below the base of the tentacles, retaining its peculiar pointed terminal basal extremity. We could thus have a free Hydroid differing but little from *Velella* and *Porpita*; that is, our *Corymorpha* would be transformed to a Hydroid, with a crown of marginal tentacles below the chitinous disk, in which there are canals of the vascular system. Between this row of marginal tentacles and the large central opening of the polypite we find clusters of reproductive *Medusæ*. Imagine the same transformation in a colony of *Hydractinia*, or of *Podocoryne*, in which we find the chitinous disk already formed and traversed by a network of canals, and add to it a central sterile polypite, and we have all the structural features of a modified *Porpita*, namely, a disk, rows of tentacular polypites, next rows of reproductive and feeding polypites, while if we make the same comparison with *Hydractinia* we add a third kind of appendage, the so-called spiral zooids.

An examination of very young stages of Tubularians, such as, for instance, the very early stages of *Endendrium* figured by Allman (Pl. XIII, Figs. 14-16, Tubularian Hydroids), show such a chitinous disk to be compared in

every respect with the float of *Velella* or of *Porpita*, with broadly open ramifications communicating with the base of the single large central polypite, and differing from it only in being fixed. We can readily imagine a slightly more advanced stage, with the additional proliferous polypites, or others, developed at the base of the central polypite and setting the attached disk free, we have to all appearances a somewhat modified *Porpita* or *Velella*.

But *Porpita* seems to be also allied to another group of Hydroids, with which thus far no attempt has been made to compare them. I mean the *Hydrocorallinae*. My basis for this comparison rests upon the presence of the singular white plate of Kölliker, and of its peculiar structure,— which reminds us of the porous structure of the corallum of *Sporadopora*, *Allopora*, *Millepora*, etc., although, of course, not having the regular horizontal floors of the latter, yet possessing, like these genera, large pits, the whole mass being riddled with passages and openings, forming the spongy mass of the white plate. Although some of the proliferous polypites of *Porpita* appear to rise from the larger of these pits, yet the others do not seem to hold any definite relation to them, beyond the fact that these proliferous polypites are limited to the ring occupied by this white plate. If this homology is correct, it shows how far-reaching are the affinities of the *Porpitiidae*,— on the one side recalling from the structure of their white plate the corallum of *Milleporidae*, which date back to the cretaceous period; and, on the other, the similar structure of the *Helioporidae*, which, as is well known, have been shown by Moseley to be *Halecyonoids*. Whether the *Stromatopora* have any relationship to either of these groups or are sponges cannot at present be determined; but should they be related to the *Milleporidae*, the peculiar structure of the corallum of the *Milleporidae*, *Stylasteridae*, and *Porpitiidae* would date back to the earliest Silurian. It is interesting to speculate, if the affinity of *Porpita* is greater to the *Milleporidae* with a porous fixed corallum, or to the *Tubularians* having only a chitinous fixed basis.

Porpita Linnæana LESS.

Kölliker* was the first to give a detailed description of the Mediterranean *Porpita*. The Florida species is closely allied to it. It differs from it in size, the largest of our specimens measuring no less than $1\frac{1}{8}$ " in diameter, while the medium size of the *P. Mediterranea* is only 4 to 5" in diameter.

* Die Schwimmpolypen v. Messina, 1853, p. 57, Pl. XII.

As will be seen from the accompanying description, the other differences to be noted between the Florida and the Mediterranean species are: the numerous and close corrugations of the lower surface of the disk; its great roughness on the upper side; the comparatively greater length of the primary polypites; the smaller size of the proliferous polypites. The shape of the former is very different from those of the Mediterranean species, ending in three to four large knobs, with three longitudinal rows of smaller tentacular knobs,—six in two of the rows, and from eight to nine in the central row (Pl. X, Figs. 3, 5, 7).

As in the Mediterranean species, the disk is circular, slightly cup-shaped, convex above, made up of two disks, thickest near the outer edges, and united together by a series of circular walls entirely separated from one another (Pl. XI, Figs. 7-13). The circular chambers thus formed open outwardly by small elliptical openings giving air free access to them (Pl. XI, Figs. 7-9, 12, *a*). These chambers are filled with air (gas?), giving to the *Porpita* its great buoyancy. The enclosed air (gas?), shining through the float and the thin mantle which covers the disk, gives to its upper side a strong silvery lustre. The upper side of the disk is but slightly corrugated, with radial depressions, the stigmata being placed in the centre of low projections, forming an irregular elevation in the lines of corrugations of the disk (Pl. XI, Fig. 7). There is a large central chamber, with an irregular ring of eight smaller ones (Pl. IX, Figs. 1-4), corresponding to the eight first-formed triangular chambers, placed round the central chamber. There does not seem to be beyond that first row any regular arrangement in the stigmata, or any order in their appearance or number. They vary greatly in position and in number in specimens of the same size. The stigmata are more numerous, as well as larger, near the edge of the disk, as has already been noticed by Kölliker.

The ribbed structure of the lower side of the disk does not seem to have been observed. The lower floor of the radiating chambers extends into deep longitudinal corrugations (Pl. XI, Figs. 5, 8-11, 13, 14), forming immense pouches, as it were; so that when seen endwise they present the appearance of high-pointed bags, with rounded tips, and deep spaces between them (Pl. XI, Figs. 13, 14). A transverse (Pl. XI, Figs. 8, 9) and a longitudinal view (Pl. XI, Figs. 10, 11), will give a better idea of their peculiar structure than any lengthy description. To the under side of these pouches are attached the tubules (Pl. XI, Figs. 1, 2, 3, 5, 8, 9), they are irregularly placed in single rows

(Pl. XI, Fig. 5), and are not arranged in sets of three, as they are represented by Kölliker as characteristic of the Mediterranean species. The tubules commence about the outer edge of the white plate, and are most numerous towards its central part (Pl. XI, Fig. 2), where they come to the inner surface, and literally cover it with a matting of winding silvery threads (Pl. XI, Figs. 1, 4). Towards the centre of the float they become less numerous again (Pl. XI, Fig. 3), ending with six or seven tubules, which take their origin near the eight primary chambers, and extend over that part of the float.

Porpita is not as easily upset as *Veella*; and the number of specimens thrown ashore by the winds is very small, as compared to the numbers of *Veella* stranded on the beaches after every storm. *Porpita* is capable of considerable control over its movements. Owing to the great size and power of its numerous long marginal tentacles, it can readily force itself back again into a normal attitude, if upset by the wind or waves. It can, by bringing its tentacles together over the disk (Pl. X, Fig. 1), and throwing up the free edge of the mantle slowly in a given direction, then expanding the tentacles of one side far over on the opposite direction beyond the central part of the disk, it can thus readily change the centre of gravity, and tilt the disk back again into a normal attitude, should it from any cause have been set afloat with the tentacles uppermost. The larger outer marginal polypites are arranged in three or four irregularly concentric rows, with two to three inner rows of smaller knobbed tentacles, in all stages of development (Pl. X, Figs. 2, 3, 10, 11). Inside of these are arranged, in from five to six similar rows, round the base of the large central polypite, the small, stout, flesh-colored feeding and reproductive polypites (Pl. X, Figs. 4, 5). These have a slightly rectangular head (Pl. VIII, Figs. 1-4), capable of considerable expansion, with four clusters of lasso-cells at the thick rounded angles of the terminal opening. At the base of these polypites are found *Medusæ* buds in all stages of development (Pl. VIII, Figs. 1-4; Pl. X, Fig. 4). At the time when in full reproductive power these clusters of *Medusæ* completely fill the whole space between the small polypites, giving to the ring which they occupy on the lower surface of the float a dark yellowish tint from the color of the yellow cells found along the rudimentary proboscis of the *Medusæ* buds, as well as along the chymiferous tubes (Pl. VIII, Figs. 1, 2, 4-11).

The large marginal tentacles (Pl. X, Fig. 3) are of a bluish tint, the tentacular knobs of a darker color. The internal cavity of the tentacles has a

somewhat greenish tint, and connects at the base of the tentacle with the vascular system of the lower part of the mantle at the point of attachment. The cavity leading to the tentacular knobs is very slender (Pl. X, Fig. 8).

The smaller polypites (the feeding-reproductive polypites) occupy on the lower surface that portion of the mantle which covers the ring formed by the so-called white plate of Kölliker, round the base of the single central polypite. These polypites are sometimes seated in cavities of this white plate, and projections of the plate itself also extend sometimes far up into the lower part of the small polypites.

The white plate consists of an irregularly anastomosing system of needles and spurs, or of bars of greater or smaller size, leaving a series of openings for the passage of the tubules (Pl. VIII, Figs. 1, 1^a; Pl. IX, Fig. 11). These tubules take their origin on the lower side of the disk, run in all possible directions between the interstices of the white plate, and come out as blind sacs on its lower side (Pl. XI, Fig. 1). Some of these tubules extend along the side of the small polypites (Pl. VIII, Figs. 1, 2, 4), while others follow the extension of the white plate (Pl. X, Figs. 4, 6), into the base of the central polypite some distance up its walls, forming a most delicate frill (Pl. X, Fig. 4, *z*) of silvery radiating lines, extending towards the mouth of the central polypite. If this white plate is a kind of kidney, as Kölliker suggests, its openings lead outwards through the cavity of the central polypite, as well as through the openings of the smaller reproductive polypites, which are placed on the ring it forms round the central polypite (Pl. VIII, Fig. 1), and into the base of which this white plate extends a considerable distance. Although Kölliker calls it the "white plate," it is in reality of a pinkish color toward the periphery, and blueish towards the interior edge, the whole of the part which lies within the base of the central polypite being of that color. The inner part of the ring of the white plate is composed of heavier bars, the edges only being spongy (Pl. XII, Fig. 15).

The liver (Pl. VIII, Figs. 1, 1^a, *z*, 16) occupies the whole of the space between the lower surface of the disk, the level of the white plate and the base of the central polypite. It fits closely into all the corrugations of the lower side of the disk, as well as into the upper ramifications of the white plate. It sends out a complicated system of radiating and anastomosing tubes from the centre towards the margin of the mantle, in which the circulation is kept up very actively (Pl. XII, Fig. 13). In younger specimens the radiating tubes are still quite simple (Pl. IX, Figs. 1-4; Pl. XII, Figs. 1-8). With

increasing size, they become somewhat more branching, and finally, near the outer edge, more or less parallel with the ring of marginal glands, there is formed a series of irregular horizontal canals connecting the radiating tubes, and forming thus a more or less apparent circular canal (Pl. XII, Figs. 9, 10, 11), till finally the whole free edge of the mantle is covered by a most intricate set of anastomosing tubes (Pl. XII, Fig. 13). The free edge of the mantle terminates by a row of large elliptical glands (Pl. XII, Figs. 4-13; Pl. VIII, Fig. 16), the interior of which is filled by fatty, globular cells (Pl. VIII, Fig. 17). The whole of the free edge of the mantle is of a beautiful clear blue color, with a dark band at the line of contact of the disk and the inner seam of the free edge of the mantle, with a darker blue line on the outer edge of the glands, both at the exterior and interior edge (Pl. VII, Fig. 2). The whole surface of the mantle is covered by a close reticulation of irregular epithelial cells.

The smallest (so-called *Velellæ*) *Rataria* examined by Pagenstecher measured between 0.8 and 2.25^{mm}, and it seems impossible from what I have said here regarding the young stages of *Velella* and of *Porpita* which I have had the opportunity to examine to consider *Rataria* as anything but the young of *Porpita*, as Burmeister had already done. The young *Porpita* in its *Rataria* stage passes through an embryonic stage, in which the young *Porpita* has a prominent sail fully as marked as the sail of the corresponding stage in *Velella*. This stage, to a certain extent, recalls strongly *Velella*. That this embryonic character gradually disappears with age, has been shown by Pagenstecher; but the succeeding stages do not lead, as has been supposed, to *Velella*, but to *Porpita*. At no time in the development of *Velella* do we have any trace of eight compartments arranged round a central chamber. We have, it is true, a central chamber; but there are only concentric chambers in the earliest stages I have seen; while the *Rataria* stages figured by Pagenstecher* correspond admirably to the young stage of *Porpita* I found at the Tortugas, in which the eight central chambers occupy the greater part of the disk, and can, as is well known, still be traced in fully grown specimens.


* *Zeits. f. Wiss. Zool.*, XII., 1863, Pls. XL, XLI, p. 496.



PLATE I

FIG. 1. *Verella mutica*, seen in profile ; natural size.

FIG. 2. The same, seen from above.

FIG. 3. The same, seen from below, with a small cluster of reproductive polypites, magnified four diameters.









PLATE II.

- Fig. 1. Larger kind of reproductive polypite of adult *Velella*: *a*, actinal opening; *b*, clusters of lasso-cells forming patches over the surface of the upper extremity of the polypite; *mm*, clusters of medusæ buds in different stages of development, magnified.
- Fig. 2. Actinal opening of feeding polypite. Fig. 4.
- Fig. 3. Another view of the actinal opening of a reproductive polypite.
- Fig. 4. Magnified view of larger feeding polypite; in these the patches of lasso-cells are but slightly developed towards the free extremity; near the base, however, they form huge spheres *bb*, studded in part with gigantic lasso-cells.
- Fig. 5. Smaller kind of reproductive polypite, more slender than Fig. 1, with a smaller number of medusæ buds. The patches of lasso-cells are clustered together so as to form hemispherical projections, which become slightly pedunculated towards the base.
- Fig. 6. Magnified view of two of the hemispherical clusters of lasso-cells of the surface of one of the smaller prolific polypites. These polypites are open at the extremity.
- Fig. 7. Another view of a smaller prolific polypite distended with food.
- Fig. 8. Small and slender feeding polypite. Lettering as in Fig. 4.
- Fig. 9. Actinal extremity of one of the smaller kinds of feeding polypites greatly extended.
- Fig. 10. Medusæ of *Velella* freed twelve hours from the reproductive polypites: *d*, digestive cavity; *ch*, chymiferous tubes; *y*, yellow cells; *l*, one of the large lasso-cells forming the four lines running from the tentacles to the apex; *tl*, large and small rudimentary tentacle; *m*, actinostome.
- Fig. 11. Somewhat older than preceding stage. Lettering as above.
- Fig. 12. Fig. 10 seen from above with the sides of the bell slightly drawn in.
- Fig. 13. Young *Velella* just before it separates from the prolific polypites.
- Fig. 14. Magnified view of the base of the rudimentary tentacular bulb: *l*, outer line of lasso-cells in the periphery of the bell; *tl*, rudimentary tentacle; *y*, clusters of yellow cells.
- Fig. 15. Medusa of *Velella* copied from drawings for L. Agassiz's Contributions to the Natural History of the United States.
- Fig. 16. Same seen from the abactinal pole.






PLATE III.

- Fig. 1. Float of young *Verella* seen from above, $\frac{1}{8}$ of an inch in length: ff' groove in which the main branch of the circulatory system passes from the fixed edge of the float to the base of the keel. In this stage the keel is a slight ridge surmounting the single central chamber: $o'o'$, apertures opening externally from the fifth and seventh chamber on the right, and from the second and fourth on the left.
- Fig. 2. The same as Fig. 1, seen in profile, showing the central chamber round which are arranged, in an irregular elliptical shape, the eight chambers of which the float, at this stage, is composed: tt , so called air tubes attached to the lower side of the float; f , groove of main branch of circulatory system.
- Fig. 3. Float of a young *Verella* measuring $\frac{3}{8}$ of an inch. Lettering as above; $o''o'''$, apertures opening externally from the ninth and sixth chambers on the right and left.
- Fig. 4. Profile view of the float of a young *Verella* having eight chambers, measuring $\frac{1}{4}$ of an inch in length. Lettering as in Fig. 2. The keel, a' , rises quite perceptibly above the central chamber on a slight conical projection: a^1-a^8 , successive chambers of the float.
- Fig. 5. Profile view of Fig. 3. Lettering as before. The keel has greatly increased in height and the second line of growth is apparent.
- Fig. 6. Float of young *Verella* with twelve chambers measuring $\frac{1}{4}$ of an inch in length and having seven lines of growth in the keel, which, in this stage, is a high conical lamella of entirely different proportions to the float than in the adult or larger specimens.
- Fig. 7. Profile of central part of the float and keel of a young *Verella* with twelve chambers, but only four lines of growth in the keel; this is also much broader than is usually the case at this stage, — $\frac{5}{16}$ '' in length.
- Fig. 8. Profile of young *Verella* with seventeen chambers in the float, measuring $\frac{7}{16}$ '' in length, with six lines of growth in the keel.
- Fig. 9. Portion of horizontal surface of the float of young *Verella* with twelve chambers, showing the openings leading from one chamber to the adjoining ones through the common walls; the opening in the last chamber opens externally.
- Fig. 10. Showing on a somewhat larger scale the passages leading from one chamber to the other in the eight outer chambers of the float of the preceding figure.
- Figs. 11, 12, 13. Opening in the outer chamber of three young *Verella* varying in size from $\frac{1}{4}$ to nearly $\frac{1}{2}$ an inch in length. This opening is always in continuation of the line of openings forming the communication between adjoining chambers.
- Fig. 15. Diagrammatic transverse section of young *Verella* with twelve chambers, showing the position of the keel a' , the central chamber a , and the concentric chambers a^1-a^{12} with the groove f for the main branch of the circulatory system.
- Fig. 16. Diagrammatic section of the float of a *Verella* with twenty-two chambers.
- Fig. 17. The corresponding section of the opposite side showing the position of the groove f . This groove is only a fold of the upper walls of the chambers, and does not divide the chambers into distinct spaces; they form each a continuous ring.
- Fig. 18. General view of the position of the clusters of tubules arising from the lower side of the four inner chambers; some of the tubules branch, a rare occurrence in our *Verella*.
- Fig. 19. Inner view of the conical part of the float of a large *Verella*, showing the position of the tubules and the corrugations and folds of the lower walls of the inner chambers; the corrugations ccc become less prominent in proportion to the distance from the central chamber; near the margin of the float the walls merely bulge out and form slight undulations.
- Fig. 20. Tubules of a large *Verella* which have penetrated through the so-called liver, and appear on the top of the liver when seen from the lower side of the cone of the float.
- Fig. 21. View from the inside of the central part of the float showing the tubules of a young *Verella* measuring about $\frac{1}{4}$ '' in length.
- Fig. 22. Basal part of three tubules near the attachment to the central chamber in a *Verella* measuring about $\frac{1}{2}$ an inch in length.






PLATE IV.

- FIG. 1. Profile view of the keel and float of small *Veella*, measuring about $\frac{1}{3}$ inch in length: *ca*, outer chamber of the float; *a*, one of the concentric chambers; *l*, so-called liver; *p*, folds of the mantle covering the keel extending from the canal at the base of the keel to the canal *e* at the edge of the mantle.
- FIG. 2. Small *Veella*, about $\frac{1}{8}$ of an inch in length, (seen from above) showing the few ramifications of the so-called liver system, *l*, seen through the float; *s*, rudimentary keel; *f*, free edge of the mantle.
- FIG. 3. Slightly more magnified view of a part of the edge of a young *Veella*, somewhat older than the preceding figure: *r*, principal ramifications; *r'*, secondary ramifications of the free edge of the mantle.
- FIG. 4. Portion of the edge of a small *Veella*, measuring $\frac{3}{8}$ of an inch in length: *v''*, primary ramifications; *v'*, secondary branches; *g*, marginal glands of the free edge of the mantle, seen from above.
- FIG. 5. Portion of the edge of a small *Veella*, slightly older than the preceding figure, in which the principal ramifications still extend to the free edge of the mantle.
- FIG. 6. Portion of the edge of the float of a small *Veella*, about in the same stage as the preceding figure, with proportionally larger marginal glands, but shorter secondary ramifications, than in Fig. 4.
- FIG. 7. Portion of the edge of the mantle of a *Veella* measuring about one inch in length, with long marginal glands *g*, with a simple free edge of the mantle *e*, extending beyond, and small clusters of yellow cells *c*, moderately closely packed along the outer edge of the mantle.
- FIG. 8. Portion of the edge of the mantle of a *Veella*, measuring about two inches in length, with a fringed free edge of the mantle *e*, beyond the marginal glands *g*, with large patches of yellow cells *c*, closely packed together over the whole of the free edge of the horizontal mantle beyond the float.
- FIG. 9. Somewhat more magnified view of a part of the horizontal mantle of the float, to show the shape of the ramifications of the so-called liver canals extending between the patches of yellow cells.
- FIG. 10. Single cluster of yellow cells with the fan-shaped terminations of the liver canals.
- FIG. 11. Portion of the outer edge of the horizontal mantle of a *Veella*, in which the ramifications of the vascular system terminate in sacs more or less filled with clusters of yellow cells.
- FIG. 12. Separate clusters of terminal extremities of the vascular system of a *Veella* nearly in the stage of the preceding figure, taken near the edge of the float.
- FIG. 13. Pouch-like terminal clusters of the vascular system filled with yellow cells, seen from above near the glandular band of the mantle.
- FIG. 14. Shows the delicate dendritic ramifications of the large vascular (liver) canals seen from the lower side of the float, entirely free from accumulations of yellow cells.
- FIG. 15. Part of one of the main vascular tubes, with large star-shaped ramifications and large clusters of yellow cells, seen from the upper surface, near the float.






PLATE V

- FIG. 1. Half of the keel of a *Veella* measuring one and a half inches in length, showing the principal tube r , with its anastomoses extending to the broad marginal canal, from which proceed the dendritic tubes extending to the edge of the mantle of the keel of Figs. 4, 5.
- FIG. 2. Somewhat more magnified view of a portion of the keel of the preceding figure, showing more distinctly the principal central branch, with its anastomoses, and the large blind sac-like pouches (see Fig. 10) formed along certain parts of the main circular canal of the keel. The pouches are filled with clusters of yellow cells; there are also smaller patches of yellow cells in the circular canal.
- FIG. 3. Still more magnified than preceding figure, to show the position of the clusters of yellow cells y , near the edge and in the circular canal.
- FIG. 4. Half of the keel of a *Veella* slightly larger than the preceding figures, showing the main central vascular branch r , the anastomosing branches v'' , extending over the whole surface of the keel to the circular canal v^o , running nearly parallel to the outer free edge of the keel. From this circular canal branch off the dendritic canals v' , which unite again along the free edge of the mantle of the keel.
- FIG. 5. Magnified view of two of the dendritic canals of the free edge of the mantle of the keel; v^o , circular canal, from which arise the main branches v' , of the dendritic canals v'' , which unite again in a marginal canal v''' , along the outer edge of the keel mantle.
- FIGS. 6, 9. Ramifications of the vascular system from the lower side of the float, with clusters of apparently decomposed or dead yellow cells.
- FIG. 7. Magnified view of the frill-like ramifications, $v v'$, of the so-called dendritic branches of the vascular system of Fig. 5, with small clusters of yellow cells $c c'$.
- FIG. 8. Enlarged view of tubes of the vascular system of the keel, to show the mode of ramification and anastomosis over the surface of the keel.
- FIG. 10. (See Fig. 2.)
- FIG. 11. General appearance of the net-work and principal branches of the so-called liver system ll' , under the conical part of the float a ; seen in profile.
- FIG. 12. General view of the ramifications of the same system; extending under the float; seen from above.
- FIG. 13. Branch of liver system and anastomoses at the junction of the float and of the horizontal mantle.







PLATE VI.

- FIG. 1. Young *Velella* seen in profile, measuring about $\frac{3}{16}$ of an inch in length: *c*, central chamber; *l*, central part of the liver in the conical part of the float; *f*, flexible keel formed by the free edge of the mantle; *cp*, central feeding polypite; *t*, prehensile tentacles; *fp*, small feeding polypites.
- FIG. 2. Same seen from the lower side, with single ramifications of the liver system extending to the edge of the horizontal mantle.
- FIG. 3. Somewhat younger than the preceding stage. The ramifications of the liver system are less pronounced; the central feeding polypite is not expanded as in the other stages.
- FIG. 4. One of the large blue prehensile marginal tentacles, *t*, of the preceding figure.
- FIGS. 5, 6, 7. Different stages of growth of the small feeding polypites of Fig. 3.
- FIG. 8. Smallest *Velella* found at the Tortugas, with a hood-like free mantle, with only four chambers, a conical float, much like that of Pl. III, Fig. 1, in shape, surmounted by a flexible flat elliptical crest, with a large central polypite, eight large prehensile marginal tentacles, and the same number of small rudimentary feeding polypites.
- FIG. 9. The same as Fig. 8, seen from above, showing the simple tubes of the liver system of that stage.
- FIG. 10. Central polypite of a large *Velella* slightly drawn in.
- FIG. 11. Different attitude of the same.
- FIG. 12. Basal part of the same when fully expanded.
- FIG. 13. Distal extremity of the same when fully extended.
- FIG. 14. Terminal part of one of the prehensile tentacles, showing the band of lasso-cells running along the edges.
- FIG. 15. Somewhat more magnified view of a part of such a tentacle, showing the terminal band of lasso-cells.
- FIG. 16. Part of the base of a prehensile tentacle where the band of lasso-cells becomes broken up in patches.
- FIG. 17. One of the prehensile tentacles showing the passage of the terminal band of lasso-cells into patches near the base.
- FIG. 18. Young prehensile tentacle, with large patches of lasso-cells, but without a terminal band.
- FIG. 19. Portion of prolific polypite with young medusæ buds *m*, and the extremity of the tubule *t*.
- FIGS. 20, 21. Ramifications of the vascular system at the base of the feeding polypites on the mantle of the lower side of the float.
- FIG. 22. Enlarged view of the central feeding polypite of an adult *Velella*, showing the rudimentary lips of the actinal opening and the longitudinal rows of patches of lasso-cells *b*.



PLATE VII.

- Fig. 1. Porpita seen in profile fully expanded, $1\frac{1}{8}$ " in diameter.
Fig. 2. The same seen from the upper side of the float.





PLATE VIII.

- FIG. 1. A portion of the disk of a *Porpita*, showing the white plate with its spongy ramifications *i*, extending into the base of the feeding and reproductive polypites (*c p*, *f p*), with the so-called liver *l*, placed between the lower floor of the chambers of the float *c*, and the spongy upper part of the white plate; with the tubules *t*, extending from the lower side of the float through the mass of the liver, many of them passing through the white plate and extending along the base of the feeding and reproductive polypites.
- FIG. 1*. General view showing the relative position of the white plate *i*, the liver *l*, the chambers of the float *c*, and of the tubules of the upper and lower side of the plate.
- FIG. 2. A cluster of feeding and reproductive polypites: *t*, tubules extending along the base; *m*, medusæ buds in different stages of development.
- FIG. 3. A single reproductive polypite with two very young medusæ buds *b b*.
- FIG. 3*. Actinal opening of one of the feeding polypites.
- FIG. 4. A single reproductive polypite with clusters of medusæ *m*, in different stages of development and the tubule *t*, extending more than half the length of the polypite towards its extremity.
- FIG. 5. Different view of a single reproductive polypite, with two clusters of medusæ.
- FIG. 6. Young medusa just freed from the reproductive polypite: *d*, canal leading to the cavity of the polypite through the base of attachment; *y*, clusters of yellow cells; *c*, large lasso-cells.
- FIG. 7. Young medusa, somewhat older, more elongate, lettering as in the previous figure. The lasso-cells extend along the sides of the bell.
- FIG. 8. Young medusa, stage intermediate between Figs. 6 and 7.
- FIG. 9. Same seen from above.
- FIG. 10. Fig. 6 seen from above.
- FIG. 11. Fig. 7 seen from above.
- FIG. 12. Abactinal part of the bell of Fig. 7: *c h*, chymiferous tubes; *d*, rudimentary digestive cavity.
- FIG. 13. Portion of the edge of the bell of a medusa in the stage of Fig. 7, lettering as before.
- FIG. 14. One of the large lasso-cells of the surface of the bell with the coil partially thrown out.
- FIG. 15. Young medusa still attached to the reproductive polypite.
- FIG. 16. A portion of the free edge of the mantle: *d*, edge of the disk; *g*, one of the marginal glands.
- FIG. 16*. View from the lower side of the disk, to show a portion of the ramifications of the liver.
- FIG. 17. Fatty globules filling the marginal glands.

1880

1881

1882

1883

1884

1885

1886

1887

1888

1889

1890

PLATE IX.

- FIG. 1. Young *Porpita* with the eight primary chambers, surrounded by simple straight vascular canals with eight large primary four-knobbed marginal tentacles t , sixteen secondary tentacles t' , of about half the length of the first set, and a third set of still smaller tentacles t'' , from thirty to fifty in number, alternating between them and scarcely projecting beyond the margin of the disk. The disk measures only $\frac{1}{32}$ " in diameter.
- FIG. 2. A somewhat older stage, the disk measuring about $\frac{1}{3}$ more in diameter than the preceding stage. The tentacles have greatly increased in length and are proportionally more slender; the primary and secondary tentacles also have two coils of tentacular knobs.
- FIG. 3. Still older stage, the primary chambers occupy a comparatively smaller area of the disk; the secondary tentacles, t' , are nearly as long and slender as the eight primary ones, t ; and the third set of tentacles, t'' , have also developed tentacular knobs.
- FIG. 4. Ramifications of the vascular system in an older stage than the preceding, in which the eight primary chambers, with their openings o' , are now surrounded by irregular circles of openings, o'' , o''' , leading to the second and third concentric chambers of the disk.
- FIG. 5. Extremity of the primary tentacle of a young *Porpita*, in the stage of Fig. 2.




PLATE X.

- Fig. 1. Profile view of a large Porpita measuring $1\frac{1}{16}$ " in diameter, with the tentacles swung under the disk and the disk thrown up.
- Fig. 2. Disk seen from above to show the arrangement of the tentacles of a young Porpita, measuring about $\frac{1}{2}$ " in diameter.
- Fig. 3. Magnified portion of a part of the disk of Fig. 2, and of the corresponding marginal tentacles; *pp*, proliferous polypites.
- Fig. 4. Portion of the lower side showing the mouth of the central polypite *m*, and the rows of reproductive polypites *pp*, with the tubules *tt*, extending over the base of the central polypite.
- Fig. 5. Magnified view of a portion of the disk showing the central polypite *cp*, fully expanded, with the rows of reproductive polypites *pp*, between the base of the central polypite and the marginal tentacular polypites *tp*.
- Fig. 6. Magnified view of a small reproductive feeding polypite *pp*, with a cluster of medusæ in different stages of development, and with the meandering tubules extending over its base.
- Fig. 7. Extremity of one of the largest marginal tentacular polypites, showing the arrangement of the tentacular knobs.
- Fig. 8. Magnified view of one of the tentacular knobs.
- Fig. 9. Folds of the actinal opening of the central polypite.
- Fig. 10. Profile view, showing the arrangement of the tentacular marginal polypites.
- Fig. 11. Different stages of growth of the smaller marginal tentacular polypites.




PLATE XI.

- Fig. 1. Portion of the lower side of the disk, showing a part of the white plate *ss*, with the ramifications of the liver *ll*, and the course of the tubules *tt*, which have passed through the white plate, with portions of the corrugations of the lower side of the circular chambers *c*.
- Fig. 2. Diagrammatic sketch showing the relative position of the disk, of the edge of the white plate, and of the mantle.
- Fig. 3. Central chamber *a*, with the adjoining chambers from which lead the first rows of tubules *t*.
- Fig. 4. Cluster of tubules on the dorsal side of the white plate.
- Fig. 5. Corrugations of three of the inner chambers of the lower side of the disk, showing the short tubules and their mode of attachment, about under the white plate.
- Figs. 6, 6'. Greatly enlarged view of a few joints of one of the tubules.
- Fig. 7. Part of the outer portion of the disk, showing the arrangement of the stigmata *o*, on the upper surface.
- Fig. 8. Slightly oblique view of the cross-section of the outer part of the disk showing the position of the stigmata *oo*, of the successive chambers *aa*, and of the primary and secondary corrugations on the lower side, *c* and *c'*, of the circular chambers, and the mode of attachment of the tubules *tt*.
- Fig. 9. Different profile view of a portion of the disk. Lettering as before.
- Fig. 10. Shows the general arrangement of the simple corrugations, *c*, of the lower side of the circular chambers, in a young specimen.
- Fig. 11. Portion of the lower side of the disk showing the folds of the lower side of the chambers, *c c'*, as older far as the last outer circular wall, *ppp*, of the disk.
- Fig. 12. Portion of the disk of a young *Porpita*, $\frac{3}{8}$ " in diameter, with four circular chambers; *o'-o'*, stigmata, *ff'*, fold in the chambers corresponding to the original eight chambers, and forming in older specimens the irregularly radiating lines of the upper side of the disk.
- Fig. 13. Slightly oblique end view of a part of the corrugations, *c c'*, of the last chamber of the edge of the disk.
- Fig. 14. An end view of a few corrugations of a similar part of the disk.



PLATE XII.

- Fig. 1. Portion of the disk of a young *Porpita*, measuring over $\frac{1}{8}$ " in diameter, seen from the upper side.
- Fig. 2. Same seen from the lower side.
- Fig. 3. Portion of the disk of a young *Porpita*, measuring $\frac{3}{8}$ " in diameter.
- Fig. 4. Young *Porpita*, measuring $\frac{3}{8}$ " in diameter, in which a portion of the disk of the vascular system has remained simple. Seen from above.
- Fig. 5. Portion of the disk of a young *Porpita*, nearly in stage of Fig. 4, seen from above, but the vascular system is branching.
- Fig. 6. Portion of the disk of a young *Porpita* at about the same stage with simple vascular system, but proportionally longer tubes.
- Fig. 7. Portion of the disk of a young *Porpita* about in stage of Fig. 6, seen from the lower side of the disk.
- Fig. 8. Portion of the disk of a young *Porpita* somewhat more advanced, the vascular system more branching.
- Figs. 9, 9'. Portion of the disk of a young *Porpita* measuring about $\frac{1}{4}$ " in diameter. The ramifications of the vascular system are limited to the margin of the mantle.
- Fig. 10. Portion of the disk of a young *Porpita* measuring $\frac{5}{8}$ " in diameter.
- Fig. 11. Portion of the disk of a *Porpita* measuring not quite $\frac{1}{2}$ " in diameter.
- Fig. 12. Marginal folds of the edge of the mantle.
- Fig. 13. Portion of the disk of a *Porpita* measuring 1" in diameter, showing the system of anastomosing tubes of the mantle.
- Fig. 14. A portion of the same, showing the star-like irregular termination of this anastomosing system.
- Fig. 15. Magnified view of a part of the white plate.



ERRATA.

- Page xxix. Transpose from line 11 to line 10, so as to read "occur the following variations : one" etc.
- " 27. TROP. LEBERIS, line 3, read "Nasal grooved or divided."
 - " 53. Under *Scotophis*, and p. 54 under *S. obsoletus*, read "Anal bifid."
 - " 64. Sixth line, for 23, read 25.
 - " 65. After ELAPSOIDEUS, insert "pl. V, f. 2."
 - " 66. After DOLIATUS, erase "pl. V, f. 2."
 - " 85. Erase CHEILORHINA and C. VILLARSII.
 - " 153. Under var. VULPINUS, read "p. 299."
 - " 154. Under OPHIOLUS DOLIATUS, erase reference to Dum. Bibr.
 - " 155. Var. CALLIGASTER, line 1, after "Harl." insert "1827, Jour. Phil. Ac., V, 359, —"
 - " " Under var. DOLIATUS, line 3, after "f. 2," insert "D. & B., 1854, Erp., 621."



Memoirs of the Museum of Comparative Zoölogy

AT HARVARD COLLEGE, CAMBRIDGE, MASS.

VOL. VIII, No. 3.

THE
REPTILES AND BATRACHIANS
OF NORTH AMERICA.

By SAMUEL GARMAN.

PUBLISHED BY PERMISSION OF THE KENTUCKY GEOLOGICAL SURVEY.

WITH TEN PLATES.

YEOMAN PRESS, FRANKFORT, KY.:

MAJOR, JOHNSTON & BARRETT.

1883.

PRELIMINARY NOTE.

The preparation of the following memoir was begun by Mr. Garman during my term of service as Director of the Kentucky Geological Survey. Circumstances have made it necessary to delay its publication to the present date.

In the act of 1874, providing for the continuance of the Geological Survey, the Legislature of Kentucky permits "that the scientific results of the Survey may be published in any scientific journal by permission of the Director." Although this memoir will form a part of the second volume of *Memoirs of the Kentucky Survey*, it seems to me highly desirable that it should secure the wider circulation that will be given to it by this publication in the memoirs of the Museum of Comparative Zoölogy. The Kentucky Survey is fortunate in being able, by this arrangement, to acknowledge the large debt it owes to this Museum for most effective aid in every step of its scientific work.

N. S. SHALER.

Formerly Director of the Kentucky Geological Survey.

CAMBRIDGE, MASS., June, 1883.

ON THE
REPTILES AND BATRACHIANS,

By SAMUEL GARMAN.

INTRODUCTION.

REPTILES AND BATRACHIANS.

The Reptiles and Batrachians belong to what are commonly called the Cold-blooded Vertebrates. This name is applied because the heat-producing capacity of their bodies is so low as to render them very susceptible to thermal variation in the surrounding air or water, and unable to maintain any standard temperature. In Batrachia that breathe by means of gills, the small amount of heat the body is capable of producing is limited by the amount of oxygen in the water. A heart in which ventricles or atria are more or less incompletely separated mixes venous with the arterial blood supplied the tissues of lung-bearing Reptiles and Batrachians; and thus, in consequence of partial oxygenation, their heat-producing capacity is reduced. Animals of these classes are, to a considerable extent, dependent on external heat; they are more active during the warmer portions of the season or of the day. During the winter of the temperate zones or the dry season of the tropics they are comparatively inactive.

The Turtles, Crocodiles, Lizards, Amphisbaenians, and Snakes are Reptiles. They are hatched or born with the shape of the adult, breathe by lungs, and generally are covered by a skin the outer layers of which are folded so as to resemble scales. The Batrachians include such as the Toads, Frogs, Salamanders, Newts, Sirens, and Cæcilians. Nearly all of them breathe and progress like fishes during the earlier portion of their existence, and the majority go through a metamorphosis, taking on the form of the adult and breathing by lungs later in life. They are without the scale-like folds of the Epiderm.

From the Turtles to the Snakes of the one, and from the Frogs to the worm-like Cæcilians of the other of these classes, there is a great diversity of forms and habits. In all the intelligence is of a low order. Commonly one or more of the senses is feebly or not at all developed. Many are slow in growth, and live to great ages. Most are tenacious of life, and able to do without food or drink for long periods. The greater portion are harmless; the exceptions are such as the Crocodiles, the few venomous, and a

few large non-venomous serpents. Depending so much upon small vertebrates and insects for food, the majority are important checks upon the increase of the enemies of garden and field. Occasionally farmers recognize the compensation received for protecting and gathering these despised creatures; more often, unfortunately, in prejudice and ignorance of the friendly habits, they exert themselves in favor of extermination. Even the lizards that dart about on the bark, branches, and leaves in search of grubs, borers, and other pests of the orchard, are included in the general massacre. The dangerous serpents of the United States are the Rattlesnake, the Moccasin, and Copperhead, and, though the error is on the side of safety, the popular fear of these is much greater than is warranted by fact.

The flesh of most of the Turtles, many of the Lizards, and that of numerous Batrachians, is excellent for the table. Aside from their value on account of flesh and habits of feeding, the Batrachia form an item of food for fishes that is not to be overlooked by those engaged in stocking ponds and streams.

REPTILES.

There is a great number of Reptiles among the fossils. Some of these belong to recent genera; others are of enormous size, and pertain to genera long ago extinct. It is the purpose, however, to confine attention in this work to living forms, and in the main to such as now exist in North America. These are included in four orders: TESTUDINATA, Turtles; RHIZODONTA, Crocodiles and Alligators; SAURIA, Lizards; and OPHIDIA, Serpents.

TESTUDINATA.

The form typical of this order has a short, clumsy body, of which the vital portions are inclosed in a bony case, formed by the expansion and consolidation of the backbones, ribs, and sternum. Perhaps it might be more exact to say the shell or box is formed by a union of dermal and neural skeletons. Young turtles have spaces covered by fibro-cartilage between the ribs, near the margins. Among fresh water turtles, *Trionycida*, this peculiarity is persistent; the upper shell, *carapace*, occupies only the central portion of the back, and the lower, *plastron*, is also incomplete. The Trunk-back, *Sphargis*, has a leathery case rather than a shell. The epiderm has

the form of thin plates or scales; these are much thickened in the Hawk-bill, *Eretmochelys*, and form the tortoise shell of commerce; they are hardly noticed in such genera as *Trionyx* and *Sphargis*. The feet and limbs vary greatly according to habits. Marine species have broad paddles in which the toes are not distinct, and the claws appear as hooks on the margin. Certain fresh-water species have the toes broadly webbed and furnished with strong claws for tearing to pieces the creatures on which they feed. And in the land Turtles, *Testudo*, foot, ankle and leg resemble clubs or the feet of elephants. The skull is short, massive, and bears but a single condyle. Ordinarily the neck is long and flexible, and the tail is short; in sea turtles, however, the neck is comparatively short, and the tail of the "Snapper" is long.

The sight is keen. The eyes are usually somewhat large and very mobile; they are protected by two lids and a nictitating membrane. The hearing is tolerably acute; the tympanum is not protected by valves or cavity. As would be expected from the mode and infrequency of breathing, the sense of smell is dull. The tongue is thick and fleshy. From the manner in which domesticated species select their food, there can be little doubt of the possession of taste. This sense is probably more developed in the vegetable-eating land tortoises. No specialized tactile organs are known, unless, it may be, in the fleshy lips of some or barbels in exceptional cases, as the Matamata. Respiration is effected by muscular exertion or swallowing. Many, if not all, turtles have voice. The most recent verification of this has been in *Clemmys Muhlenbergii*, by Dr. Abbott. Few creatures are more sensitive to changes of the weather. A passing cloud often drives them to shelter.

All turtles are oviparous. The eggs are fertilized within the oviduct; they are deposited in the sand or earth, and left to be hatched by the heat of the sun. The ovaries and testes are paired; the intromittent organ of the male is simple. When in copulation the male is carried on the back of the female. On the paddle of male sea turtles the first claw is modified so as to form a hook to grasp the edges of the shells of their mates. Males of many species are distinguished by a depression or concavity in the plastron.

Land Turtles generally have a well-ossified, high-arched shell, club feet, and long necks. They travel slowly, feed on vegetables, and are able to endure fasts of great length. Sometimes they make long journeys for

water along accustomed routes. When drinking the head is plunged into the liquid, which is taken slowly in gulps, as if to prolong the enjoyment as much as possible. South America and the Galapagos Islands possess the largest living species. Our largest is the "Gopher" of the Southern States, *Xerobates*. "Box Tortoises," *Cistuda*, are not uncommon east of the Rocky Mountains. In these both valves of the plastron are hinged, and, when the head, tail, and limbs are drawn within, can be closed tightly against the carapace.

Mud Turtles have a more elongate shell than the preceding; and the height, compared with the length, is less. In structure the plastron varies considerably, most often it is rigid; one or the other, or both, of the lobes may be hinged. Their food is mixed—animal and vegetable. Species of *Emys*, *Cinosternum*, and *Ozotheca* are common examples. One of the largest that might be placed in this group is the gigantic river tortoise of the Amazons, *Podocnemys*. The largest North American inland turtles are the "Snappers," *Chelydra*, and *Gypochelys*. In them the head is large, tail long, and plastron reduced in size.

Fresh-water Turtles, such as belong to the genera *Amyda*, *Platypeltis*, and *Aspionectes*, are seldom seen on the land. Their shells are much depressed and the margins are leathery. The nostrils are produced beyond the snout in a sort of proboscis, which admits of taking air without disturbing the surface of the water or showing the head. They are carnivorous, and feed on fishes, batrachians and worms. The feet are webbed, furnished with strong claws, and the marine species are hardly more expert in swimming. Some of these are quite as savage as the snappers.

The flesh of the majority of the land, mud, and water species is excellent. When brought to market they are ordinarily known by the name "Ter-rapin," though that best known by this title is the marsh turtle, *Malacoclemmys palustris*.

Sea Turtles are numerous off the coasts of Florida. "Trunk-backs" or "Leather-backs," *Sphargis*, are the largest, attaining more than a thousand pounds in weight. As the common name suggests, carapace and plastron are not ossified as in other marine genera. *Sphargis* is very erratic. Specimens from the Gulf of Mexico find their way northward to the coasts of New England, and are supposed to cross the Atlantic, individuals having been taken at the British Isles. Loggerheads, *Thalasseochelys*, are more common. These reach a weight of four hundred pounds. They are said

to eat almost anything—seaweed, sponges, crustacea, mollusks, fish, etc. A massive sponge growing around the Florida Keys is called Loggerhead sponge, because, as the fishermen say, the turtles are very fond of it. They take Conchs from their shells by biting off the small end of the spiral. The Bastard Turtle, *Colpochelys*, is smaller than, and intermediate between, Loggerheads and Green. Green Turtles, *Chelonia*, are reported to weigh as much as eight hundred pounds. They are most prized for the markets. A peculiar grass, *Zostera*, growing on the bottom in shoal places, is said to be their main dependence for food. A great many are shipped from Florida and the West Indies to the Northern States and Europe. Hawkbills or Shell-backs, *Eretmochelys*, are valued as the source of the shell used in the arts. One hundred and sixty pounds is a large weight. All of the marine species are used as food, but the Green is most sought. Many men do little else than supply the markets with the meat and eggs. According to an expert in these matters, Richard M. Kemp, turtlers most often make their captures by means of the peg. Bits of the grass floating above the grazing turtles betray their whereabouts, and if they are not startled it is not difficult to fasten the peg in the shell. A peg is an iron instrument like a blunt nail, to which a line is attached, and which, when driven into the shell, easily slips out of the end of the long pole in which it is placed. By means of the line the turtle is drawn on board the boat. Mr. Kemp says that the Bastard and the Leather-back couple and lay in December, January and February, and the Hawkbill, Loggerhead, and Green in April, May and June. When about to deposit the eggs, the only time the shore is visited, the female selects some sandy beach, and in the night drags herself out above high-water mark. Here she digs a hole of one to two feet in depth, in which she drops seventy-five to two hundred eggs. She then covers the nest carefully, packing down the sand with her body, and retreats to the water. Fourteen or fifteen nights afterward she returns to make another nest near the first. Three to five times in a season, the fishermen say, she returns in this way. If a turtler going his rounds in the morning finds the tracks made on the way to and from the nest, he takes a short stick and carefully thrusts it down here and there in the trampled space until pushed through an egg, the yolk upon the probe discovers the nest. Fourteen nights later he is on the ground waiting till the creature comes ashore, when he turns her on her back, and in the morning carries her to the markets. Persons in the business claim that there is no

diminution in numbers, which, in view of the wholesale destruction, seems hardly possible.

The eggs are spherical and soft-shelled—that is, they are covered by a thick skin, on which the limy covering is thin. The young are furnished with a sharp point in front of the nostrils, like that on the beak of a young bird, with which they tear their way out. At once, on freeing themselves from the sand, they take to the water. I am told these animals are very fond of the “Portuguese Men-of-War,” *Physalia*; and that, when engaged in eating them, they shut their eyes to avoid the stings, and may be taken into a boat without pegging. Nets are also used in taking them.

RHIZODONTA.

Crocodiles have long bodies, long tails, short necks, and rather short legs. The heads vary greatly in shape; in some the snout is long and narrow; in other genera it is pointed, and in the alligators it is broad. The condyle is single; generally the vertebrae are concave in front. A complete separation exists between the ventricles, but communication between the atria mixes venous with the arterial blood going to the posterior portion of the body. In the skin of the dorsal portions there are bony plates. The epiderm is thin and corneous, closely attached to the skull, and arranged over body and tail in transverse series of rectangular plates. The mouth is large, and the jaws are strong. The teeth are numerous, conical in some, compressed in others, and in all received in sockets by their bases. The tongue is fleshy, short, non-extensile, and adherent to the floor of the mouth. Nostrils, eyes, and ears are situated prominently on the top of the head, which enables the animal to breathe, see, and hear without exposing itself. The eye is provided with three lids; the pupil varies from horizontal to vertical in the different genera. By means of valves the tympanic cavity may be closed against the water. At the back of the tongue a valvular arrangement closes the mouth cavity against the water when struggling with the prey below the surface. In some respects the stomach resembles that of birds; it is rounded and the coating is thick. The abdominal is separated from the pectoral chamber by a muscular diaphragm. Crocodiles are carnivorous; they commonly drown their prey, afterward going to the shore to eat it. They are said sometimes to wait until it is partially decomposed. Their flesh is rarely, if ever, eaten. Recently the skins have come into use to a considerable extent for leather. The eggs are oblong,

and have a hard shell; they are left in holes in the earth and sand to hatch. The cloacal aperture is a longitudinal slit; the penis is single. The Rhizodonts are long lived; the young grow rapidly, adults more slowly. Specimens have been taken of lengths greater than fifteen feet. When swimming they propel themselves by the tail, the webs between the toes in many species being of more service in lifting them from the mud.

Voice in this group seems to be limited to a sort of suppressed bellow or blowing noise. Habits vary in different genera; some are nocturnal, others diurnal. In regions affected by protracted drouths some bury themselves in the mud before it hardens, and reappear when the rains have again softened it sufficiently to allow them to release themselves. Some are very dangerous. Natives in crocodile-infested regions are able to free themselves when seized by the creatures by thrusting the fingers into the eyes. Few animals are harder to kill. A hook baited with flesh and tied to a number of small cords, which get between the teeth, is the best means of capture. Caught in this way the powerful blows of the tail are effectively used in defense.

The common North American representative of the order is the Alligator, ranging from Florida to Tehuantepec. Stragglers have been reported as far north as the mouth of the Ohio river. A crocodile, *C. americanus*, ventures north along the coasts of Mexico from Central America. Another, probably *C. intermedius*, from the West Indies, is occasionally found in Florida.

SAURIA.

The transition from this order to the preceding and to the following is so gradual that it is next to impossible to cite characters that shall be general and exclusive. In a general way animals belonging to this order are distinguished from the Rhizodonts by dentition, squamation, structure of heart, structure of ear, transverse anal aperture, and male sexual organs, and from the Ophidians by lack of motion of maxilla and other bones of the head, elastic symphysis of mandibles, structure of ear, structure of eye, and presence of limbs. It will be seen at once that there are numerous exceptions, especially in regard to the separation from the serpents. Scolecophidia have the bones of the head more firmly united than other snakes, and some agree with the Amphisbenaë of the Sauria in the

possession of rudiments of limbs not visible externally. A few lizards have the ears covered by the skin, and others have no eyelids. There are serpents and lizards which have similar forms, motions, and habits. The extent of variation may best be seen by comparison of the examples cited below.

Development of the senses varies according to habits. In the serpent-tongued lizards we meet with great quickness of sight, hearing, scent and touch, accompanied by lack of taste—the tongue simply being a very sensitive tactile organ. Again, in species living upon vegetation, the touch is deficient and the tongue is short and thick, evidently an organ of taste. The keen-sighted *Chamaeleon* has a tongue which is probably an organ of taste as well as touch.

Many of the Saurians are noted for transient variations of color. The cause of these changes is found in the presence of pigments of different colors at various depths below the surface of the skin. The expansion or contraction of one or more of the layers, in consequence of muscular action, nervous irritation, or contraction or inflation of the lungs, changes the proportion of the different pigments visible at a particular instant. If the upper layer contains the dark pigment, the contraction of its chromatophores lessens the visible amount of this color and exposes a greater amount of that beneath it. These changes are not to be confounded with that occurring on soils of light color, or in regions where there is great reflection of the rays of light; in such localities permanent lightening of the colors is apparently due to a bleaching process undergone by the pigments. It seems as if the effect of polarized light upon the pigment differed from that of the direct unpolarized rays. Commonly each vertebra is concave in front and convex behind. To this there is a very marked exception in the case of the RHYNCHOCEPHALIA, a suborder founded on a family represented by a single genus of New Zealand lizard, *Hatteria* or *Sphenodon*. Externally this animal resembles the species of the European genus *Lacerta* so much that one can hardly believe it more than generically distinct. Skeletal structure and details of anatomy discover differences of sufficient importance to warrant the establishment of a different order. Most noticeable of its peculiarities are the series of palatine teeth, the structure of the skull, the biconcave vertebrae, the presence of a cartilaginous rib beneath each transverse fold of the skin of the abdomen, and the absence on the male of intromittent sexual organs. Of those examined the females were lighter in color, and

possessed much lower crests than the males. Specimens from the Ru Rima Rocks are smaller and darker than others from the main island. The stomachs contained nothing but insects.

Heloderma, one of the largest North American lizards of the Sauria proper, inhabits the arid region extending from Utah to Tehuantepec. It is a clumsy, slow-motioned creature, and presents a repulsive appearance. The skin is covered with transverse series of thick plates, rounded and separated somewhat on the back, quadrangular and close together on the ventral surface. As if better protected from below, the *Heloderma* is said to turn himself on his back when attacked. The teeth are long, slender, sharp, and grooved. The saliva is very irritating when introduced into a wound, as is almost certain to be the case when the animal is enraged. It is generally considered to be fatal to the smaller animals. These are probably the only venomous of the Saurians. They are terrestrial and carnivorous; not at all particular as to kind and condition of food. Two species of the genus are all that are known. By some authors they have been placed in the *Varanidae*; for others they form the family *Helodermidae*, which disposition is to be preferred.

The *Varanidae* include the largest lizards of the old world. They are elongate and slender in build, and live near the water. The nostrils can be closed by valves, and are provided with air pouches, arrangements which greatly favor diving and remaining below the surface.

True *Chamaeleonidae* are not found in America. The home of the family is Africa and Madagascar. One or two species have found their way northward in Southern Asia and Europe. That wrongly called by the name in the Southern United States is an *Anolis*. Chamaeleons have compressed bodies, short necks, and prehensile tails. The head is angular, often crested or provided with one or more proboscis-like processes in front. The skin is covered with granular folds or scales. The tongue is long, slender, and very extensible; it has a club-shaped extremity, prehensile and viscous in front. The eyes are large, globular, very mobile, covered by a lid through the center of which there is a narrow opening. A Chamaeleon is able to watch an object ahead of him with one eye while closely examining with the other something that has attracted his attention in the opposite direction. The tympanum is covered by the skin, but as the latter is exceedingly sensitive to irritation of any kind, it is possible the hearing is not greatly interfered with. The limbs are slender, compressed, and each bears five toes disposed

in two opposable groups. Such an arrangement of the toes, with the prehensile tail, enables the animals to walk lengthwise of the small branches with readiness. They spend their lives in the trees, and feed mainly upon insects, capturing them with the tongue, which can be thrust forward several inches for the purpose. The females lay from eight to a dozen eggs under the fallen leaves. In this group the transient variations of color are excessive in amount and rapidity; they often differ on opposite sides. Ability to take on at will the color of any object upon which an individual may be placed does not exist.

Most of the *Gecconidae* have rudimentary eyelids, and the eyeball covered by a transparent membrane under which it moves with freedom. A few have connivent lids. The pupil is most often oblong and erect. The tongue is short and thick. The skin is covered with granular or tubercular prominences, which are not imbricated. In the greater number of the species the feet are provided with adhesive apparatus under the toes in the shape of expansions or transverse series of plates, with which they are able to cling to vertical and smooth surfaces. These disks vary greatly in the different genera. Sometimes there are no disks, and sometimes the claws are absent. Occasionally the claws are retractile as in cats. The body and head are commonly depressed. When broken or lost, the tail grows out again; it may be reproduced a number of times in the life of the individual. This organ takes on fantastic shapes in some species; in all it is very fragile. *Ptychozoon* is marked by fringed dermal expansions on sides of tail, body, and head, which form a sort of parachute, answering a similar purpose when leaping to that of the membranes of *Draco*. The name *Gecko* is given in imitation of the voice. Geckoes live in the tropics of both hemispheres. Some frequent houses, where they are very useful on account of their insectivorous habits. In the United States a single species is represented, *Sphaeriodactylus notatus* Bd., at Key West, Florida. Three or four others, belonging to *Coleonyx*, *Diplodactylus* and *Phyllodactylus*, are reported from Sonora and Lower California. Farther South they are more common.

The *Agamidae* belong to the Eastern Hemisphere. In this family the eye and eyelid are well developed, the teeth are generally planted upon the upper edges of the bones of the jaws, the tongue is thick, and slightly, or not at all, extensile, the scales are imbricate, and the tail not nearly so fragile as in other Saurians. The toes are without disks. Of the odd forms in the various subfamilies, probably none is more striking than that of the

Dragons or flying lizards, *Draco*. These possess a membranous expansion on each side of the body, supported by the posterior or false ribs, which are much lengthened for the purpose. This membrane may be folded back against the body or expanded to serve as a parachute, as in case of the flying squirrel. The ordinary mode of progression is by means of short jumps. A dermal expansion or goitre depends from the throat; it is distended when in use by the hyoid apparatus. The tail is long and slender. All the species of Dragons are small; the total length seldom exceeds seven or eight inches. *Lyriocephalus* has its ears hidden under the skin, and its snout surmounted by a globular prominence. *Ceratophora* has a lobe or conical pointed horn above the tip of the snout. *Moloch*, an Australian lizard, literally bristles with spines. *Chlamydosaurus*, also Australian, has broad fan-like membranes extending backward from the sides of the back of the head.

The *Iguanidæ* belong almost entirely to the new world. They are diurnal. Their eyelids are valvular, and their tongues thick and slightly notched. This is one of the families in which forms and habits are most diversified, as will be seen by the instances cited below. It contains some of the largest species of the order. Among these are the terrestrial cactus-eating *Conolophus* and the amphibious *Amblyrhynchus*, which feeds on sea weeds and when pursued secretes itself beneath the rocks under the water, genera inhabiting only the Galapagos Archipelago. *Metapoceros*, the Navassa lizard, found also in Hayti, is very large and strong. To aid in burrowing, older specimens have peculiar scrapers under the basal joints of the third and fourth toes, and the second joint of the third toe of the hind feet. These scrapers are on the front side of the toes as they are pushed backward; they are merely expanded and solidified scales, the lower extremities of which form a sharp serrated edge of five or more denticulations arranged in a curve, convex forward. In *Thysanodactylus* the scales along the sides of the toes are so much expanded as to answer the purpose of a web in swimming. *Iguana*, though venturing north into Mexico, is properly a South American genus. It is also a common lizard in the West Indies. The species are large, rivaling the Galapagos lizards in bulk but much longer. Their flesh and eggs are much prized by the natives of the localities in which they are found. Where they have not been disturbed they are very tame and enjoy sunning themselves in pleasant weather extended along leafless branches of trees. They are hunted with dogs, and taken from the tree

with the noose, the arrow, or the gun. The range of *Basiliscus* also extends northward into Southern Mexico. Crested on head, back, and tail, slender and spider-like in build, in novelty of appearance it rivals the *Chlamydosaurus*. In some genera, as *Polychrus* and *Sphaerops*, the skin is to such an extent unprotected by the scales or granulations as to allow the transient color variations to be very noticeable.

Nine or ten genera of the family are found north of Central Mexico. Only one, in a single species, *Sceloporus undulatus*, occurs east of the Mississippi. The undulated lizard ranges northward from the Gulf as far as Southern Illinois and Maryland. It is common in parts of Kentucky and Virginia. *Phrynosoma*, a viviparous genus, ranges from Mexico to Dakota; it contains nine species, none of which appear East of the plains. One, *P. douglassii*, is common in Southern Dakota. Early in August the female gives birth to about eight young ones. Ants appear to form their principal food, though they are not by any means limited to them. They are vulgarly called "Horned Toads." The subfamily *Anoliidae* is represented in the Southern States by a common Saurian, the little greenish, flat-crowned, slender-tailed, goitred lizard of the trees and bushes, sometimes called "Chamaeleon" or "Scorpion."

The *Lacertidae* belong to the old world. They are diurnal land lizards, with polygonal symmetrically-arranged shields on the head, slender, forked exertile tongues, eyes provided with lids, nonimbricate scales, four limbs, and long fragile tails. They are the common lizards of Europe. The genus *Zootoca* is viviparous.

The *Teiidae* are Saurians of the Western Hemisphere. Their heads are covered with angular shields; and their tongues are free, notched, and exertile. The only genus representing the family in the United States contains the slender, short-limbed, six-lined lizard *Cnemidophorus sexlineatus*, of the South. Species of this genus are more numerous in the Rocky Mountain district, from Montana to Mexico. *Teius* contains several large South American species; they are the "Tupinambis" of the Amazon basin. A young individual has the teeth fixed on the inside of the jaw bone, but as he grows older the bone surrounds them more and more.

The Scincs, *Scincidae*, generally have fusiform or subcylindrical bodies, and are covered with glossy scales, which are either smooth, keeled, or grooved. The limbs are short, the body and tail rather long. In some there are no limbs, *Anguis*; in others there are two, *Ophioles*; and in the

majority there are four. The eyelids are connivent. Transparent lower lids occur in species of *Eumeces*. The tongue is slender, exsertile, and provided with a pair of pointed extremities. In habits the Scines are terrestrial. They secrete themselves under logs, bark, rocks, leaves, or in shallow burrows in loose earth or sand. Their eggs number ten or a dozen, and are laid in these situations. East of the Mississippi *Eumeces fasciatus*, the "Blue-tail," is the most common; it is found as far North as Illinois and New York. Specimens ten inches in length are very large. A second species, *E. anthracinus*, is found in the mountains of Pennsylvania and Southward. Two others have been described from Florida. Westward to Mexico the number is much increased. *E. leptogrammus* is taken in Dakota. A species of another genus, *Oligosoma laterale*, has a distribution somewhat similar to that of *E. fasciatus*, probably not extending quite so far North. The family is found in all tropical and subtropical countries. *Trachydosaurus*, the "Stump-tail," and *Cyclodus*, Australian genera, are of the largest.

A Californian genus, *Aniella*, furnishes a foundation for the family *Aniellidæ*. This lizard has a long snake-like body and tail, and is without limbs.

The Acontias, *Acontiadæ*, are from the Eastern Hemisphere. *Acontias* has no limbs, and the upper eyelid is rudimentary. *Ecesia* has short limbs, and the toes are not separate. *Nessia* has only three toes to the foot.

One of the common lizards of Southern California and Mexico, *Gerrophonotus*, belongs to the *Zonuridæ*, a family of which the greater portion belongs to the old world, and which is specially marked by a distinct longitudinal fold or groove along each flank. Another member of this family is the footless snake-lizard, *Ophisaurus*. The latter has a long, slender tail, which is easily broken, and being longer than the body, more than half the total length can be carried away without disabling the animal, which, by a second growth, soon replaces the portion lost. It is to this peculiar genus that we owe the fiction of the "Glass-snake." *Pseudopus*, an allied genus of Europe and Southern Asia, resembles the preceding, but has on each side of the vent a small limb, on which the toes are not separated.

The AMPHISBÆNIA form a very distinct suborder of the Sauria. In the typical forms the body is long and subcylindrical, and the tail short and thick. The bones of the skull are firmly articulated, and the symphysis of the lower jaw is nonelastic. The tongue is flat, thick, and notched at the end; the eyes are small and covered by the skin; the ears are hidden; and

the surface of the skin is divided into small rectangular spaces by transverse folds traversed by longitudinal grooves or furrows. Limbs are absent, or only present in front. *Chirotos*, a Mexican genus, has the fore limbs; the other genera are footless. Most of the species are said to be found in or about the nests of ants. One species, *Lepidosternon floridanum* Bd., inhabits the Southeastern United States.

OPHIDIA.—Serpents.

The elongate spindle-shaped or subcylindrical form of the snake is a familiar one. There are species which present no distinction between head, neck, and tail, and there are others in which these parts are well marked. The majority have no limbs; a few have rudimentary hind limbs, which appear as claws on each side of the vent. A sternum is lacking, but in the worm-snakes there are rudiments of a pelvic arch. Most of the serpents have teeth on jaws and palate. In consequence of the elastic articulations of the bones of the skull and the symphysis of the lower jaws, they are able to swallow animals considerably larger than would otherwise be possible. All the species of the order are carnivorous. Some crush their prey in the folds of the body, others kill the creatures on which they feed by venomous secretions, but the greater number swallow living food. During the operation of swallowing, small pores along the jaws give out a saliva which renders the act more easy. Frequently the snake disgorges a partly-swallowed meal to change its position in the mouth, or to take breath; the slimy appearance of such morsels has given rise to the erroneous idea that they have been deliberately covered with slime before the attempt to dine. Movable eyelids do not exist in the Ophidia. The epiderm is continuous over body and head, and forms a transparent cover for the eyeball. With the slough or cast skin the outer envelope of the eye is carried away; this happens at least once in a season.

Near the time of sloughing or moulting, the dermal covering of the eye is loosened, more opaque, and interferes, to some extent, with the sight; at such times snakes are said to be blind. Size of ball, shape of pupil, keenness of sight, etc., vary according to the habits of the species. Hearing is dull. There are no external evidences of ears, and it is most likely the jar felt over the whole body from an approaching tread often gives the needful warning. The hissing sound produced by expelling the breath from the lungs has been supposed to be the nearest approach to a voice. Lately, however, some

Yellow Boas, *Chilabothrus inornatus*, in my possession have been repeatedly heard to give utterance to a low plaintive whine as the breath has been forcibly emitted. Apparently it is by the sense of smell that the snake finds its companion. About the time of coupling many species are possessed of a powerful and very disagreeable odor. The sense of taste is probably lacking. The tongue is a tactile organ; it is soft, slender, provided with a pair of flexible tips, and can be retracted into a sheath at the bottom of the mouth. As serpents move about they are constantly feeling ahead of them with the tongue, and the forward thrust and peculiar appearance of this organ has given rise to the false idea that with it the "stinging" is done. The stomach is formed by a widening of the alimentary canal; its sides are thicker than those of the œsophagus, and have longitudinal folds. Digestion is quick or slow according to the temperature; venom hastens the process. One lung is often rudimentary; in species of *Boa*, *Naja*, and *Crotalus*, both are developed; in *Boa* they are about equal in size. The ovaries and testes are paired, the right often larger and placed a little in advance of the left. The male is furnished with a pair of intromittent organs, one of which is placed on each side of the vent under the base of the tail. They are tube-like, and bear a groove on the side; when in use they are everted like the finger of a glove, and the groove becomes an external furrow by which the seminal liquid is conducted into the oviduct. Careless observers have mistaken these organs for feet. In certain species their extremities are surrounded by series of strong, sharp spines or hooks. The eggs are oblong, and have a soft, leathery envelope, for the rupture of which in hatching the young are provided with an egg-tooth. Oviparous serpents generally leave the eggs to hatch and care for themselves; the Pythons or rock snakes of the Eastern Hemisphere are exceptions to this rule; after the eggs are laid the female coils her body round them and incubates. Viviparous species are those in which the eggs are hatched in the oviduct; there are those in which hatching and laying happen so nearly at the same time that they are at times oviparous and at others apparently viviparous. The ribs are very numerous, in some species numbering hundreds, and are loosely articulated to the vertebrae. They furnish the main dependence in locomotion. In reality, there are three methods of progression used by Ophidia, and each of these may be employed separately. When a serpent glides he brings the lower ends of opposite ribs forward

at the same instant. Each step is short, hardly equal the length of one of the ventral scutes. The ribs are attached to the scute in such a way that its free posterior edge, braced against projections on the surface over which the animal is moving, serves as a support from which the body is pushed ahead. A track left by a gliding snake may be a right line. This motion can be executed so that the entire body moves along without apparent effort. In walking opposed ribs are moved forward alternately, and the edge of the end of the scute under the rib moved holds what is gained by the motion while the opposite extremity is stepped ahead of it. This mode of traveling gives rise to flexures of the body, and the track is undulating. By the third method, the creature pushes or elbows its way with the body and tail, as do the eels or snake-like batrachians. Most of the worm snakes progress in this way. In fact, any snake hurriedly moving among grass, bushes, or rocks, takes advantage of any support he can get for his sides or extremity. Except when gliding, serpents in motion commonly use the three methods at the same time.

The outer dermal layers are generally folded into the semblance of scales; occasionally they are tubercular or granular. The scaly folds are most often imbricated on the body, and occasionally over the head. Generally labial and other plates of the head are non-imbricated. When the epiderm is to be shed, the snake rubs it loose at the lips, and, creeping against some object, manages to strip it off entire, and ordinarily inside out. The number of times the skin is shed in a season is variously stated by different authorities. Serpents that were kept for study have sloughed but once in a year, and I am inclined to believe this is the common practice. Nocturnal serpents are few, and belong to the torrid zone. Most, if not all, are affected by a period of comparative inaction; during the winter this takes the form of hibernation or a winter sleep. Such a sleep or rest is not an absolute necessity; when waked and kept lively every day of the winter they do not seem to be any the worse for it in the spring.

In coloration these reptiles are excessively varied; individual variation is great in the same species; the colors are more bleached in exposed situations; and of terrestrial species, those portions of the body that are habitually raised from the ground are commonly whiter beneath than the balance of the ventral surface.

Certain good authorities have adopted the idea that some snakes swallow their young for protection in times of danger. The evidence hardly appears

sufficient to warrant the belief. Instances reported to the writer concerned some known to be snake-eaters; others known to lay eggs, paying no farther attention to them, and others of undeterminable species noticed by incompetent observers.

The Museum of Comparative Zoölogy possesses a specimen of *Coluber constrictor*, the "Black Snake," taken in Essex county, Massachusetts, by Prof. Putnam, the stomach of which contained a half-digested specimen of *Cyclophis vernalis*, the "Green Snake," and another of *Storeria dekayi*, DeKay's "Brown Snake." It is a well established fact that this snake and others feed upon their own species at times. Specimens of viviparous species, in which the young were about ready for delivery, have led some to the mistaken belief that the little ones had been swallowed.

The classification adopted in the following synopsis includes all serpents in four groups or suborders: SCOLECOPHIDIA, worm snakes; ONYCHOPHIDIA, clawed snakes; ACACOPHIDIA harmless snakes; TOXICOPHIDIA, venomous snakes.

SCOLECOPHIDIA.

Those of the first suborder are subcylindrical, wormlike, burrowing species, which feed upon worms, larvæ of insects, and the like, living beneath the surface of the ground. The body is usually elongate, and the tail short and thick. On account of their subterranean habits, the eyes are protected by thickened plates, and are more or less imperfectly developed. Probably it is by the sense of smell that the presence of food is detected. The mouth is comparatively small, and the bones of the skull are more firmly articulated than in other Ophidians. There is great similarity between the scales of the back and those of the ventral surface. The body is marked by a certain rigidity, necessary in burrowing, which makes these snakes very awkward travelers upon the surface of the ground. As if to aid in pushing, the tail is in cases provided with a sharp spine, and in others it is truncate and tubercular. Venomous Scolecophidia are unknown. North American representatives of the group are found in ten or eleven species of the family *Typhlopidae*. They range from California and Texas southward and westward. Worm snakes belong to tropical and subtropical regions of both hemispheres. Certain species possess a rudimentary pelvis.

ONYCHOPHIDIA.

Serpents of the second suborder, *Onychophidia*, are distinguished by rudimentary posterior limbs, the only external evidence of which is a small claw or prominence on each side of the vent. This group contains the largest of the snakes, the Boas and Anacondas of the Americas and the incubating Pythons of the old world, species of which are said to reach a length greater than thirty feet. The section of the group containing species with thick non-prehensile tails is represented in California and Mexico by several burrowing snakes belonging to the genus *Charina* of the *Erycidae*. *Eunectes*, containing the Anacondas, is a South American genus, and its species reach greater size than other new world snakes; it belongs to the section with prehensile tails. Of this section there are three species of *Boa* in Mexico; a doubtful fourth is reported in the Yellow Boa, *Chilabothrus inornatus*, properly of the West Indies. These Onychophidia are expert climbers, and many are equally active in the water. None of them are venomous. They are able to swallow animals of much greater diameter than their own. Ordinarily they seize the prey with the teeth and crush it in folds of the body.

ACACOPHIDIA.

The innocuous snakes comprise all the common, harmless species of the ground, fresh water, or trees. Being not at all harmful, and feeding on mice, moles, insects, etc., many of them are important aids in restraining the increase of the most destructive enemies of the husbandman. The vulgar fear of these animals is mainly the result of education or of prejudice, which lacks foundation in facts. In reality, some of the most dreaded have no existence. Such as the "Ring" or "Hoop Snake," which is said to take the end of its tail in its mouth and roll over and over like a hoop, killing everything it touches with its venomous horn, and the "Blow Snake," the breath of which is deadly, are mere fictions. A few members of the division prey upon others, and are said to conquer and eat species of the Toxicophidia.

Several are possessed of considerable curiosity, and sometimes follow objects that have attracted their attention. The common black snake, *Coluber constrictor*, occasionally chases men in this way, but if turned upon at once seeks safety in flight. It happens at times that harmless species

put themselves in the attitude of defense, and resent attack when unable to escape; their bites are mere scratches, and should occasion no anxiety.

Although typical earth, water, or tree snakes are very distinct, the intermediates are so numerous as to make it impossible to classify them according to habits. It is well, however, to note the differences of the types, without losing sight of the fact that there are forms of such structure and habits as to make their position doubtful in either of the three divisions.

Ground Snakes are rather stout and thick in body and tail. Their bodies are usually cylindrical, and the tails most often short and conical. For the most part they have smooth scales. The eye is of moderate size to small, and the pupil generally round. Certain species which burrow, as those of *Heterodon*, have valvular nostrils, and rostrals shaped somewhat like shovels or plows.

Water Snakes are well represented in the genus *Nerodia*, in which the body is stout and fusiform, the tail tapering, the nostrils valvular and near the top of the snout, and the scales keeled. One of the most peculiar forms of the division is *Herpeton*, which has a pair of feelers or tentacles extending forward from the muzzle.

Tree Snakes are very long and slender. Their eyes are comparatively large, and the pupil is either round or oblong, erect or horizontal. Tactile appendages to the snout are possessed by some species. In general they feed upon the small animals that make their homes among the leaves and branches. *Rachiodon*, an African genus, feeds on eggs; it has very weak teeth in the jaws, but the inferior processes of the vertebrae of the neck extend into the œsophagus, and are tipped with enamel for the purpose of crushing the shells after so far swallowed that the contents may not escape by the mouth and be wasted. Green is a very common color in this division.

In the list of families in this suborder north of Tehuantepec, several really belong to South America, being introduced on account of straggling species reported from the Mexican region. Of the *Dipsadinae*, a single species ranges north of Mexico into Arizona and Texas. From Southern Mexico two species of *Scytalinae* have been noted. There are two species of *Dendrophinae*, which also range into Mexico. The most abundant North American snakes belong to the *Natricinae*, of which species, as *Eutania sirtalis*, range over the whole continent, excepting the far North. The

genera *Eutania*, *Nerodia*, *Regina*, and *Storeria*, are North American; *Helicops* and *Hydrops* are better represented in Central and South America.

Although the family names in common use are retained here, they are ranked as subfamilies of one great family, including all the Acacophidia or Colubrine snakes, the *Colubridæ*. The family *Colubrinæ* contains in the genus *Coluber* several species very common in the United States: the Black snake, *C. constrictor*, is found over a great portion of the continent, and the section of the genus containing the Coach Whip snake (*Masticophis*), *C. flagelliformis*, includes half a dozen species, from the Southern States, Mexico, and as far south as Chili. Two species of the South American genus *Spilotes* are found in the Gulf States. *Pityophis*, *Cyclophis*, *Phyllophilophis*, and *Salvadora* are North American. As the name *Salvadora* does not seem to be used elsewhere in the animal kingdom, it has been retained. *Dromicus* is South American and West Indian, but contains one species peculiar to the Southeastern United States and several others found in Southern Mexico. The old world genus *Elaphis* claims several widely distributed species from the southern half of the continent.

Tachymenis, *Erythrolamprus*, *Liophis*, and *Xenodon* are South American *Coronellinæ*, and are represented by a few species in the southern part of Mexico. *Ophibolus*, *Diadophis*, *Heterodon*, *Rhinocheilus*, *Cemophora*, and *Hypsiglena* are classed as North American; the first three are found in all the States, with, perhaps, a few exceptions in the northern tier.

The *Calamarinæ*, of the genera *Tantilla*, *Contia*, *Sonora*, *Virginia*, and *Carpophis* are Northern; *Ficimia*, *Cheilorhina*, *Stenorhina*, *Ninia*, *Geophis*, and *Elapomorphus* are shared by Mexico with South America.

Tree snakes are found in the *Dipsadinæ*, *Scytalinæ*, *Dendrophinæ*, and *Colubrinæ*; water snakes in the *Natricinæ*; and ground snakes in the *Calamarinæ*, *Coronellinæ*, and *Colubrinæ*. But in each family there are species whose habits are so uncertain as to render their positions doubtful.

TOXICOPHIDIA.

Grouping all the venomous serpents in a single suborder, brings together animals of considerable diversity. Because of their differences they have been arranged in various sections: one, the *Proteroglypha*, in which the fangs are erect, immovable, and grooved in front; and another, the *Solenoglypha*, which are distinguished by an erectile perforate fang. The Pro-

teroglypha are subdivided still farther into the *Conocerca*, the conical tailed or colubriiform, and the *Platyicerca*, flat-tailed or sea snakes. The Solenoglypha are also subdivided into *Abothropha*, pitless vipers, and *Bothropha*, which are marked by a pit on the side of the face between the eye and nostril.

The cosmopolitan genus *Elaps* contains all the North American species of *Conocerca*. There are two species in the Southern States, and a half dozen or more in Mexico. Our species are not considered dangerous. Among the old world species of the division are some of the most deadly of the order, such as the Cobras, *Naja*, of Southern Asia and Northern Africa, the favorites of the Indian jugglers. As in other sections, there are species which feed upon other snakes.

The *Sea Snakes*, *Platyicerca*, have the nostrils on the top of the snout, and the tail compressed so as to form a paddle. Three to five feet in length is the common size; ten feet is a great length for the largest. They are numerous in parts of the Indian Ocean. As far as known, a single species has been found in fresh water, *Hydrophis semperi*, from Lake Taal in the Philippines. *Pelamys bicolor* has been taken off the coasts of Southern Mexico. Sea snakes are said to approach the shores only when about to give birth to the young. They feed upon marine animals of various kinds, and are sometimes taken in the nets of the fishermen, who fear them very little.

The *Abothropha* belong to the Eastern Hemisphere. Serpents of this division differ from those of the next principally in the absence of the pit or lachrymal fossa. With this exception, structure, habits, effects of venom, etc., are similar.

The pit vipers, *Bothropha*, are the most venomous of the new world Ophidia. A single family, *Crotalidæ*, has heretofore included all of them. As in the pitless vipers, the fangs are the only teeth upon the maxillaries. The latter are short, and rotate upon the lachrymal and frontal bones in such a manner as to allow the fangs to recline against the roof of the mouth, where they are covered by an elastic membrane when not in use. The fangs are replaced when shed or broken by others, which grow in a reclining position behind those in function. A section of one of these fangs indicates that the tooth has been folded from its sides forward and closed around the groove which opens a short distance above the point. Another opening at the base is opposed to the extremity of the duct from the venom

gland at the side of the skull behind the eye. This gland is hollow, and contains the venom it secretes. Venom glands vary considerably in size and position; they extend back upon the neck in certain Conocereæ. Compression by means of the muscles upon the gland forces its contents through the duct and tooth. This compression is entirely independent of opening the mouth or erecting the fangs. At the will of the serpent the venom may or may not be delivered with the bite. As if the creature was economical, and did not wish to waste its supply, it is frequently the case that it strikes one or more times without using the poison, and is able to strike a more effective blow when more thoroughly exasperated. If an angry snake is held up by the neck, and its mouth kept open, it will raise and lower the fangs from time to time, and occasionally spurt the venom through them with considerable force. Usually on being disturbed and unable to retreat, the snake throws himself into an attitude of defence by coiling his body so that the upper half can be readily straightened in the act of striking. A Rattlesnake will place the tail in the coil with the rattle directed upward. Very angry individuals strike whether coiled or not. They do not jump; the hinder part of the body remains in position, and none of our serpents are in the habit of reaching more than half of their length. If a stroke is successful, and the fangs enter the flesh of the victim, a backward pull throws the weight of the body on the fangs, tending to make free a space in front of them for the introduction of the venom. A blow is sometimes given when the fangs are not erected, sometimes with the mouth wide open, and sometimes, apparently, with the jaws closed, the fangs passing on the outside of the lower jaw which then serves as a lever with which to disengage them. After wounding it fatally, the snake watches the prey very intently until the venom has produced the desired effect and the victim falls, when it appears as if an estimate of the size was made with chin and tongue by passing from one end to the other of the body before the attempt to swallow begins at its nose. This intent gazing at squirrels, birds, etc., in their dying agonies, has given rise to the erroneous ideas concerning fascination. All there is of fascination or charming is to be accounted for by the fact just mentioned, together with the quiet, almost imperceptible motion of the snake and the recklessness and terrorism of the prey. Venom hastens decomposition, and thus renders digestion more rapid.

In general the venomous snakes of North America are much less dangerous than is supposed. What is fatal to the small animals which form their food becomes much less troublesome to larger ones. The degree of danger to men varies according to the size of the snake, the number of blood vessels in the part bitten, and the condition of the system at the time. There is a great variation in the amount of venom injected. After it has bitten several times, the supply of venom is reduced, so that a bite from a large specimen is not attended by serious consequences. The immediate effects of the poison are a terrible pain and rapid swelling about the bitten part, followed by general nervous prostration or exhaustion, retarded respiration, and enfeebled action of the heart, accompanied by vomiting, perspiration, and partial loss of control of the muscles. When not fatal, recovery is very rapid, usually being complete in about three days. A peculiar swelling on the jaws remains for a long time in cattle that have been bitten. Fatal cases in which men have been victims probably do not number as many as one in twenty. The remedies are cutting out the bite, burning with red-hot iron, caustic or nitric acid, suction by the mouth or cups, ligatures to prevent the rapid dispersion of the poison through the system, ammonia used as a wash and taken internally, and alcoholic stimulants. If there are no cuts or sores in the lips or mouth, the poison can be drawn out by suction; it is only when introduced into the blood that it is effective. On receiving a bite, the first thing to be done is to suck out as much of the poison as is possible, enlarging the wound with a knife for the purpose. At once a bandage should be put on so as to keep the venom from the rest of the body as long as possible. For a bite on the finger, it should be tied between the bite and the hand; another ligature should be placed around the wrist, and possibly a third around the arm; these should be loosened gradually and for an instant at a time, the aim being to control the dispersion, and introduce the dangerous matter into the circulation little by little. It is well to cauterize immediately if possible. Washing the wound with ammonia is recommended, as also drinking a dilute mixture of ten or fifteen drops in a glass of water. Alcoholic stimulants are undoubtedly beneficial if not taken to excess, in which case they become sedatives, and assist rather than counteract the poison.

The rattle of the Rattlesnake is formed of horny rings or shells, of which the first is firmly attached to the end of the tail, and each of the others loosely grasps that in front of it, and is itself grasped by the next in order.

It is believed that one joint is added to the number each year; this is but approximately true; there are variations connected with the different ages and rates of growth, the extent of which have not been fully determined. The joints are so easily worn out and broken that it is seldom the case that a large snake has the complete series.

The probable use of the organ has given rise to a great deal of discussion. Because of the resemblance of the sound to the crepitation of the grasshopper, and to the shrilling of the Cicada or harvest fly, though much less rapid in its vibrations than the latter, it is claimed that it draws insect-eating birds within reach. This is possible; but the fact that the snakes are usually so quiet when feeding, and that birds are so rarely found in their stomachs, bears against the supposition. The idea that the rattle produces terrorism in the prey is hardly tenable, since it appears to be least used when feeding, and frightens the game away. The rattles are used in time of coupling, but not with the energy marking their use in anger. In reality, the attitude of the serpent is defensive at all times, unless it be when seeking food; it prefers to economize venom and take no risk of losing fangs; it does not wish to strike unless driven to do so; and the rattling is a threat or warning that it is ready if it must. In economizing venom and fangs the rattle is undoubtedly beneficial, and prevents a great many useless strokes; at the same time its action is disadvantageous in that it calls the attention of enemies to its owner. A snake appears to realize how much its subsistence depends on an ever ready supply of venom and perfect fangs, and how helpless it becomes without them.

It is often asserted that deer destroy a great many snakes. On what foundation the assertion rests I do not know. Hogs are certainly fond of them, and large fowl are especially destructive to the young.

Ancistrodon, Copperheads and Moccasins, and *Crotalus*, Rattlesnakes, are about equally to be feared. These genera properly belong to North America. Five species of Rattlesnakes are found East of the Mississippi.

Certain species having plates on the crown resembling those of the Colubers have been given the generic names *Crotalophorus*, *Caudisona*, and *Crotalus* by different authors. These names were synonymous at a time when the species of which we speak were unknown. They were originally applied to species which have not been separated from the genus as first constituted. In view of this, it seems best to regard them as synonymous, and to give to the species with the shields a name less liable to confuse on

account of previous association. The name proposed for the species grouped under *Crotalophorus* of Gray, Holbrook, and others, but not of Linné or Gronow, is *Sistrurus* (*σειστρον*, a rattle).

BATRACHIA.

The Batrachians are vertebrates, such as the Cæcilians, Salamanders, Toads, and Frogs, the most of which are affected by a metamorphosis, during which a branchial and fish-like respiration is changed to one in which the main dependence is placed on lungs. Whether breathing by means of gills in the earlier stages, or by lungs in the later, these animals are possessed of a supplementary cutaneous respiration. It is by the aid of the latter that they are enabled to pass periods of several months to a couple of years buried in the earth or mud during hibernation. Moisture is at all times absolutely necessary; deprived of it they soon die. The skin is naked, either smooth or rough. In cases the skin over the whole body is glandular; in others the cutaneous glands are aggregated in particular portions of the body, as the parotoids of Toads and some Salamanders. A few of the snake-like batrachians have rudimentary scales hidden in the skin. The slough is stripped off more or less entire, and generally eaten. The gills are retained by certain forms throughout their whole existence; in these, however, the lungs are partially developed, and the respiration is at the same time branchial and pulmonary. The heart has but a single ventricle, and the atria are incompletely separated. There are two occipital condyles. For the most part, the very young feed on vegetation; in such the intestine is elongate. Those beginning their independent existence after the tadpole stage has been passed, and the later stages of all members of the class, are carnivorous. The prey is swallowed entire. There are a few instances in which the embryo is developed and the young hatched in the oviduct. Commonly the eggs are fertilized externally, as or after they are laid. Eggs of batrachia have been favorites of embryologists in their researches; it was in them Prevost and Dumas first noticed the cleavage masses, and in them Newport saw the spermatozoon creep through the outer envelopes to the yolk. We know no species of which the bite is poisonous. The nearest approach to venom is in the acrid, milky secretion of the parotoids, which produces considerable irritation when brought in contact with the membranes of eyes or mouth.

Young specimens reproduce lost portions of the body with great readiness. The flesh of many species is good for food.

This class contains three orders. To the first or footless order the name *Apoda* is applied; another title for the same has been *Cecilia*. To the second, in which feet and tail are present, the name *Urodela* is given; this has also been called *Ichthyodi*. And the third, having feet but lacking a tail in the adult forms, is known by the name *Anura*.

The APODA are snake-like or vermiform. Some of them are very long and slender, others short and thick. As the name indicates, they are without feet. The tail is short. The skin is smooth, slimy, and arranged in transverse folds, between which rudimentary scales are sometimes found. Usually but a single lung is developed. As might be expected from their habits, burrowing in the ground to feed on worms, insects and the like, the sight is somewhat imperfect, the eyes being partially hidden under the skin. The young resemble the adults; the metamorphosis is slight. Apoda are found between the tropics of both old and new worlds. The only North American species yet found were taken in southern Mexico.

In URODELA limbs and tail are present. The number of limbs varies. When there are but two, as in *Siren*, the hinder are lacking. The earlier or larval stages resemble the fishes in means of breathing and progressing. *Siren*, *Proteus*, and *Necturus* have persistent branchiæ, and their lungs remain rudimentary. The majority develop lungs and lose the gills. After the gills are resorbed, the branchial openings close in most cases; a few, as *Menopoma*,* retain a small opening through life. Very young larvæ have a cutaneous gular flap, free at the hinder margin, which extends backward over the branchial arches and the isthmus between them; it unites first with the skin of the sternal region, and later, as the gills disappear, with that of the shoulders. The process is similar in larvæ of those Anura in which one side is entirely closed and a small passage is left for the passage of the water on the other. Many species spend the greater portions of their lives in damp localities on the land rather than in the water. Those preferring the land are marked by greater roundness of the tail, the aquatic forms having this organ compressed and expanded. There are teeth on both jaws, and most often on the palate. The tongue is pedicellate in some, has free margins in others, or is attached along the center in others. The toes are without claws.† The fore legs appear first.

* *Cryptobranchus*.

† Except *Onychodactylus*.

When first seen they are fleshy buds. These elongate, and the rounded extremity shows signs of dividing into two. One of the two becomes the inner finger; on the outside of the other the third finger pushes out, and beyond the latter toward the wrist the fourth. The hind limbs appear in similar manner. A single toe is formed of the inner lobe, the outer being one, or becoming two, three, or four according to the genus. In certain genera there is much variation in regard to number of digits; *Proteus* is a good instance; *Amphiuma* is another. Of twenty-six specimens of *Amphiuma tridactylum*, twenty have three toes, the normal number, to each foot, and among the others occur the following one has one toe on each hind foot; one has one variations: to the right hind foot and two to the left; two have single toes on the right fore foot; another has one, and still another has two on the left fore foot.

Salamandra atra is one of the cases in which the young are hatched in the oviduct. The eggs of some are fertilized in the oviduct during copulation; those of others are thrown free into the water to come in contact with the sperm emitted by the male at the same instant. According to accounts recently published by M. Gasco, the female Axolotl finds the spermatophore which has been deposited by the male, and placing the cloacal opening over it introduces a portion, and thus secures fertilization. It is generally conceded that the males are more numerous than the females of species of this order. One of the oddest forms is *Pleurodeles*, in which the ribs extend through the skin on the flank.

The *Urodela* belong to the temperate regions of the Northern hemisphere. Most of them are found in North America. The largest known is the *Megalobatrachus* of Japan. Of the North American, the largest is the aquatic genus *Menopoma*.*

The ANURA or tailless Batrachia have short bodies and short necks. The vertebræ are few, and in body or neck comparatively immovable. Generally the hind limbs are longer and stronger. There are four digits on the hand and five on the foot. Except in a few species of those living in trees, there are no webs between the fingers. *Rhacophorus* is the most marked exception; it has such extensive webs between fingers and toes as to be able by their aid to sail long distances in the manner of flying lizards or squirrels. Webs are common to the aquatic species, as in frogs, *Ranidæ*. Terrestrial frogs are more or less completely without the palmation. In *Cystignathidæ* the toes are free. *Pipa*, the Surinam Toad, one of the group *Aglossa*, or

* *Cryptobranchus*.

tongueless Anura, has star-shaped extremities to the fingers, which are probably sensitive tactile organs. *Dactylethra*, the clawed frog of Africa, belonging to the same group, has claws on its toes. Of the group possessing tongues, *Opisthoglossa*, those living in trees have fleshy bulbs or disks on the ends of the digits, by means of which they cling to smooth surfaces. The tree frogs, *Hylidæ*, exhibit this structure at its greatest development. Many species of the order have horny processes on the feet for burrowing. The males of some develop horny claws or tubercles on the inside of the thumbs or on the breast in the coupling season.

All of the *Anura* begin life as tadpoles, breathing and moving like fishes. They undergo a complete metamorphosis in taking on the form of the adult. Fertilization takes place externally, the male grasps the female tightly around the body, and the spermatic liquid is emitted as the eggs are extruded. Usually the males are smaller and more numerous. Habits vary much in regard to care of eggs. Our common Toads and Frogs leave them to care for themselves in the pools in which they are laid. The male of the obstetric Toad, *Alytes*, carries the eggs until they are about to hatch attached to his thighs, and then he enters the water. The male of the Surinam Toad, *Pipa*, places them carefully on the back of the female, and the skin rises around each, enclosing it in a little cup, within which the metamorphosis is effected. *Nototrema* carries the eggs in a pouch on the hinder part of the back. *Hylodes martinicensis* lays the eggs in the axils of leaves, and transforms so soon after hatching that it can hardly be said to have a tadpole stage. The larva of the Bull Frog, *Rana catesbyana*, remains a tadpole from one to several years. Many of the young feed upon vegetation for a time, afterward becoming carnivorous. M. Fernand Lataste has recently called attention to series of pectinate fringes or teeth outside of the horny lips of the tadpoles, and M. Héron-Royer finds so much variation in the various genera as to aid him in classification.

As the lungs and limbs develop, the intestine is shortened and the tail resorbed. The lower jaws are toothless.* As in toads, *Bufo*, some are entirely without teeth in the adult forms, and others, as the frogs, *Rana*, are in possession of teeth on upper jaws and on the roof of the mouth between the inner nostrils. The eyes are most often prominent and very keen. The inner eyelid is transparent. An East Indian genus, *Megalophrys*, and another from South America, *Ceratophrys*, have supraciliary


* Except in *Amphignathodontis* and *Hemiphraetide*.

prominences, which give them a fantastic appearance. The hearing is very quick. The tympanic membrane is generally exposed on the side of the head, and in certain species is very large. In *Opisthoglossa* the tongue is used in capturing the prey. Most often it is attached in front, and the hinder portion can be thrown forward out of the mouth; when not so free the whole tongue is pushed forward. The food consists of worms, insects, etc., and occasionally of small vertebrates. Several large species are said to catch fish. The latter, however, are well able to retaliate, and the batrachians furnish food for numbers of the fishes. A large number of the species have voice. A special apparatus for its production, possessed by the males, consists of a pair of membranous gular sacs, sometimes a single one, under the floor of the mouth, with which they communicate by a couple of slits. These sacs are either covered by the skin or, in a few instances, there are openings through which they are protruded when inflated with air from the lungs. The largest species of the order is the Marine Toad, *Bufo aqua*, of South America; the largest North American is the Bull Frog, *Rana catesbyana*. The number of species north of Mexico is about forty for each of the Anura and Urodela.

NORTH AMERICAN REPTILES.

PART I.

OPIHIDIA.—SERPENTS.



- | | |
|------------------------------------|-------------------|
| 1. Rostral. | 11. Parietals. |
| 2. Nasals. | 12. Occipitals. |
| 3. Loreal. | 13. Labials. |
| 4. Preoculars or Anteorbitals. | 14. Infralabials. |
| 5. Postoculars or Postorbitals. | 15. Gulars. |
| 6. Temporals. | 16. Mental. |
| 7. Internasals. | 17. Submentals. |
| 8. Prefrontals. | 18. Ventrals. |
| 9. Frontal. | 19. Dorsals. |
| 10. Supraillaries or Supraoculars. | 20. Tongue. |

SYNOPSES AND DESCRIPTIONS.

OPHIDIA.

Bronniart, 1800.

Vertebrate animals with elongate more or less cylindrical bodies, in which head, neck, trunk and tail have the same axis; blood aerated in lungs; bones of the head articulated by means of elastic ligament, which, in the jaws especially, admits of considerable distention; limbs absent or rudimentary; without a movable eyelid or external evidence of ears.

Not venomous;

no fold under the chin;

wormlike; head and tail not distinct from the trunk;

no claws at the side of the vent

SCOLECOPHIDIA.

a fold under the chin; head more or less distinct;

a claw at each side of the vent

ONYCHOPHIDIA.

without claws or venom glands

ACACOPHIDIA.

Venomous;

without claws; head more or less distinct;

with venom-secreting glands and fangs

TOXICOPHIDIA.

SCOLECOPHIDIA.

SCOLECOPHIDES.

Dumeril & Bibron, 1844.

Body elongate cylindrical; head short, indistinct; tail short, thick. Eyes imperfect, in some species invisible, covered by the ocular shields. Mouth small, inferior. Ligamentary attachments of the bones of the head possessing but little elasticity. No fold under the chin. Teeth few, on either the upper jaw or the lower, none on the palate. Tongue forked, exsertile. Scales smooth, imbricate, similar on back and belly.

Found under rocks or pieces of wood, or in the earth, where they burrow to feed on worms, larvæ of insects, etc.

TYPHLOPIDÆ.

TYPHLOPINÆ.

Teeth in the upper jaw

TYPHLOPS.

STENOSTOMINÆ.

Teeth in the lower jaw;

crown-shields resembling those of the
colubers ANOMALEPIS.

crown-shields scale-like

STENOSTOMA.

TYPHLOPIDÆ.

Fitzinger, 1823.

TYPHLOPINÆ.

TYPHLOPS.

Schneider, 1801.

Teeth on the upper jaw. Rostral shield well developed, reaching the upper part of the snout. Nasal vertical, divided or entire. Crown with scales similar to those of the back. Labials four, rarely three, anterior small. Hab. All tropical and subtropical regions.

TYPHLOPS LONGISSIMUS.

*OPHTHALMIDIUM LONGISSIMUM Dum. & Bibron, 1844, Exp. Gén. VI, 263.**TYPHLOPS LONGISSIMUS Jan, 1861, Arch. per la Zool. I, 182.*

Body long, slender, cylindrical; head depressed, rounded; tail short, rounded at the extremity, armed with a small spine. Rostral linguiform.

Nasal small, rectangular, half as wide as the rostral at the mouth. Preoculars twice as high as wide. Oculars smaller than the preoculars, similar in shape. Crown-shields small; one prefrontal, one frontal, two supraoculars, and two parietals. Scales small in 22 longitudinal rows; transverse rows 512.

Anterior portion of the head yellowish; body uniform ashy gray. Hab. North America. (From descr.)

TYPHLOPS PERDITUS.

Peters, 1869, Monatsberichte Berl. Akad., 435.

Eyes invisible. Nostrils at the end of the snout. Head shields similar to those of *T. reticulatus*, except that the posterior margin of the nasorostral is much less concave and the preocular less bent forward. Scales in 18 longitudinal rows. Dorsal scales olivaceous brown with yellowish-green margins. Beneath and on the end of the snout greenish-yellow. (From descr.) Said to be from Orizaba, Mexico. Total length 0^m, 285; tail 0^m, 004.

TYPHLOPS BASIMACULATUS.

Cope, 1866, Pr. Ac. N. Sc. Phil., 320.

Body compressed posteriorly; head depressed, muzzle from above rounded truncate; tail narrowed, obtuse. Rostral narrow. Eye invisible, ocular extending to labials. A preocular, little wider than ocular. Supralabials four. Scales in 18 longitudinal rows. Yellow, scales of seven dorsal rows with a large brown spot at the base. Top of head and end of tail immaculate. Cordova and Orizaba, Mexico. (From descr.)

TYPHLOPS (IDIOTYPHLOPS) EMUNCTUS *spec. nov.*

Very long, slender, enlarging toward the tail; head not distinct from the neck, slightly depressed, rounded, muzzle extending beyond the lower jaw; tail not longer than the thickness of the body at the vent, rounded to a point. Eye small, indistinct or invisible. Shields of the side of the head small. Rostral large, bent far back on the head, lateral margins concave, rather narrow, widening toward the top of the head, followed by two short, broad verticals, acute angled laterally, of which the posterior is shorter. Nasal deeper than long, divided diagonally from end to end; nostril opening forward. Above the nasal a small plate has the form of a spherical triangle with concave sides. Labials four; posterior scale-like; anterior largest, beneath the nasal and extending forward in a long point

between it and the rostral; third extending obliquely backward beneath the ocular, with which it is in contact at the angle. The triangular internasals are widely separated by the rostral. Below the internasal, a small rhomboid plate rests in a notch between the second and third labials, and behind this plate lies the small preocular.

Scales small, hinder margin rounded, in 22 rows, similar above and beneath. Silvery brown, snout and lips light yellow. Hab. Panama and northward.

ANOMALEPIS *Jan.*

Teeth on the upper jaw. Rostral plate narrow, hardly reaching the top of the head. Nasal grooved behind the nostril. Head with large plates on the crown. Labials two. Scales smooth, lustrous, small in the ventral rows. Anal divided.

ANOMALEPIS MEXICANUS.

Jan., 1861, *Icon. Gén. Ophid., Livr. I, pl. V.*

Body long, slender, subcylindrical; head indistinct; tail very short, bluntly rounded on the end. Muzzle broad, rounded. Mouth inferior, outline not regularly curved. Eyes visible through the ocular. Rostral narrow, scarcely reaching the top of the head. A pair of broad frontals. Frontal shield-shaped, nearly as broad as long. Supraoculars larger than the ocular. Ocular smaller than the preocular. Nasal with a groove behind the nostril. Loreal larger than the first labial. Labials two, anterior small, quadrangular; posterior larger, separated from the preocular by a narrow, elongate plate, and from the ocular by another that is similar but shorter. Lower labials five, small. The scales in contact with the head shields are larger than those farther back. Scales small, smooth, glossy, those of the belly not larger. Anal bifid.

Color reddish-brown, very light below. A yellowish-white border on the scales produces the appearance of reticulations. (From *Jan.*) Hab. Mex.

STENOSTOMA *Wagler.*

Body slender, cylindrical; tail short. Muzzle large, extending beyond the lower jaw. Mouth inferior. Teeth on the lower jaw (3-5). Eye visible through the ocular. Nostrils anterior. Nasal grooved or divided. Rostral large, reaching backward on the top of the head. Ocular large.

Labials two to three, rarely four, not in contact with the rostral. Anal entire, larger than the adjacent scales.

Tropical and semi-tropical regions.

- *Scales in 14 longitudinal rows;
 two labials, separated by the ocular;
 a small plate between the ocular and the medial row;
 lower labials four *dulce*.
 no small plate between ocular and medial row;
 lower labials five *tenuiculum*.
 three labials, two in front of the ocular;
 a small plate between ocular and medial row
myopicum.
 Scales in 13 rows *phenops*.

STENOSTOMA DULCE.

RENA DULCIS Baird & Girard, 1853, *Cat. N. A. Serp.*, 142.

STENOSTOMA DULCE Jan, 1861, *Icon. Ophid.*, *Livr. II*, pl. V.

Body long, slender, cylindrical; head indistinct, slightly depressed, muzzle rounded; tail short, thick, ending in a spine. Mouth semilunar, inferior. Eyes distinctly visible. Rostral reaching on the top of the head nearly as far back as the eyes. Nasal obliquely divided in two parts, nostril between, separated from the ocular by a single labial. Ocular large, separated from the medial series by a single plate. A large labial behind the ocular. Lower labials four. Scales smooth, lustrous, in 14 rows, ventral rather smaller, median dorsal row in contact with the rostral and separating the supraoculars.

Reddish-brown above, reddish-white beneath. Hab. Texas.

STENOSTOMA TENUICULUM, *sp. n.*

Body long, slender, cylindrical; head indistinct, slightly depressed, rounded on the muzzle; tail short, thick, ending in a point. Mouth semilunar, inferior. Eye small, distinctly visible. Rostral reaching on the head nearly as far back as the eyes. Nasal in two parts, nostril between, separated from the ocular by a single labial. Ocular large, in contact with the vertebral series (*i. e.*, oculars separated by a single scale on the top of the head). A large labial behind the ocular. Lower labials five. Scales smooth, glossy, posterior margins rounded, in 14 rows, medial row of the

* See Synopsis, p. 130.

belly larger, median dorsal row in contact with the rostral and separating the oculars.

Light greyish-brown, white beneath.

Described from San Luis Potosí, Mex.

STENOSTOMA MYOPICUM *spec. nov.*

Body long, slender, subcylindrical; head indistinct, slightly depressed, muzzle rounded; tail short, thick, abruptly rounded downward and from the sides to the terminal spine. Mouth semilunar, inferior. Eyes distinctly visible. Rostral broad, reaching back on the head to a vertical from the eye. Nasal in two parts, nostril between. Two labials between nasal and ocular, anterior lower and beneath the nasal, posterior higher, extending upward in a sharp angle in front of the eye. Ocular large, rather narrow, separated from the medial row by a small plate. A large labial behind the ocular. A pair of temporals, anterior resting on the posterior labial and in contact with the ocular. Lower labials five, anterior and posterior small. Scales smooth, glossy, rounded on the posterior margin, in 14 rows, medial row of belly slightly larger, median dorsal in contact with the rostral, separated from the oculars by a small scale. Anal entire.

Dark reddish-brown above, white tinged with purple beneath. Largest specimen 8 inches; tail, .43 in. Hab. near Tampico, Mex. Dr. Edw. Palmer.

This species bears some resemblance to *S. bicolor*, Jan. It is distinguished by the height and shape of the anterior labials, and the narrowness of the upper half of the nasals.

STENOSTOMA PHENOPS.

Cope, 1875, *Jour. Ac. N. Sc., Phil.*, 128.

"Scales in 13 rows. The eye is distinct, and the nargal suture extends to the rostral plate. The lower surfaces are without marking, but the scales of the upper surface are black, with pale borders. There are three white spots: one on the end of the rostral plate, one at the apex, and one on the under side of the tail. Total length, 0^m.156; tail, .009. The same species was obtained near Coban, Guatemala, by Henry Hague." Tehuan-tepee. Known only from the description.

ONYCHOPHIDIA.

With rudimentary limbs, the only external evidence of which is a claw on each side of the vent. Pupil vertical.

Body nearly cylindrical;
head rather indistinct from the neck;
tail usually short, thick, non-prehensile

ERYCIDAE.

Body compressed;
head distinct from the neck;
tail generally prehensile; larger teeth in front

BOAEIDAE.

ERYCIDAE.

Body nearly cylindrical; head small, rather indistinct; tail non-prehensile, short, thick, usually blunt. Eyes small, pupil vertical. Head-shields irregular. With a spur on each side of the vent.

Asia, Africa, E. Indies, America.

CHARINA *Gray*.

Body stout, subcylindrical; head small; tail short, thick, blunt. Eyes small, pupil vertical. Head-shields irregular. Nasals in contact behind the rostral. Nostrils lateral. Labials varying in number, reaching the orbit, or subdivided and separated from it by small suborbitals. Loreals one or more. Scales smooth, imbricate. Subcaudals one row, sometimes irregular. Anal entire. Spurs small. Puget Sound to Mexico.

CHARINA BOTTAE.

TORTRIX BOTTAE *Blainville*, 1835, *Nouv. Ann. du Mus. (III)*, 290, pl. 26, f. 1-1b.

CHARINA BOTTAE *Gray*, 1849, *Cat. Snakes Brit. Mus.*, 113.

Body moderately elongate, subcylindrical, slightly compressed, little fusiform; head small, rather indistinct; tail short, thick, blunt. Eye small, pupil vertical. Mouth cleft moderate, outline nearly straight. Snout prominent, broad, rounded. Head-shields and number of scales on the body varying greatly. Rostral broad. Prefrontals varying, usually two pairs. Frontal broader than long. Parietals broad and short, or dissected. Supraorbital short, broad. Nasal in two parts, nostril between; the anterior portions are longer, and meet on the top of the snout, or are subdivided to form a pair of internasals. One or more loreals. One or

more ante and two or more postorbitals, or the eye surrounded by a ring of eight or ten scales. Labials nine or more, anterior commonly higher. Infralabials ten or more, anterior broader or subdivided. Two or three pairs of small submentals, decreasing in size backward. Scales small, smooth, number of rows varying greatly (highest noticed, 45). Ventrals varying in width and number (near 200). Anal entire.

Color uniform leaden to reddish-brown, light yellowish below.

CHARINA TRIVIRGATA.

LICHANURA TRIVIRGATA Cope, 1861, *Pr. Ac. N. Sc., Phil.*, 304.

Rostral prominent, elevated, recurved, five-sided. Loreals five. Ocular ring of ten scales. Labials 14-15, anterior higher. Infralabials 15, anterior five longest. A short mental fissure. Scales in 40 rows, inferior a little larger. Total length, 25 inches; tail 4. Brownish-yellow, lighter below. Belly and flanks speckled with reddish-brown. Back with three deep, reddish-brown bands from muzzle to end of tail, median four, lateral five, and separating spaces three and a half scales in width. (From descr.)

Hab. Lower Cal.

Possibly a variety of the preceding.

BOAEIDAE.

Body more or less compressed; head distinct; tail generally prehensile. Pupil oblong, erect. Larger teeth in front. Submental fold conspicuous. A spur or rudimentary limb on each side of the vent.

BOA Linné.

Body elongate fusiform, slightly compressed; head distinct, flattened on the crown, broad posteriorly, muzzle prominent; tail prehensile. Nasals entire or divided. Nostril lateral. Teeth smooth, larger forward. Labials not pitted, short, numerous. Scales covering the top of the head and the loreal region and surrounding the eye, small. Dorsal scales small, smooth, flat. Anal entire. Subcaudals simple.

BOA IMPERATOR.

Daudin, 1803, *Hist. Rept.* I, 150.

Body robust. Muzzle truncate, outline of extremity nearly vertical. Scales of head, face, and temples small, rather larger than those of *B. constrictor*. Eye surrounded by 16 to 18 scales, of which several are in

contact with the labials. Labials about 20, infralabials 20 to 22. Scales in 65 to 69 rows, outer broader. Ventrals 233 to 240; subcaudals 52 to 57. Brownish, varying from light to dark (in alcohol), with a dorsal series of quadrangular brown spots, emarginate anteriorly and posteriorly, separated from the series of triangular brown spots on the upper portion of the flank by an interrupted narrow line of light color, which is most distinct opposite the spots. Upper spots of the flank larger and more distinct. An irregularly placed series on the outer rows of scales and the edge of the abdomen. Spots darkening toward the tail, often with lighter centers. A brown line on the head from the frontal region to the neck. A brown line from behind the angle of the mouth through the eye to the loreal region, spreading toward its extremities. A bar from the eye to the lip. A pair of spots on each side of the lower jaw. Rostral with a spot. Scales irregularly punctulate or spotted. Mex.

I have little doubt that this will eventually be degraded to the rank of a variety. Specimens at hand from the Isthmus of Darien are intermediates between this species and *B. constrictor*.

BOA CONSTRICTOR.

Var. isthmica.

Body stout, compressed; head distinct; snout broad, truncate; tail short, prehensile. Rostral large, broader near the upper edge, concave at the sides and below, rounded above, surrounded by scales larger than the others of the head. Nasal divided. Eye surrounded by eighteen or nineteen scales. Several larger scales in front of the oculars. Two rows between oculars and labials. Labials 22-23. Infralabials 25-27. Scales small, flat, smooth, in 78 rows, outer broader. Ventrals narrow, 246. Subcaudals 57.

Ground color light greyish (in alcohol). Pattern of markings as in *B. constrictor*. Transverse brown bands of the back twenty-five, posteriorly united by the darker margins with the triangular spot on the upper part of the flank. Beneath the latter on the flank there is a series of oblong, rounded spots, each enclosing a rounded spot of lighter color. A series of more or less broken quadrangular black spots on each side of the belly includes scales of the outer row. The spots unite on the tail so as to form a dorsal and subcaudal row. A light brown band from rostral to neck includes several light spots on the occipital region. A brown band with

white margin extends from a point close behind the angle of the mouth through the eye to the nasal, broadening on the loreal region, and descending across the mouth to the submentals. A short distance in front of the angle of the mouth on the lower jaw there is a rounded brown spot surrounded by a white ring, by which it is separated from the edge of the lip. An angular black spot descending from the eye does not reach the labials. Central America.

BOA MEXICANA *Rapp.*

BOA DIVINILOQUAX var. *Mexicana* Jan, 1864, *Icon. Ophiüd., Livr. V, pl. II.*

The figure by which this species is known is that of a serpent more closely allied to *B. imperator* than to *B. diviniloquax*. The muzzle is broader and more vertically truncate than in the former, and the ground color is more bleached. The pattern of coloration is the same, but some of the lines of the head and the spots and punctulations of the flanks have become obsolete. Nasals divided. Suborbitals in contact with labials. Scales in 55 longitudinal rows.

ACACOPHIDIA.

Head and tail more distinct from the body, and eyes and teeth more perfectly developed than in the *Scolecophidia*; without claws at the side of the vent, as in the *Onychophidia*; and without poison-secreting glands and fangs, as in the *Toxicophidia*.

This group includes all the common, non-venomous snakes of the ground, trees, or fresh waters. It contains but a single family. For convenience this is divided into subfamilies, nearly corresponding to what have heretofore been accepted as families, and for which the old names have been retained.

COLUBRIDAE.

- Elongate, tapering, compressed; head broad; pupil generally elliptical
DIPSADINAE.
- Moderately slender, subcylindrical; head moderate; pupil elliptical
SCYTALINAE.
- Slender, long; head long, narrow; eyes large; pupil round
DENDROPHINAE.
- Fusiform, slender to stout; head broad behind; posterior maxillary teeth usually larger; scales generally keeled
NATRICINAE.
- Elongate, tapering to head and tail; head distinct from the neck, moderately broad; crown-shields regular; loreal usually present; teeth smooth
COLUBRINAE.
- Subcylindrical; head more or less distinct, short, shields commonly regular; scales rarely keeled
CORONELLINAE.
- Cylindrical; head short, rather indistinct; usually some head-shields united; scales smooth or keeled
CALAMARINAE.

DIPSADINAE.

Tree snakes. Bodies more or less elongate and compressed, tapering to head and tail; head large, distinct, somewhat triangular, broad behind, depressed, muzzle broad and rounded; tail short to long and slender. Eye large to very large, pupil generally elliptical. Nostrils lateral. Teeth varying much. In the majority of the species the posterior maxillary teeth are longer and grooved. Head-shields nine, short, broad. Scales imbricate, smooth, rarely keeled, generally the vertebral row larger.

Teeth equal, smooth;
anal entire; scales in 13-17 rows,
vertebral series larger

LEPTOGNATHUS.

Posterior maxillary teeth longer, grooved;
anal entire; scales in 19-27 rows,
vertebral series larger

DIPSAS.

anal divided, rarely entire; scales in 19-23 rows,
dorsal series similar

SIBON.

LEPTOGNATHUS.

Duméril, 1852.

Body long, more or less compressed; head subquadrangular, high, distinct from neck; tail medium. Eye moderate, pupil erect. Teeth equal, smooth. Head-shields nine, short, broad. Rostral broad, low. With or without a distinct loreal. Orbitals 1-2+2-3, suborbitals sometimes present. Scales moderate, rarely keeled, in 13-17 rows, vertebral generally larger. Anal rarely divided. Subcaudals in two rows.

Hab. Mexico to the Argentine Republic; India.

Scales keeled in vertebral series, rows 15-17;	
loreal distinct	<i>fasciatus.</i>
Scales smooth, rows 17;	
loreal distinct	<i>dumérilii.</i>
Scales smooth, rows 15;	
loreal and anteorbital fused;	
transverse bands narrow	<i>nebulatus.</i>
transverse bands broad	<i>dimidiatus.</i>

LEPTOGNATHUS FASCIATUS.

TROPIDODIPSAS FASCIATA, *Gthr.*, 1858, *Cat. Col. Snakes*, 181.

Slender; head oval, crown slightly flattened, muzzle broad; tail long, slender. Eye large, pupil erect. Head-shields broad. Frontals bent on the side. A loreal. Orbitals 2-2. Temporals 1+2. Labials 6-7, penultimate largest. Infralabials 7, fifth largest. Submentals two or three

pairs, second half as large as first. Scales in 15–17 rows, some of the dorsals with keels. Anal entire. Subcaudals in two rows.

Dark brown or black. The light color of the throat and chin-shields extends upward as an occipital collar. Flanks with widely separated, wedge-shaped, vertical bars of light color, which frequently meet in their points on the back, forming transverse bands to the sides of the belly. Belly much spotted with dark posteriorly.

Hab. Mexico and Central America.

LEPTOGNATHUS DUMERILII.

Jan, 1870, *Icon. L'ivr. XXXVII*, pl. V, fig. 2.

Scales smooth, in 17 rows. The principal difference between this species and the preceding is in the carination of the scales. It is altogether likely that individual variations in this respect will do away with its value, in which case the form cannot even stand as a variety.

LEPTOGNATHUS NEBULATUS.

PETALOGNATHUS NEBULATUS *Duméril*, 1852 (1853), *Mém. de l'Acad. Sci. XXIII*, 466.

Slender, elongate, much compressed; head large, distinct, crown slightly convex, muzzle high; tail slender, tapering. Eye large, pupil erect. Head-shields short, broad. Rostral as broad as high. Nasal in two parts, nostril principally in anterior. Loreal and preocular fused. One preocular, sometimes a small plate at its lower posterior angle. Postoculars two. Temporals 1+2+3. Labials seven, fourth and fifth below the eye, sixth largest. Infralabials nine, sixth largest. Submentals three pairs (2–3), anterior larger. Scales smooth, in 15 rows, several of the dorsal broader, the median broader than long. Ventrals 185–197. Anal entire. Subcaudals 87–90 pairs.

Greyish-brown, mottled with brown, with more or less irregular narrow transverse white-edged bands of brown. A series of large blotches on each side of the abdomen, apparently separated at the median line, more or less confluent with the dorsal bands, and frequently with them forming complete rings around the body. Belly yellowish, clouded and spotted with brown. Head brown, irregularly marked with white (yellow). First brown band immediately behind the head. Tail darker. Total length 17 inches; tail 4. Total length $14\frac{1}{2}$ in.; tail $3\frac{3}{4}$. Hab. Mexico to Brazil; West Indies.

LEPTOGNATHUS DIMIDIATUS.

Günther, 1872, *Ann. & Mag. Nat. Hist.*, IX, 31.

Long, slender, much compressed; head broad, short; tail less than one third of the total length. Eye rather large, pupil erect. Internasals small. Prefrontals large, bent down on the face, entering the orbit. Outer edges of frontal nearly parallel. Loreal and preocular fused. Postoculars two. Temporals 1 + 2. Submentals three pairs, anterior largest, crescent-shaped; middle longer than broad, posterior divergent. Labials eight, fifth and sixth in orbit. An azygos scale-like shield under the chin, surrounded by the first pair of infralabials, the mental and submentals. Second to sixth infralabials in contact with anterior submentals. Scales smooth, in 15 series, vertebral not larger. Ventrals 186. Anal entire. Subcaudals 98 pairs.

Body and tail with broad black rings, separated by narrow, whitish interspaces. Twenty-five rings on the trunk, sixteen on tail; white interspaces subdivided by a narrow black transverse line. Upper parts of head black; with small, irregularly placed whitish spots, and a pair of large spots of the same color on the neck, forming a sort of collar. Anterior chin-shields black. (From Günther.) Hab. Mexico.

DIPSAS.

Laurent, 1768.

Slender, elongate, compressed; head very large, distinct, depressed, broad behind, angles rounded, muzzle blunt; tail slender, tapering. Eye moderately to very large, pupil erect. Posterior maxillary teeth longer, grooved. Rostral broad, low. Head-shields nine, broad. Nasal divided. A loreal, rarely fused with orbital. One preocular. Postoculars two to three. Scales smooth, rarely keeled, in 19—27 rows, vertebral larger. Anal entire or divided. Subcaudals in two rows. Hab. Mexico to Brazil; Australia; Africa.

(*Himantodes*.)

DIPSAS CENCHOA.

COLUBER CENCHOA Linné, 1758, *Syst. Nat. Ed. X*, 226.

DIPSAS CENCHOA Fitzinger, 1826, *New Class Rept.*, 59.

Very long and slender, compressed, neck and tail very small; head large, very broad, muzzle short, crown convex; tail near one third of the total. Eye large, pupil erect. Mouth deeply cleft, curved. Internasals

about one third as large as the prefrontals. Rostral subtriangular, broader than high. Nasal divided, nostril mainly in anterior portion. Loreal small. Preocular narrow, high. Postoculars two (sometimes one), small, lower resting on the fifth labial. Labials eight, fourth and fifth in orbit, sixth and seventh larger. Infralabials nine, sixth larger. Submentals two pairs, anterior larger, posterior separated by small scales. Scales lozenge-shaped, smooth, in 17 rows, medial and outer broader. Ventrals 244 (228—259). Anal bifid. Subcaudals 134 pairs (134—174).

Light greyish-brown, scales more or less punctulate with brown. Anteriorly with dark-edged transverse blotches of brown; posteriorly these become smaller, do not descend on the flank, and are very irregular and broken. Hinder portion of body with spots irregularly distributed on the lower portions of the flanks. Specimen described from Acapulco, Mex. Total length 33 inches; tail 10 in. Hab. Mexico to Brazil.

DIPSAS LEUCOMELAS.

HIMANTODUS LEUCOMELAS Cope, 1861, *Proc. Ac. N. Sc. Phil.*, 296.

Lateral borders of frontal nearly parallel, longer than the anterior. Loreal higher than long. Two preoculars, not in contact with the frontal. Infralabials ten, sixth largest. Scales smooth, in 17 rows, vertebral larger, broader than long. Total length about three and one half times that of the tail.

Greyish, crossed by about twenty-nine black bands, which encroach on the ends of the ventrals. About eighteen spots on the tail. Spotted or dotted with black on the belly. A pair of black spots from the supraciliaries over the occipitals to the neck. A band of black across the prefrontals. Scale pores single. (Unknown to me.) Vera Cruz.

SIBON.

Fitzinger, 1826.

Moderately long, slightly compressed, tapering to head and tip of tail; head distinct, large, broad, depressed, with broad muzzle; tail moderate. Eye medium, pupil oblong, erect. Posterior maxillary teeth longer, grooved. Head-shields regular. Nasal in two parts, nostril between. One or more loreals. Oculars 1 to 3—2 to 3, sometimes with suboculars. Scales smooth, imbricate, in 19—23 rows, dorsal not larger. Anal and subcaudals divided. Hab. Texas and Mexico to Brazil.

Long, slender;

Scales in 23 (21—23) rows, ventrals 266 (\pm)

biscutatatum.

Scales in 21 rows, ventrals 206—210 (\pm)

var. *latifasciatum.*

Moderate;

Postmaxillary tooth longer, grooved;

Scales in 19—25 rows, ventrals 181—188 (\pm)

annulatum.

dorsal series broad transverse bands

var. *septentrionale.*

dorsal series of spots rather small

torquatatum.

Postmaxillary teeth in continuous series, not grooved;

Scales in 19 rows; ventrals 179—182 (\pm)

discolor.

SIBON BISCUTATUM.

SIBON BISCUTATA *Dun. Bibr., 1854, Eqp. Gén. VII, 1153.*

Long, slender, compressed; head distinct, broad oval, depressed, snout prominent, broad, rounded; tail moderate, tapering. Eye large, pupil erect. Mouth-cleft deep, slightly curved. Anterior maxillary teeth longer; posterior large, isolated, grooved. Head-shields nine, broad. Rostral broader than high. Nasal divided. Loreals two to four. Orbitals 2 to 3—2 to 3. Temporals irregular. Labials 9—10, fourth and fifth usually under the eye, the four preceding the last larger. Infralabials 11—13. Submentals two pairs; posterior small, not in contact. Scales in 23 rows. Ventrals 264 (\pm). Anal bifid. Subcaudals 94 (\pm) pairs.

Light brown, punctulate, with broad transverse bands or rhombs of brown, which are paler in the center and separated by spaces about equal to their width. The bands become narrower toward the abdomen, at the edge of which there is an alternating series of spots. Abdomen sprinkled with brown; subcaudals with blotches. A light band across the head behind the internasals; another across the frontal passes through the eyes to the angle of the mouth. A V-shaped light spot on the vertex and a light collar behind the occiput. (From D. & B. and Jan.) Hab. Mexico and southward.

VAR. LATIFASCIATUM.

DIPSAS BISCUTATA VAR. LATIFASCIA *Peters*, 1869, *MB. Berl. Acad.*, 877.

Rows of scales 21—22. Ventrals 206—210. 13—14 brown bands on the body; 5—6 on the tail. (From *Peters*.) Puebla, Mexico.

SIBON ANNULATUM.

COLUBER ANNULATUS *Linné*, 1754, *Mus. Ad. Fridr. Tab. 8, fig. 2*.SIBON ANNULATUS *Fitzinger*, 1826, *Neue Class. Rept.*, 60.

Moderate, tapering to head and tip of tail; head large, distinct, depressed, crown flattened, temples swollen; muzzle broad; tail medium. Eye moderate, pupil oblong, erect. Posterior maxillary teeth longer, grooved. Nasal in two parts; nostril lateral. Loreal present. Preoculars two; upper large, reaching the top of the head, lower small in a notch between the third and fourth labials, sometimes absent. Postoculars two, upper large. Temporals 1+2. Labials eight, fourth and fifth in orbit, seventh largest. Infralabials 10 (9—10), sixth largest. Submentals two pairs, nearly equal. Scales smooth, in 23 rows (19—23), similar. Ventrals 188±. Anal bifid.

Light reddish-brown, with a series of brown spots on each side of the median line, which are alternated and form a zigzag line or opposed and form transverse bands or rhombs. The number of spots varies much; in the specimen before me there are fifty on the body alone. Flank mottled with brown, lower portion with a series of spots alternating with those of the dorsal. Head of the ground color, with a brown band behind the eye, above the angle of the mouth. Belly uniform light color. Scales lustrous. Hab. Mexico to Brazil.

VAR. SEPTENTRIONALE.

DIPSAS SEPTENTRIONALIS *Kennicott*, 1859, *U. S. and Mex. Bound. Surv., II, pl. VIII, f. 1, p. 16*.

Preoculars three (2—3); upper large, reaching the top of the head; lower very small. Scales in 19—23 rows, smooth.

Yellowish white (brownish), with irregular transverse bands of brown, which extend across the back to the ventrals. The bands vary much in shape and size; the specimen at hand has thirty-eight in all. Old specimens with darker colors. A band from the nostril through the eye to the hinder labial. Head mottled or blotched with brown, sometimes with a shield-shaped spot on the occipitals. Hab. Texas to Central America.

SIBON TORQUATUM.

LEPTODEIRA TORQUATA *Günther*, 1860, *Ann. & Mag. Nat. Hist.*, pl. X, 169.

Posterior maxillary tooth larger, separated by an interspace. Scales in 21 rows. Ventrals 174. Anal bifid. Subcaudals 44—56. Total length 16 in. 5 l.; tail 3 inches. Total length 17 in. 5 l.; tail 2 in. 3 l.

Brownish-grey, with a vertebral series of brown spots, some of which are confluent into a zigzag band; neck with a white collar, behind which there is a large brownish-black blotch. Hab. Nicaragua; Island of Laguna. (From Günther.)

SIBON DISCOLOR.

LEPTODEIRA DISCOLOR *Günther*, 1860, *Pr. Zool. Soc. Lond.*, 317.

Posterior maxillary tooth larger, in a continuous series with the other teeth, not grooved. Scales in 19 rows, smooth, rhombic, those of the sides similar to those of the back. Ventrals 179—182. Anal bifid. Subcaudals 87—88. Total length 21 in. 1 l.; tail 6 in.

Dirty white, with numerous black cross-bands extending on to the ventral plates; belly uniform whitish. Fifty-one to fifty-four black bands cross the trunk and extend to the edge of the belly; they are broader than the interspaces between, and become interrupted and spot-like on the tail. Upper part of head brown, with a whitish collar behind the occipitals. Separated from *S. annulatum* by the teeth. Hab. Oaxaca, Mex. (From Günther.)

SCYTALINAE.

Moderate, cylindrical to subcylindrical; head distinct from the neck, broad behind, crown flat; tail medium, tapering. Eye medium, pupil oblong or elliptical. Crown-shields nine. Nasal generally divided. A loreal. Preoculars one to two. Postoculars two. Posterior maxillary teeth longer, grooved. Scales smooth, in 17 to 19 rows, occasionally the vertebral series larger. Anal entire. Subcaudals simple or bifid.

OXYRHOPUS.

Wagler, 1830.

Elongate and subcylindrical to slender and slightly compressed; head moderate, a little larger than the neck, depressed, muzzle rather broad, crown flat; tail short to moderate. Eye medium, pupil elliptical, erect.

Posterior upper maxillary teeth longer, grooved. Anterior teeth equal. Crown-shields regular. Nasal divided. A loreal. Preocular one, sometimes two. Two postoculars. Scales smooth, lustrous, in 19 rows, varying rarely to 17 or 15. Anal entire. Subcaudals in two rows.

Light reddish-white; scales more or less tipped with brown;
a yellowish collar *cloelia*.

Reddish-white, with about thirty black rings around the body.
doliatus.

OXYRHOPUS CLOELIA.

COLUBER CLOELIA *Daudin*, 1799, *Hist. Rept. VI*, pl. 78, p. 330.

OXYRHOPUS CLOELIA *Günther*, 1858, *Cat. Col. Serp.*, 189.

Elongate, slightly compressed, body tapering toward the tail; head larger than the neck, depressed, rounded, crown flattened, muzzle broad; tail rather less than a fourth of the total, tapering. Eye moderate, pupil erect. Mouth-cleft but slightly curved. Crown-shields nine, broad. Rostral as high or higher than broad, bent back in a blunt angle between the internasals. Prefrontals broad, bent down on the side of the face. Frontal broad, angle between the parietals a little less than right. Parietals large. Nasal in two parts, anterior acute-angled between the rostral and first labial. Loreal small, narrow in front. Oculars 1—2. One temporal in contact with postoculars, a second below it in a notch between the labials. Labials seven to eight, third and fourth or fourth and fifth in orbit, the two preceding the last larger. Infralabials eight to nine, fifth larger. Submentals two pairs, hinder smaller. Scales smooth, lustrous, with two pores, in 19 rows. Ventrals 242. Anal entire. Subcaudals 77 pairs.

Light reddish-white, whiter below. Tips of scales slightly dotted with brown. Crown-shields and sides of face dark brown. A triangular brown patch on the back and sides of the neck, in front of which there is a yellow collar which extends forward under the eye on the labials. Throat and chin uniform white. Behind the dark patch on the neck the scales become gradually lighter. Central America.

OXYRHOPUS DOLIATUS.

Duméril & Bibron, 1854, *Erp. Gén. VII*, 1020.

Elongate, compressed; head distinct from the neck, crown flat, broad behind; tail moderate, less than one fourth of the total, tapering. Snout short, broad, rounded. Eye moderate, pupil elliptical, erect. Posterior

maxillary tooth longer, grooved. Crown-shields regular. Rostral broad, low, reaching the top of the snout. Internasals much smaller than prefrontals. One preocular, not in contact with the frontal. Postoculars three (2—3). Loreal elongate. Labials eight, fourth and fifth in orbit. Two temporals in contact with the oculars. Scales smooth, in 19 rows, medial rather larger. Anal entire.

Yellowish in alcohol, reddish in life, with thirty or more black rings—more or less complete—encircling the body. The intervals on the back are of varying width, reddish-white, and spotted with black. A dark spot on the crown, somewhat broken by lighter streaks on the edges of the plates. Belly yellowish, black bands often broken or incomplete. Central and South America.

DENDROPHINAE.

Tree snakes. Slender, elongate; head very distinct from the neck, narrow, long, depressed, snout prominent; tail not distinct from the body, very slender. Eye moderate to large, pupil round. Nostrils small, lateral. Posterior upper maxillary teeth longer, grooved or smooth. Scales long, narrow, in 15 to 21 rows. Ventrals commonly with a slight keel at the side of the abdomen. Subcaudals in two rows. Tropics of both continents.

Posterior upper maxillary teeth longer, smooth; scales smooth or keeled	LEPTOPHIS.
Posterior maxillary teeth longer, grooved behind; scales smooth	OXYBELIS.

LEPTOPHIS.

Bull. 1825.

Very slender, long; head long, narrow, depressed, distinct from the slender neck. Snout produced beyond the lower jaw. Eye large, pupil round. Postoculars 2—3. Abdomen slightly keeled at the sides. Postmaxillary teeth longer, smooth. Scales in 15 to 19 rows, smooth or keeled. Subcaudals in two rows.

LEPTOPHIS MEXICANUS.

LEPTOPHIS MEXICANUS *Donvil & Bibron*, 1854, *Exp. Gén.* VII, 536.

Two postoculars. Loreal present. Labials eight. Scales in 15 rows, carinate on trunk and tail. Ventrals 157—169. Anal bifid. Subcaudals 138—154 pairs. Mexico. (From descr.)

OXYBELIS.

Wagler, 1830.

Body much elongate, slender; tail long. Head long, slender; snout very long, produced. Rostral inferior. Pupil round. No loreals. Nasal entire. Oculars 1—2. Scales smooth. Subcaudals bifid. Posterior maxillary teeth longer, grooved.

OXYBELIS AENEUS.

DRYINUS AENEUS Wagler, 1824, *Spir. Serp., Brazil*, 12, pl. III.OXYBELIS AENEUS Wagler, 1830, *Syst. Amph.*, 183.

Very slender, slightly compressed; head distinct from the neck, long, narrow, crown convex; snout elongate and narrow, sides concave; tail very long and slender, a little shorter than the body. Eye large, pupil round. Mouth deeply cleft, outline sinuous. Crown-shields nine, anterior elongate, parietals broad. Rostral reaching the top of the snout, but not bent backward. Internasals long. Prefrontals long, bent down on the side. Frontal narrow, tapering backward. Supraciliaries long, broad posteriorly. Nasal long, narrow, entire. No loreal. One anteocular, as broad as long. Postoculars two, lower small. Labials eight, posterior very long, fourth to the sixth in the orbit. Infralabials eight, fifth and sixth larger. Temporals large 1+2. Submentals two pairs, posterior larger. Scales long, pointed, in 17 rows, outer and vertebral broader. Ventrals 183. Anal bifid. Subcaudals 197 pairs.

Light iridescent greyish or silvery punctulate with brown, sides and belly slightly darker. A narrow line of light color along the middle of the belly, more or less distinct. Irregularly sprinkled with brown spots on the back and flanks. Anteriorly the belly, throat, chin, and labials are nearly white, above this a dark brown border extends from the nostrils to the sides of the neck. Top of head bronzed and clouded. Specimens described from Obispo, Cent. Amer. Total length $41\frac{1}{2}$ inches; tail $18\frac{3}{4}$. Total length $23\frac{1}{2}$ inches; tail $11\frac{1}{4}$.

NATRICINAE.

Small to large, slender to stout, generally fusiform, approaching the subcylindrical in *Hydrops*; head distinct, subtriangular, muzzle narrow across the nasals; tail slender to stout, commonly tapering. Eye medium to large, pupil round. Maxillary teeth subequal in a continuous series, or posterior isolated, equal or larger, smooth or grooved. Internasals small,

subtriangular or fused, sometimes displaced by the nasals. Scales keeled, except in *Hydrops*.

Scales keeled;	
nasal divided;	
teeth smooth in continuous series;	
loreal present	TROPIDONATUS.
loreal absent or present	STORERIA.
nasal grooved or divided;	
postmaxillary teeth larger and isolated	HELICOPS.
scales smooth;	
nasal grooved or divided;	
postmaxillary teeth little larger	HYDROPS.

TROPIDONOTUS.

Kuhl, 1826.

Stout to slender, tapering to head and tail, belly round; head distinct, crown flat, occipital tract broad, snout narrow; tail tapering to a point. Eye moderately large, pupil round. Teeth smooth and equal, or posterior a little larger. Head-shields normal, anterior small. Nasal in two parts, nostril between. Internasals small, narrowed anteriorly or triangular. A loreal. Preoculars 1—3. Postoculars 2—5. Suboculars present in *T. cyclopium*. Scales keeled, pointed, truncate or emarginate, in 17—33 rows, outer broader. Subcaudals in two rows.

Anal entire;	
series of rufous spots; rostral resembling that of <i>Salvadora</i> ;	
of scales 21	<i>rufipunctatus</i> .
striped; series of scales 19 (17—19); slender	<i>saurita</i> .
striped and spotted; series of scales 17—21	<i>sirtalis</i>
Anal bifid;	
banded transversely; series of scales 23—25	<i>fasciatus</i> .
transverse bands and spots; series 29—31	<i>tarvispilatus</i> .
transverse bands to uniform; with suborbitals; series 27—33	<i>cyclopium</i> .
irregularly spotted; series 19—21	<i>compressicaudus</i> .
spots in longitudinal rows; series 19	<i>kirtlandii</i> .
striped; orbitals 2—2; series 19	<i>leberis</i> .

TROPIDONOTUS RUFIPUNCTATUS.

CHILOPOMA RUFIPUNCTATUM Yarrow (*Cope MS.*), 1875, *Zool. Wheeler's Survey*, p. 544, pl. XX, fig. I.

Moderately slender, tapering to head and tail; head distinct, elongate, narrowed anteriorly; tail about one fourth of total. Eye moderate, pupil round. Crown-shields regular. Rostral turned over the superior face of the muzzle, posterior border truncate. Internasals longer than wide. Prefrontals decurved laterally. Nasal long, narrow. Preocular higher than wide, in contact with the frontal. Two preoculars on one side. Frontal elongate, obtuse behind. Parietals elongate, bounding the upper postocular behind. Postoculars three. Temporals 1+3. Labials eight, last very small, fourth and fifth bounding the orbit below. Infralabials nine. Submentals two pairs. Scales keeled, in 21 rows, first and second smooth, poreless. Ventrals 177. Subcaudals 4+83 pairs.

Light brown above, olive shaded on the head. Anterior half of body marked with six rows of small alternating bright rufous or orange spots, each of which occupies one and sometimes an adjoining scale. They stand on the first and second, the fifth, and on the eighth rows respectively. On the posterior third of the length they are wanting, and are indistinct posterior to the middle. Lower surfaces pale brownish-grey; the base of each ventral with blackish markings. Labials light; head without spots. The figure does not agree with the description. Southern Arizona. (From descr.)

TROPIDONOTUS SAURITA, *pl. III, fig. 2.*

COLUBER SAURITA Linné, 1766, *Syst. Nat.*, ed. XII, 385.

TROPIDONOTUS SAURITA Schlegel, 1837, *Ess. Phys. Serp.* II, 321.

Body slender; head distinct, crown flat; tail long, near a third of the total length, slender. Eye large, over the fourth and fifth labials. Nine head-shields. Frontal hexagonal. Rostral broader than high. Nasal in two parts, nostril between, near the upper edge. A loreal. Oculars 1—3. Temporals 1+2. Mouth-cleft deep, curved. Labials 8 (7—8), third and fourth in orbit when there are but seven. Infralabials 10 (9—11), fifth and sixth large. Submentals two pairs, elongate, narrow, hinder larger. Scales keeled, in 19 rows, outer broader. Ventrals broad, 150—160 (with var. proximus 150—180). Anal entire. Subcaudals 110—120 (100—120).

Brown to black, olivaceous beneath. With a dorsal and two lateral narrow lines of yellow. Lateral lines upon the third and fourth rows of scales; below them on the outer rows the color is brown, sometimes very

much faded, and shades into the olive beneath. Skin between the scales black. Short white lines to be seen on stretching the skin. Specimens from the Mississippi Valley show black spots in the brown of the flank. A vertical spot of yellow on the anteorbital. Dark color of the head extending to the rostral and upper labials. A pair of small yellow spots near the inner margins of the occipitals. Closely allied to *T. sirtalis*. Valley of the Mississippi and eastward.

VAR. PROXIMUS.

Stouter. Ventrals 168—178. Subcaudals 100—108. Mississippi Valley to Mexico.

TROPIDONOTUS SIRTALIS, *pl. III, fig. 3.*

COLUBER SIRTALIS *Linnaë*, 1758, *Syst. Nat.*, ed. X, 222; ed. XII, 1766, p. 383.

TROPIDONOTUS SIRTALIS *Holbrook*, 1842, *N. A. Herp. IV*, 41, *pl. XI.*

Young slender, old moderately stout; head distinct, broad behind, narrow in front, flat on the crown; tail medium, slender posteriorly, one fourth to one fifth of the total length. Nine head-shields. Vertical hexangular. Rostral broader than high. Nasal in two parts, nostril between—in anterior portion. Loreal quadrangular. One anteorbital. Postorbitals 3—4. Eye large, over the third and fourth or fourth and fifth labials. Mouth-cleft deep, curved. Labials 7 or 8, antepenultimate largest. Infralabials 10. Submentals two pairs, hinder larger. Scales keeled in 17—21 rows, dorsal narrow, outer broad. Ventrals broad 140—155 (140—180 with the varieties). Subcaudals 50—90 pairs.

Brown above, varying from light ashy to very dark; olive beneath. A dorsal and two lateral narrow lines of yellow, sometimes faded or obliterated. In the brown space between the yellow lines on the flank are two series of quadrate alternating spots; sometimes the upper of these forms a band (*var. collaris*), and in cases the spots of the lower are joined to it (*var. pickeringii*). A brown band on the outer row (sometimes obsolete), with or without a series of spots. Black spots on the base near the extremities of each ventral are visible on bending the body. When the skin is stretched numerous short white lines are to be seen on the flanks of most specimens. A small yellow spot near the inner margin of each parietal shield. Chin and throat lighter colored, yellowish. Labials more or less bordered with black. Kentucky specimens show a light collar extending toward the occiput around the angle of the jaw, behind which

the dark color is produced toward the throat. On account of the amount of variation of this species in different localities, it is hardly to be separated from the preceding. Canada to Central America.

sirtalis. Rows 19. Spots 70—90. Ventrals 140—165+60—85 pairs.

Mississippi Valley and eastward.

marcianus. Rows 21. Spots 50—75. Ventrals 145—175+50—90 pairs.

Texas to Mexico.

parietalis. Rows 21. Spots 70—110. Ventrals 160—185+50—90 pairs.

Valley of the Missouri and westward.

infernalis. Rows 17—19. Spots 100—120. Ventrals 145—175+75—95 pairs. California and northward.

collaris. Rows 19. Ventrals 151—165+58—84 (Jan.) Mexico to Panama.

ordinatus. Agrees with *sirtalis*, but has the lower series of spots more distinct, while the stripes are almost obsolete, and is found in various isolated localities of the same district, as Alabama, South Carolina, Martha's Vineyard, and Nova Scotia.

TROPIDONOTUS SIPEDON, *pl. II, fig. 3.*

COLUBER SIPEDON *Linnaeus*, 1758, *Syst. Nat.*, ed. X, 219.

TROPIDONOTUS SIPEDON *Hallbrook*, 1842, *N. A. Herp. IV*, p. 6, *pl. 29.*

Stout, fusiform, belly broad; head distinct, narrow forward; tail smaller than the body, tapering. Eye medium, pupil round. Mouth-cleft deep, abruptly curved near the angle. Nine head-shields. Rostral low, broader than high. Nasal in two parts, nostril between. A loreal. One anteorbital, sometimes two. Postorbitals two to four. Temporals 1+2, large. Labials 8 (7—9), sixth and seventh large. Infralabials 10 (9—11), fifth and sixth large. Submentals two pairs, large, not in contact with the mental. Scales strongly keeled, in 23 to 25 rows, dorsal narrow, outer broader than long, keeled to smooth. Ventrals 130—155. Anal divided. Subcaudals 40—75 pairs.

Varying from ashy to very dark brown, or to brownish-red; with three series of dark-edged brown spots; dorsal about ten scales in width, varying in shape from irregular rounded to nearly square, distinct or confluent with the lateral into transverse bands, which widen toward the vertebral rows. There are usually twenty-five to thirty spots on the flank, wider than long, and from fifteen to twenty on the tail. Commonly the spots on the flanks are more or less opposed and confluent anteriorly, but posteriorly alter-

nated with those of the dorsal series. A faded band of dark color from the eye to the angle of the mouth. The markings are subject to great variation, often very indistinct or obsolete. Belly reddish, yellowish, or white, more or less spotted or mottled with brown, lighter under neck and chin. Canada to Mexico.

sipalon. Greyish or brownish. Rows of scales 23 (23—25). Maine to Sonora.

erythrogaster. Reddish. Rows of scales 23—25. Southeastern U. S.

rhombifer. Brownish, with transverse dorsal rhombs. Rows of scales 25.

TROPIDONOTUS TAXISPILOTUS.

Holbrook, 1842, N. A. Herp. IV, p. 35, pl. VIII.

Stout, tapering to head and tail; head moderate, broader than the neck, snout, narrow; tail about one fourth of the total length, somewhat distinct, slightly compressed, tapering. Head-shields normal. Rostral broad, low. Nasal in two parts, nostril between. A loreal. One anteorbital (1—2.) Postorbitals 2 (2—3.) A single temporal in contact with the orbitals. Labials 8 (7—9), the two preceding the last large. Infralabials 10 (9—11), fifth and sixth larger. Scales keeled, notched at the extremity, in 29—31 rows, outer rather large. Ventrals broad, 130—141. Anal bifid. Subcaudals 80 pairs. Brown, with three series of dark blotches, separated by spaces of nearly equal width. Lateral series little smaller than the dorsal. Belly brownish, marked with yellow on the fore edges of the scales, more yellow anteriorly. Southeastern United States.

TROPIDONOTUS CYCLOPIUM, *pl. II, fig. 4.*

Dunéril & Bibeon, 1854, Erp. Gén. VII, 576.

Moderately stout, elongate fusiform; belly broad, back slightly compressed; head distinct, broad behind, short in front of eyes; tail about one fourth of the total, subtriangular, tapering. Head-shields nine. Rostral broad, low. Nasal in two parts, nostril between. Lower side of loreal longer, upper short. One ante, two post, and two to three suborbitals. Temporals large, a single one in contact with the orbitals. Labials 8—9, three preceding the last large, occasionally one or more in contact with the orbitals. Infralabials 10—12. Submentals two pairs, anterior larger. Scales notched at the end, keeled, in 27—33 rows, outer broader, faintly keeled. Ventrals 138—150. Anal bifid. Subcaudals 60—75 pairs.

Clouded olivaceous brown, with narrow, irregular, more or less indistinct vertical bands of about two scales in width, separated by narrow spaces. In places on the body these bands appear to meet on the back, in others they alternate, and posteriorly most of them are divided on the middle of the flank to form alternating series of quadrate blotches, similar to those of *sipedon*. Belly brown toward the vent, with semicircular yellow spots on the hinder edges of the scutes; anteriorly it is yellow, with spots of brown, somewhat similar in shape on the bases of the plates. There is great variation in intensity and amount of marking. A very large specimen from Florida shows only a narrow band of brown on the base of each ventral. Specimens of this species from Florida are hardly to be separated from *taxispilotus*. Ohio to Florida.

TROPIDONOTUS COMPRESSICAUDUS.

NERODIA COMPRESSICAUDA *Kennicott*, 1860, *Pr. Ac. N. Sc., Phil.*, 335.

Moderate, belly broad, back slightly compressed; head distinct, narrow; tail compressed, subtriangular, about one fourth of the total. Head shields normal. Internasals as long or longer than wide, narrow in front. Nasal divided obliquely, nostril near the upper margin. One anteorbital. Postorbitals two (2—3). A large temporal in contact with the orbitals. Labials eight (7—9), the two preceding the last large. Eye medium, pupil round. Infralabials ten (9—11). Hinder pair of submentals larger. Scales keeled, truncate or notched at the end, in 19—21 rows, outer broader, keeled to smooth. Ventrals broad 125—135. Anal bifid. Subcaudals 66—81 pairs.

Dark greyish-brown, much spotted and dotted with yellowish. Ventrals, chin-shields, and labials with central rounded or oblong spots of yellow. Flanks with faintly outlined vertical bands, appearing in parts of the body as three or four series of spots, sometimes united at the neck into as many short longitudinal bands. Head nearly black. Florida.

TROPIDONOTUS LEBERIS, *pl. II, fig. 1.*

COLUBER LEBERIS *Linnaeus*, 1758, *Syst., ed. X, l. 216; ed. XII, 1766, 375.*

TROPIDONOTUS LEBERIS *Holbrook*, 1842, *N. A. Herp. IV, 49, pl. XIII.*

Body moderately stout and long; head very little larger than the neck, depressed, crown flat; tail near one fourth of the total, tapering to a point. Head-shields normal. Rostral broad, low. Nasal divided. Mouth deeply cleft, curved posteriorly. A loreal. Anteorbitals two (2—3.) Postorbi-

tals two (2—3). Temporals 1+2. Labials seven (7—8). Eye over third and fourth, fifth and sixth larger. Infralabials 10 (9—11), fifth and sixth larger, posterior very small. Submentals two pairs, hinder larger. Scales keeled, truncate or notched at the end, in 19 rows (19—21 var. *validus*), dorsal narrow, outer broad, faintly keeled. Ventrals 140—150. Anal bifid. Subcaudals 64—86 pairs.

Reddish-brown, more or less olive, with a band of light color from the rostral along the upper labials the first and second rows of scales on the flank; below this on the first row and ends of the ventrals there is a narrow brown space. The light bands are usually margined with a dark line. Anteriorly the belly is yellowish, more or less mottled with olive, or bearing two lines of olive near the middle; posteriorly the ventrals are olive, with or without a narrow space of yellow in the middle. Chin, throat, and labials light yellow. With or without vertebral and lateral narrow lines of brown. In Kentucky specimens these lines are almost obsolete. Bright fresh ones just taken from the water have the appearance of being striped longitudinally, above and below, with narrow bands, the dark separated by lighter of about the same width, which do not form very marked contrasts, if we except the light stripe on the flank. South Carolina to California and Mexico.

rigidus. Dorsal vittae distinct; colors dark. Southeastern United States.

grahamii. Vittae distinct; bands of yellow on the flank; a light band on the middle of the back; belly yellow. Texas.

validus. Vittae obsolete; light brownish ash above; belly uniform yellowish. California to Mexico.

TROPIDONOTUS KIRTLANDII, *pl. I, fig. 3.*

REGINA KIRTLANDII Kennicott, 1856, *Pr. Ac. N. Sc., Phil.*, 95.

Small. Moderately stout, tapering little to head and tail, belly broad, back slightly compressed; head small, sub-oval, little larger than the neck, depressed, crown convex, snout short, rounded; tail short, about one fifth of the total, tapering regularly to the tip. Eye medium, pupil round. Mouth-cleft deep, curved. Head-shields nine. Parietals and frontal large. Prefrontals broad, reaching down the side of the face. Rostral broad, low. Nasal divided. A loreal. One anteorbital. Postorbitals two. Temporals 1+2. Labials 6, eye over third and fourth, fifth and sixth larger. Infralabials 7 (7—8), fifth and sixth larger. Submentals two pairs, subequal. Scales keeled, in 19 rows, dorsal narrow, truncate, outer and caudal

broad, rounded at the end. Ventrals 129—133. Anal bifid. Subcaudals 50—59 pairs.

Light greyish-brown, scales punctulate with black, with three alternating series of rounded black spots on each side. The median series of spots are the largest, four or five scales in width, about fifty from head to vent, and twenty-five on the tail. The spots of the upper series are much smaller and less distinct than those of the median; and those of the lower are still smaller and more faint. Belly yellowish, with a series of black spots near each edge. These spots are on the bases of the ventrals, a short distance from the extremity. Head to upper edge of labials dark brown, with small yellow spots on the occiput; yellow beneath. Illinois.

STORERIA.

Baird & Girard, 1853.

Small species. Moderately elongate, rounded; head short, narrow, very little larger than the neck, crown flat; tail tapering. Head-shields normal, anterior short. Loreal absent or present. Nasal in two parts, nostril between, or grooved below the nostril. Teeth equal, smooth. Scales keeled, in 15—17 rows. Anal divided or entire. Subcaudals in two rows. Hardly to be ranked higher than as a subgenus of *Tropidonotus*. Canada to Mexico.

Anal bifid;

a loreal; orbitals 2—3;

scales in 15 rows

storerioides.

No loreal; orbitals 2—2;

scales in 15 rows

occipitomaculata.

orbitals 1—2;

scales in 17 rows

dekayi.

Anal entire;

No loreal; scales in 15 rows; orbitals 1—2.

copei.

A loreal; scales in 17 rows;

nasal grooved

lineata.

STORERIA STORERIOIDES.

Tropidoclonium storerioides Cope, 1865, *Pr. Ac. N. Sc., Phil.*, 190.

Small, not slender, muzzle obtuse, in general similar to *S. dekayi*. Nasal divided or grooved. Head-shields nine. Loreal trapezoidal touching the decurved prefrontals at the superior angle, sometimes entering the orbit

between the two anteorbitals. One temporal in contact with the three postorbitals. Labials 7 (6—7), third and fourth in orbit. Infralabials 7, fourth largest. Submentals two pairs. Scales keeled, in 15 rows, vertebral narrow, outer broad, smooth. Ventrals 126±. Anal divided. Subcaudals 40 pairs.

Light to olive brown, punctulate above and below, with about fifty-four light-edged black cross-bars extending over six rows of scales, alternating with shorter ones on the sides, broken into spots on the neck. A large blotch behind the occiput on each side. Total length 13 in. 5 l.; tail 2 in. 8 l. (From descr.) Mexican Plateau, between Valley of Mexico and the Eastern Range.

STORERIA OCCIPITOMACULATA, *pl. I, fig. 2.*

TROPIDONOTUS OCCIPITOMACULATUS Storer, 1839, *Rep. Mass. Rept.*, 230.

STORERIA OCCIPITOMACULATUS Baird & Girard, 1853, *Cat. N. A. Serp.*, 137.

Small, not stout, elongate, subcylindrical, belly round; head distinct, depressed, crown slightly convex, narrow in front; tail about one fourth of the total, tapering regularly to a slender extremity. Eyes medium, pupil round. Mouth-cleft deep, slightly curved. Head-shields nine, short, broad. Rostral broader than high. Nasal divided, nostril in anterior portion. Loreal absent, rarely present (a specimen from Michigan has one on each side). Two anteorbitals. Postorbitals two. Temporals 1+2. Labials 6 (5—6), low, eye over the third and fourth, penultimate larger. Infralabials 7 (6—7), fifth largest. Submentals two pairs, posterior half as large as anterior. Scales keeled, notched at the end, in 15 rows, dorsal narrow, outer much broader. Ventrals 117—128. Anal bifid. Subcaudals 43—50.

Brownish or greyish olive. A narrow band of light color on the middle of the back, three scales in width, inclosed by a pair of narrow black lines; on the inferior row there is a similar line or vitta. Closely examined, the scales are punctulate or mottled with black. The middle of the belly is yellowish white, red in life, and becomes more olive toward the flanks. Behind each angle of the mouth, and on the occiput, there is a spot of light color. Under the head and neck the scales are more or less punctulate or clouded with brown. The markings vary from very distinct to obsolete. Mississippi Valley and Eastward.

STORERIA DEKAYI *pl. I, fig. 1.*TROPIDONOTUS DEKAYI *Holbrook, 1842, N. A. Herp. IV, 53, pl. XIII.*STORERIA DEKAYI *Baird & Girard, 1853, Cat. Serp., 135.*

Small. Body elongate fusiform, belly broad, rounded; head distinct, depressed, rounded behind, narrow and angular in front; tail short, about one fifth of the total, tapering gradually, slender near the extremity. Eye moderate, pupil round. Snout prominent. Rostral broader than high. Head-shields normal. Frontal short, broad. Nostril in hinder edge of anterior portion of the nasal. No loreal. One anteorbital. Postorbitals two, sometimes fused. One temporal in contact with the orbitals. Labials 7, eye over the third and fourth. Infralabials 7, fourth and fifth large. Submentals two pairs, subequal. An elongate shield behind the fifth and sixth infralabials. Scales keeled, slightly notched at the end, in 17 rows, dorsal narrow, outer very broad. Ventrals 120—138. Anal bifid. Subcaudals 40—60.

Light brownish olive, ashy to reddish, with two series of small black spots, about four scales apart, along the middle of the back. The spots are irregularly placed, and often united across the intervening space, forming short transverse bands of about seven scales in width. The spaces between these spots is lighter than the flanks, and separated from them by a dark line connecting the outer edges of the blotches. In the first pair the spots are larger, and extend from the occiput down the back of the head around the angles of the jaws. Head darker, a dark band across the occipitals to the hinder labial, sometimes broken on the sides. A vertical band of black under the eye. Often there is a second series of indistinct spots on the flank below the dorsal vitta, alternating with the vertebral series. Belly uniform, or with irregularly placed dots of black near the sides. Chin and throat more yellow. Spots sometimes obsolete. Maine to Mexico.

A specimen from Jalapa, Mexico, has 145 ventrals, a bifid anal, and 44 pairs of subcaudals. It is only by a close examination that it can be distinguished from Massachusetts specimens.

STORERIA COPEL.

ADELOPHIS COPEL *Cope, 1879, Pr. Am. Phil. Soc., p. 265 (Duges MSS.)*

Head little larger than neck; tail pointed, near one fifth of the total. Rostral not produced, projecting very slightly above the level of the muzzle. Internasals small, triangular. Frontal longer than wide, hexangu-

lar. Prefrontals in contact with labials. Supraoculars elongate, straight. Occipitals large. Nostril in posterior border of anterior portion of nasal. No loreal. One anteorbital. Postorbitals two. Temporals 1+2. Labials 5, third and fourth in contact with orbitals and eye. Infralabials 5 (6). Submentals two pairs. Scales keeled, truncated, in 15 rows, outer larger and smooth. Anal entire.

Light brown. A yellow band two scales in width along the vertebral rows, bordered on each side by a reddish-brown band of equal width, below which passes a narrow black line, ending in a black spot behind the eye. Posterior margins of outer row marked with black. Top of head reddish-brown. Lips and flanks yellow. Guadalajara, Mexico. (From descr.)

STORERIA LINEATA, *pl. I, fig. 4.*

MICROPS LINEATUS *Hallowell*, 1856, *Proc. Ac. N. Sc., Phil.*, p. 241.

Small. Stout, subcylindrical, belly broad, rounded; head indistinct, small, slightly convex on the crown; tail short, thick. Eye small, over third and fourth labials, pupil round. Mouth-cleft deep, curving near the angle. Snout narrow, rounded. Head-shields nine. Prefrontals reaching down on the side of the face. Rostral small, low, a little broader than high. Nasal grooved below the nostril. Loreal elongate, low. Orbitals 1+2. Temporals 1+1+2. Labials seven, sixth wedged between the fifth and seventh, not reaching the lip. Infralabials six, fourth largest. Submentals two pairs, posterior half as large as the anterior. Scales keeled, in 17 rows, median narrow, notched, outer two broad, smooth, or faintly keeled. Ventrals 138—145. Anal entire. Subcaudals 32—35 pairs.

Light greyish-brown. A light line from the occiput to the end of the tail on the medial three rows of scales. The base of each scale in the row next to this is black, forming two lines of dots. A white line occupies half of each of the second and third outer rows. Bases of the scales black in the first, second, fourth, and sometimes fifth rows. Under the epiderm the appearance is much darker. Belly yellowish white; on each side of the middle a series of black spots on the bases of the ventrals, more or less confluent across the belly, and meeting on the neck. Head mottled with black. Kansas to Texas.

HELICOPS.

Wagler, 1830.

Form stout, tapering to head and tail; head larger than the neck, broad behind, narrow forward, crown flat; tail tapering. Eyes moderate to small, pupil round. Teeth smooth, post maxillary larger and often isolated. Head-shields eight or nine. Internasals narrow in front or fused, generally in contact with the rostral. Nasal entire and grooved, or divided. With or without a loreal. Commonly 1—2 ante, and 2—3 postorbitals, suborbitals rarely present. Scales keeled in 19—25 rows. Anal bifid. Subcaudals in two rows.

Nasal divided;

internasals 2; scales keeled, in 19—21 rows

variabilis.

Nasal entire, grooved;

internasal 1; scales in 19 rows, keels few

allenii.

HELICOPS VARIABILIS.

TRETANORHINUS VARIABILIS *Dun. & Bibr.*, 1854, *Erp. Gén.* VII, 349.

Fusiform; head distinct, narrow across the nasals, broad behind, depressed, crown flat; tail moderate, tapering rapidly near the base. Eye small, pupil round. Jaws nearly equal in length. Posterior maxillary teeth little larger, slightly compressed. Rostral broader than high, convex. Head-shields nine. Internasals small, irregular, subtriangular, not at all or but little in contact with the rostral. Prefrontals broader in front, longer than broad. Frontal pentagonal, truncate and narrower in front. Nasal in two parts, the small nostril between, nearer the upper margin. Loreals 2 (1—2), anterior smaller. Anteorbitals 2 (2—3), lower larger. Postorbitals 2. Temporals 1+2+3. Labials 8—9, the two preceding the last largest, a single one (fourth or fifth) under the eye. Infralabials 10—11, sixth largest (fifth or sixth). Submentals two pairs, posterior larger. Scales keeled, lozenge-shaped, truncate, in 21 rows (19—21), dorsal finely striate, outer broader, nearly or quite smooth. Ventrals broad 142 (137—158). Anal bifid. Subcaudals 65 pairs (65—78).

Light greyish to dark brown, edges of scales lighter; young with irregular indistinct blotches of darker. A light line on the outer two rows of scales from the neck to the tip of the tail. Belly yellowish white, mottled

with brown, darker on throat and chin. A dark border to the lower edge of the light line on the flank is not always present. A dark band passes from the nasals through the eye to the neck. One of the specimens from which this description is taken has been named by Prof. Jan *Helicops Agassizii*. Hab. Mexico and Central America.

HELICOPS ALLENII, *pl. VII, fig. 4.*

Garman, 1874, Proc. Bost. Soc. Nat. Hist., XVII, 92.

Fusiform, stout, slightly compressed, belly rounded; head small, crown slightly convex and tapering to a point at the rostral; tail moderate, tapering rapidly near the base. Eye large, pupil round, situated in advance of the middle of the length of the mouth. Mouth-cleft deep, slightly curved. Posterior maxillary teeth larger. Head-shields eight. Internasals fused, very small. Frontal slightly narrower forward. Occipitals broad, in contact with the labials. Supraciliaries elongate, narrow in front. Prefrontals broad, reaching down on the side of the face. Rostral small, broader than high. Nasals entire, in contact between rostral and prefrontals. Nostril small, superior, a groove from its lower edge to the labial. Loreal quadrangular. One anteorbital. Postorbitals three, lower small. Temporals 1+2 (not in contact with the orbitals). Labials 8, sixth and seventh large, hinder angle of third and the fourth entering the orbit. Infralabials 10 (10—11), fifth and sixth larger. Submentals two pairs, hinder larger. Scales smooth, excepting several rows on the tail, lustrous, in 19 rows, vertebral twice as long as wide, outer wider than long, the medial two rows of the tail strongly and the next pair slightly keeled. Ventrals 128. Anal bifid. Subcaudals 58 pairs.

Dorsal rows light reddish-brown (yellowish to flesh color on the flanks). A vertebral band of rich dark brown near five scales in width, the middle of which is perceptibly lighter. A band of lighter brown on each flank, two scales and two half scales, separated from the vertebral by a band of the ground color two half scales wide. Lateral edges of each scale darker. Labials, chin and throat yellowish. Head of a uniform rich dark brown. Florida.

HYDROPS.

Wagler, 1830.

Elongate fusiform to subcylindrical; head rather indistinct, short, broad, rounded; tail short, thick, pointed. Eye moderate, pupil round. Teeth smooth, in continuous series, or posterior maxillary larger, sometimes iso-

lated, rarely grooved. Head-shields 8—9. Internasals 2 or fused. Nasal grooved or divided. Loreal rarely present. Orbitals 1—2. Scales smooth, lustrous, in 15—21 rows. Anal bifid. Subcaudals in two rows.

Nasal grooved;

no loreal;

scales in 19 rows;

head-shields normal

erythrogrammus.

internasals fused

abacurus.

Nasal divided;

a loreal;

scales in 21 rows;

head-shields normal

quinquevittatus.

HYDROPS ERYTHROGRAMMUS.

COLUBER ERYTHROGRAMMUS *Daudin*, 1799, *Hist. Rept. VII*, 93, *Tab. 83, f. 2*.

Stout, round, subfusiform; head hardly distinct from neck, slightly depressed, rounded; tail short, one sixth to one eighth of the total, thick, tapering to a point. Snout moderate, rounded. Eye medium, over the third and fourth labials, pupil round. Mouth deeply cleft, outline nearly straight. Teeth larger backward. Head-shields normal. Rostral broad, convex. Internasals small. Prefrontals broad, entering the orbit. Frontal hexagonal, lateral sides nearly parallel. Anteorbital and loreal united. Postorbitals two. Temporals 1+2. Labials 7, penultimate largest. Infralabials eight, fifth larger. Submentals two pairs, posterior smaller. Scales smooth, glossy, in 19 rows, dorsal broad, outer broader than long. Ventrals broad 167—185. Anal bifid, exceptionally entire. Subcaudals 38 to 50 pairs.

Red or reddish brown, with four bands of lustrous dark-brown or black from head to tail, each about three scales in width, separated by red spaces of about one scale on the vertebral and the sixth rows on each flank. Outer row, sometimes the first two or three rows, red. Lateral brown band often narrowed, lower margin notched. A rounded spot of black near each end of the ventrals. A row of spots along the middle of the belly, sometimes broken or absent. Labials and chin-shields with central spots of black. Head-shields more or less marked with red. Largest specimen $39\frac{1}{2}$ inches in total; tail $5\frac{1}{2}$ inches. Mississippi Valley and eastward, as far north as Illinois and North Carolina.

HYDROPS ABACURUS, *pl. I, fig. 5.*COLUBER ABACURUS *Holbrook*, 1836, *N. A. Herp.* 1st ed., I, 119, *pl. XXIII.*HYDROPS ABACURUS *Dum. & Bibr.*, 1854, *Erp. Gén.*, *Tab.* 65.

Long, moderately stout, cylindrical; head very little larger than the neck, depressed, rounded; tail short, thick, conical, one sixth to one ninth of the total. Snout short, broad. Eye medium, over third and fourth labials, pupil round. Mouth-cleft deep, slightly curved. Teeth little larger backward. Head-shields eight. Internasals united. Prefrontals reaching the orbit. Frontal hexangular. Nasal entire, grooved from the nostril to the first labial. Nostril small, visible from above. Loreal united with anteorbital. One anteorbital. Postorbitals two, lower small, in a notch between the labials. Temporals 1+2. Labials seven, sometimes fused or divided, penultimate larger. Infralabials nine, fifth larger. Submentals two pairs, hinder small. Scales smooth, glossy, rounded at the end, in 19 rows, outer broader. Ventrals large, broad, 171—203. Anal bifid. Subcaudals in 35—47 pairs.

Ground color red, yellow in alcohol. Back and usually top of head bluish-black. The black color extends down the flanks in wedge-shaped points, separated by wedges of the red, which sometimes extend to the middle of the back. The bands of black extend nearly or quite to the middle of the belly, where they meet or alternate with those from the opposite side. Red occasionally encircling the neck or scattered over the top of the head. Anterior labials and chin-shields, with rounded spots of black. Largest specimen 54 inches long; tail $5\frac{5}{8}$ inches. Mississippi Valley and eastward, as far north as Illinois and Virginia.

HYDROPS QUINQUEVITTATUS.

HOMALOPSIS QUINQUEVITTATUS *Dum. & Bibr.*, 1854, *Erp. Gén.* VII, 975.

Elongate, moderately stout, subcylindrical; head distinct, little larger than the neck, depressed, crown flat, muzzle narrow between the nostrils; tail stout, tapering regularly to the point. Eye moderate, pupil round. Mouth-cleft deep, outline curved. Posterior maxillary teeth longer, separated by an interspace. Head-shields normal. Internasals small, narrow in front. Frontal short, broad, truncate, or angled in front. Nasal divided or entire. Loreal half as large as the anteorbital. Postorbitals two. Temporals 1+2. Submentals two pairs, nearly equal, posterior diverging.

Labials 8, sixth and seventh largest. Infralabials 10. Scales smooth, lustrous, in 21 rows. Ventrals 160. Anal bifid. Subcaudals 65 pairs.

A band of black along each flank from the nostril to the tip of the tail. One or two of the outer rows lighter. Dorsal region covered by a band of greyish brown, lighter near the black bands. Labials, infralabials, and chin-shields with rounded spots of black. Each ventral has a rounded spot upon the base near each end which form a series along each side of the abdomen. Southern Mexico and Central America.

COLUBRINAE.

Elongate fusiform, stout to slender; head distinct from the neck, elongate, crown flattened, sides concave; neck somewhat small; tail rather long, not distinct from the trunk, tapering. Muzzle stout, produced. Mouth deeply cleft, outline curved. Teeth subequal or posterior larger. Eye large, pupil round. Scales smooth or keeled, imbricate; subcaudals in two rows.

The typical genera of the family are those of *Coluber* and *Elaphis*, the genera to which the common black snakes of the United States belong.

Nasal divided;

edges of rostral free; scales smooth, in 17 series;
posterior maxillary teeth longer, isolated, smooth

SALVADORA.

Nasal entire;

teeth equal, smooth; loreal present or lacking;
anal bifid; scales smooth or keeled, in 15 or 17 series

CYCLOPHIS.

Nasal divided;

anal bifid; scales smooth or keeled; loreal present;
teeth becoming longer posteriorly

COLUBER.

anal entire;

teeth equal smooth; a loreal; crown-shields nine;
scales keeled or smooth, in 15 to 23 series

SPILOTES.

scales keeled, in 27 to 35 series;

crown-shields nine to eleven

PITYOPHIS.

anal bifid or entire;

teeth equal, smooth; a loreal; crown-shields nine;
scales keeled, sometimes obsolete, in 19 to 25 series

ELAPHIS.

anal bifid:

posterior maxillary teeth longer, smooth; loreal rarely absent;
crown-shields nine

DROMICUS.

SALVADORA.

Baird & Girard, 1853.

Elongate, slender; head distinct from neck, long, snout produced; tail slender, tapering gradually. Eye large, pupil round. Posterior maxillary teeth larger, smooth, isolated. Head-shields nine. Rostral prominent, bent back on the head, edges more or less free. A loreal. Nasal divided. Anteorbitals two to three. Postorbitals two to three. Scales smooth, generally in 17 rows. Anal bifid. Subcaudals in two rows.

Labials forming the lower edge of the orbit;

with longitudinal bands *grahamii.*

Orbit separated from the labials by small shields;

with a dorsal series of blotches *decurtata.*

SALVADORA GRAHAMIAE.

Baird & Girard, 1853, Cat. N. Amer. Serp., 104.

Slender, fusiform; head moderate, distinct from the neck, with flat crown; tail about one fourth of the total, slender. Eye large, pupil round. Snout produced, truncate, inclined from a vertical. Rostral prominent, with free edges, bent back on the top of the snout. Crown-shields nine. Internasals broad, separated in front by the rostral. Prefrontals broad. Supraciliaries and frontal long. Nasal divided, nostril lateral near the margin of the prefrontal. Loreals one to three. Anteorbitals two. Postorbitals three to two. Labials nine, eye over the fifth and sixth, seventh largest. Temporals two, followed by irregular shields. Infralabials nine to ten, fifth largest. Posterior submentals small. Scales smooth, in 17 rows, medial narrow, outer broad. Ventrals 175—186. Anal bifid. Subcaudals 75 to 106 pairs.

Top of the head yellowish brown, from this to the end of the tail extends a medial band of dull yellowish one to three scales in width. On each

side from the nasal through the eye along the flank there is a black band three or more scales in width. Below the black a band of yellowish green a scale or more in width separates it from a second dark band about half a scale in width; one or the other of these is often obsolete. Outer rows dull yellowish green. Belly uniform dull yellowish. Mexico.

VAR. HEXALEPIS.

PHYMOTHYRA HEXALEPIS Cope, 1866, *Proc. Ac. N. Sc., Phil.*, 304.

Is shorter, and has the loreal divided.

SALVADORA DECURTATA.

PHYMOTHYRA DECURTATA Cope, 1860, *Proc. Ac. N. Sc., Phil.*, 310.

"This species is distinguished by many features. The head is shortened and somewhat arched, the rostral plate very broad and free, entirely separating the internasals. There is a complete annulus of scales around the eye. The tail is relatively shorter than in the known species.

"Ground color light grey; a series of elongate parallelogramic brown blotches occupies the dorsal region, from the nape to the end of the tail. Labials and under surfaces unspotted. Length about 14 inches. This serpent is remarkably different from the three already known representatives of the genus. In these the orbit rests on the labials, and the color is in bands." Upper part of Lower California.

CYCLOPHIS.

Günther, 1858.

Body slender, belly rounded; head ovoid, distinct from the neck; tail long, tapering regularly. Eye somewhat large, pupil round. Teeth equal, smooth. Head-shields nine. Loreal present or lacking. Nasal entire. Orbitals 1—2. Scales smooth in 15 rows, or keeled in 17. Anal and subcaudals bifid.

CYCLOPHIS VERNALIS, *pl. III, fig. 4.*

COLUBER VERNALIS (De K.) *Harlan*, 1826, *Jour. Ac. N. Sc., Phil.*, V, 361.

CYCLOPHIS VERNALIS *Günther*, 1858, *Cat. Col. Snakes*, 119.

Moderately slender, subcylindrical, with flat belly. Head moderate, distinct from the neck, sides concave in front of the eye. Tail rather stout, near one third of the total, tapering. Eye medium, over the third

and fourth labials; pupil round. Head-shields nine. Supraciliaries prominent. Frontal pentagonal. Muzzle prominent, extending beyond the lower jaw. Rostral bent back and sharp-angled between the internasals. Nasal entire, nostril near the center. Loreal present. Orbitals 1—2, lower small. Temporals 1+2. Labials seven, sixth larger. Infralabials eight, fifth larger. Posterior submentals shorter. Scales smooth, in 15 rows, medial narrow, outer broad. Ventrals 125—140. Anal bifid. Subcaudals 69—95 pairs.

Grass green, whiter below.

(*Phyllophilophis.*)

CYCLOPHIS AESTIVUS, *pl. III, fig. 1.*

COLUBER AESTIVUS *Liné*, 1766, *Syst. Nat.*, ed. XII, vol. I, 387.

CYCLOPHIS AESTIVUS *Güther*, 1858, *Cat. Col. Snakes*, 119.

Body long, slender, compressed, with small neck and flat belly. Head distinct from the neck, elongate, with flat crown and pointed snout. Tail slender, more than half as long as the body. Eye large, pupil round. Mouth deeply cleft. Head-shields normal. Vertical pentagonal. Snout prominent; rostral broad, reaching the top. Nasal entire, elongate, nostril in the middle. Loreal present. Orbitals 1—2, lower smaller. Temporals 1+2. Labials seven, third and fourth in the orbit. Infralabials eight, fifth very large. Posterior submentals longer. Scales narrow, in 17 rows, keeled except on the outer or outer two, which are broader and smooth. Ventrals 150—165. Anal bifid. Subcaudals 110—135 pairs.

Green, blue in alcohol, often with a reddish tint; belly white, greenish posteriorly.

COLUBER.

Liné, 1754, *Mus. Ad. Fridr.*; *Syst. Nat.*, ed. X, 1758, I, 216.

Elongate, moderate to slender, slightly compressed; head distinct from the neck, somewhat narrow, crown flat, subquadrangular in transverse section at the eye; tail moderate to long and slender. Eye moderate, pupil round. Teeth becoming larger and longer posteriorly. Crown-shields nine. Nasal divided. Loreal present, sometimes divided. Preoculars one to two. Postoculars two to three. Scales rhomboid, smooth or keeled, in 14 to 23 rows. Anal bifid. Subcaudals in two rows.

(*Basanium.*)

Scales in 17 rows;

anal entire;

back plumbeous to black, in adults;

two labials in the orbit

constrictor.

one labial in the orbit

mentocarius.

(*Masticophis*.)

Anal bifid;

dingy white, mixed with brown, varying to black on the anterior half of the body

flagelliformis.

reddish-brown; anterior scales darker in the center, posterior with a light spot at the base

testaceus.

color in longitudinal bands;

three preoculars; brown darkening anteriorly

aurigulus.

one preocular;

bands broken into transverse blotches anteriorly

mexicanus.

a yellow band on vertebral rows; a black line at ends of ventrals

spinulis.

Scales in 15 rows;

a broad band of dark color on the back; flanks with narrow lines of light color, separated by dark.

taeniatus.

COLUBER CONSTRICTOR, *pl. IV, fig. 3.*

Linné, 1758, *Syst. Nat.*, ed. X, I, 216; 1766, ed. XII, I, 385.

Elongate, slender, slightly compressed; head distinct from the neck, long, narrow, angled in front, concave on the sides, crown flat, curving downward near the snout; tail about one fourth of the total, slender. Eyes large, over the third and fourth labials, pupil round. Mouth-cleft deep, curved. Crown-shields normal. Prefrontals bent downward on the side of the head. Rostral broad, very convex. Nasal in two parts, nostril between. A loreal. Preoculars two, sometimes but one, lower very small, in a notch between the second and third labials. Postoculars two (2—3), lower resting on the fourth labial. Temporals 2+2+2. Labials seven (7—8), fourth, sixth, and seventh larger. Infralabials nine, fifth very

large, posterior two small. Submentals two pairs. Scales smooth, in 17 rows, medial narrow, outer and caudal as broad as long. Ventrals 175—190. Anal entire. Subcaudals 80—110 pairs.

Uniform lustrous black above, varying to olive or leaden, more or less yellow, on the western prairies; chin, throat, and portions of the labials white. The white extends upward behind the jaws, or is limited to the chin.

The young are olivaceous with a dorsal series of large, darker-edged, irregular-shaped brown spots, the largest of which forms a transverse band immediately behind the head. Sides and abdomen with irregularly placed spots of brown, varying in size from that of a dot to that of a couple of scales. The spots become obsolete backward. Head yellowish brown above, spotted or mottled with dark, below yellowish. Mississippi Valley and eastward.

var. FLAVIVENTRIS.

Say, 1823, *Long's Exp.* II, 185.

The "Blue Racer" of the prairies is rather stouter than the "Black Snake" of the Eastern States. On the back it is light olive, greenish or blueish; beneath yellowish, some of the scales and interspaces showing darker. The young are mottled like those of the black snake and without spots below.

var. MENTOVARIUS.

CORYPHODON MENTOVARIUS *Duméril & Bibron*, 1854, *Exp. Gén.* VII, 187.

Back greenish-brown. A single labial entering the orbit. This species resembles greatly *C. constrictor*; it is distinguished by the single labial beneath the eye. The head is comparatively more elongate, and the eye larger. The hinder submentals are longer than the anterior, which is not the case in *C. constrictor*. (Duméril.) Mexico.

COLUBER FLAGELLIFORMIS.

Hillbrook, 1836, *N. A. Herp.*, ed. I, 107, pl. XIX.

Long, slender, with flat belly; head distinct, elongate, narrow, subquad-rangular in transverse section, as deep as wide at the eyes, crown flat, bent downward slightly in front of the orbits; tail long, near one fourth of total, slender, tapering regularly to the tip. Eyes large, over the fourth and

fifth labials, pupil round. Snout prominent, produced, rounded. Mouth-left deep, curving. Teeth equal, smooth. Sides of the head concave in front of the eyes. Crown-shields nine, large, elongate. Rostral convex, little bent back between the internasals. Prefrontals large, slightly decurved upon the side. Frontal long, broad in front, becoming narrow at the middle, sides parallel posteriorly. Supraciliaries large, convex, prominent over the eye. Parietals broad. Nasal in two parts, nostril between. Loreal higher than long. Two preoculars; upper large, projecting over the loreal, upper angle in contact with the frontal; lower small in a notch between the third and fourth labials. Postoculars two, lower resting on the fifth labial. Two or three temporals in contact with the orbitals. Labials eight, posterior two largest. Infralabials ten, fifth largest. Submentals two pairs, subequal. Scales smooth, in 17 rows, outer broader. Ventrals 200—206 (with varieties 190—206). Anal bifid. Subcaudals 94 to 111 pairs. Frequently there is a short incision in the upper preocular near the upper edge of the loreal, sometimes this is met by another, cutting out a triangular piece of the shield.

Young whitish, uniform posteriorly, with narrow transverse bands of brownish anteriorly. The bands darken toward the head, which is banded and clouded with brown, darkest on the supraciliaries.

Adult dark anteriorly—brown to black, mixed white and brown medially, and whitish posteriorly. Belly dark anteriorly, clouded with brown, tinged with red in the middle and white posteriorly. Upper anteorbital with a yellow spot next the eye. The species varies considerably in the depth of the color of the anterior half of the body and in the amount of yellow about the chin. Mississippi river and eastward.

var. TESTACEUS.

COLUBER TESTACEUS. Say, 1823, *Long's Exp.* II, 48.

Elongate, tapering to head and tail, head a little larger than the neck, long, quadrangular in transverse section at the eyes, with concave sides, arched from the snout backward; tail near one fourth of the total, tapering slender. Eye large, pupil round, brows overhanging. A couple of the posterior maxillary teeth a little longer. Crown-shields normal. Prefrontals broad anteriorly, bent downward at the sides. Frontal broad in front, narrow in the posterior half. Supraciliaries broadening backward. Nasal divided. Loreal small, lozenge-shaped. Preoculars two; lower

small, in a notch between the third and fourth labials; upper large, reaching the frontal. Postoculars two. (In this specimen there is a large temporal between the truncate parietals and the posterior labials, and in front of it there are three small ones, two of which are in contact with the orbitals.) Labials eight, fourth and fifth entering the orbit, last two larger. Infralabials ten, fifth large. Submentals two pairs, subequal. Scales moderate, smooth, in 17 rows, outer broader. Ventrals broad, 191—198. Anal bifid, occasionally entire. Subcaudals 80—108 pairs.

Reddish-brown, more red below, lateral edges of scales lighter, each scale with a brown line through its middle, darkening toward the tip. On the hinder portion of the body each scale has a light spot at its base, and the margin is brown. Head brown, more or less yellow or mottled with yellow. Labials, chin, and throat yellow, more or less blotched with brown. Usually there are two rows of brown spots on the shields of the throat, sometimes extending under the body. In a specimen stripped of the epiderm there are indistinct narrow transverse lines of darker on the back. Total length 47 inches; tail $12\frac{1}{2}$ inches. Arkansas to Mexico.

VAR. AURIGULUS.

DRYMOBIUS AURIGULUS Cope, 1861, *Pr. Ac. N. Sc., Phil.*, 301.

Crown flat, muzzle elongate. Eye moderate. Rostral plate rounded, prominent, recurved above. Frontal elongate, posteriorly half as wide as each supraciliary, not in contact with the preocular. Parietals elongate. Nasals and loreal very long, the latter encroaching much on the preocular. Three pre, two postoculars. Labials eight, fourth and fifth entering the orbit; the last equal in length and elevation to the penultimate. Infralabials ten, fifth largest. Anterior submentals shorter. Scales in 17 rows.

Brown, darkening anteriorly. Head-plates light brown, shaded with yellow. A narrow yellow band around the muzzle from eye to eye. Temporal region and postoculars each with a spot. Labials, chin, and anterior portion of abdomen bright golden, as also the sides of the neck to the fifth row of scales. On the second and third rows of scales of the neck there is a black band, interrupted at intervals of about seven scales. It finally becomes continuous, and with a band on the first row almost excludes the ground color from the posterior and middle parts of the body. Abdomen dirty yellowish. Cape St. Lucas, Lower Cal.

COLUBER MEXICANUS.

ZAMENIS MEXICANUS *D. & B.*, 1854, *Erp. Gén.*, 695.

Long, slender, belly broad; head distinct from neck, elongate, depressed, crown flat, sides concave at the eye; tail near one third of the total, slender, tapering regularly. Mouth-cleft deep, sinuous. Eye moderately large, pupil round. Crown-shields normal. Rostral prominent, higher than wide, bent back between the internasals in an acute angle. Prefrontals broader than long. Frontal long, sides concave. Supraciliaries large, prominent. Nasal divided. Loreal present. Preoculars large, considerably in contact with the anterior angles of the frontal. Two postoculars, lower resting on the sixth labial. Temporals 2+2+2. Labials nine, fourth, fifth, and sixth in the orbit, hinder four larger. Infralabials ten, sixth largest. Submentals two pairs, nearly equal. Scales smooth, in 17 rows, dorsal narrow, lateral as broad as long. Ventrals 182. Anal bifid. Subcaudals 128 pairs.

Light brownish or yellowish-brown. Posteriorly with four black bands on the dorsal rows, each of the width of two scales, separated on the flank by a space of equal width and by the median row of scales on the back. Forward the bands are broken up and each scale of the dorsal rows is more or less marked with black. The first band occupies a portion of the outer row and all of the second. Anteriorly the back is crossed by transverse bands of brown, the first of which crosses the head just behind the parietals and meets the bands from the eye. Crown-shields broadly margined with brown. Parietals divided longitudinally by a band, a dot of brown on the yellow portion of each near the middle. Total length $31\frac{1}{2}$ inches; tail $10\frac{3}{4}$ inches. San Blas, Jalisco, Mexico.

COLUBER SPINALIS.

MASTICOPHIS SPINALIS *Peters*, 1866, *Monatsber. Berl. Akad.*, 91.

Very long and slender; head twice as long as broad, snout prominent, rounded; tail near one fourth of the total. Posterior maxillary tooth longer, smooth. Crown-shields nine. Prefrontals broader than long, not longer than the internasals. Nasal in two parts, nostril between, above the middle. Loreal trapezoidal, longer than broad. Two preoculars; upper large; lower small, in a notch between the third and fourth labials. Frontal long, broad and blunt-angled in front, concave at the sides. Parietals hardly longer than the frontal. Supraciliaries narrow in front,

broader than the frontal posteriorly. Temporals 1+2. Labials eight, fourth and fifth in the orbit. Infralabials ten, fifth largest. Submentals two pairs, elongate. Scales smooth, in 17 rows, dorsal with two pores, narrow, outer twice as broad. Ventrals 203. Anal bifid. Subcaudals 96 pairs.

Olive-tinted greenish-brown above; a yellow black-edged band one and two half scales in width along the middle of the back. A blackish line on the ends of the ventrals and edge of the outer row. Above the dark edges a light space on the outer row extends along body and tail as a light line. A yellow cross band on the hinder extremity of the prefrontals, the ante and postoculars. The greater part of the supralabials and ventral surface yellow. Mexico.

COLUBER TAENIATUS.

LEPTOPHIS TAENIATA *Hallowell*, 1852, *Proc. Ac. N. Sc., Phil.*, 181.

Very long and slender; head distinct from the neck, elongate, narrow, subquadrangular in transverse section, crown flattened, sides concave in front of the eye; tail slender, long, one third of the total. Snout prominent. Mouth-cleft deep, outline curved. Teeth equal, smooth. Eye large, pupil round. Crown-shields nine, large, long. Rostral bent back in a blunt angle between the internasals. Prefrontals broad, bent downward at the sides. Frontal broad anteriorly, contracted in the middle. Supraciliaries long, broad, prominent, convex. Nasal divided, nostril between. Loreal longer than high. Preoculars two; lower very small, in a notch between the third and fourth labials; upper produced above the loreal, sometimes reaching the vertical. Postoculars two, lower on the fifth labial. Two temporals in contact with the orbitals. Labials eight, seventh largest. Infralabials ten, fifth largest. Submentals two pairs, the hinder longer. Scales smooth, lozenge-shaped, truncate at the end, in 15 rows, outer broad. Ventrals 209. Anal bifid. Subcaudals 135—157 pairs.

A narrow longitudinal line of brown or black in the middle of each row of scales. A dorsal band of brown about seven scales in width obscures those of the medial rows; those on several of the outer rows are black. At the upper edge of each flank there is a narrow band about one scale in width of orange or red. The longitudinal lines on the outer rows are more or less distinctly separated by spaces of orange. A rather indistinct

band of reddish-brown extends along each side of the abdomen, separated from the first brown line by a space of yellow. Belly and throat yellow, spotted with brown anteriorly. Orbitals and crown-shields with yellow margins or spots. Wyoming to Texas, California and Mexico.

SPILOTES.

Wagler, 1830.

Body long, compressed; head distinct from the neck, muzzle rounded; tail moderate. Eye large, pupil round. Teeth equal, smooth. A loreal. Anteorbital one, rarely two. Postoculars two to three. Scales keeled or smooth, in 15—23 rows. Anal entire. Subcaudals in two rows or simple.

(*Georgia.*)

Scales smooth, in 17 rows; subcaudals generally simple;

black; labials seven; ventrals 184—190 *couperi.*

black, with reddish-white spots; labials eight *obsoletus.*

(*Spilotes.*)

Scales smooth, in 17 rows, rarely 15; subcaudals usually in two rows;

brown; lighter anteriorly, with faint indications of transverse zigzag bands of dark; ventrals about 205 *corais.*

Scales faintly keeled, in 15 (15 to 17) rows;

yellowish, marked with chevron bands of brown; darker posteriorly; ventrals about 220 *variabilis.*

16 rows of scales;

posterior two thirds of body black *melanurus.*

18 or 19 rows of scales;

hinder third of body with chevron bands of yellow *auribundus.*

21 to 23 rows of scales;

uniform brown above, a few vertebral scales yellow with black tip *poecilonotus.*

SPILOTES COUPERI.

COLUBER COUPERI *Holbrook*, 1842, *Herp. N. Amer.* III, 75.SPILOTES COUPERI *Cope*, 1860, *Pr. Ac. N. Sc., Phil.*, 342 and 564.

Elongate, stout; head distinct from the neck, high, crown arching longitudinally; tail short, near one sixth of the total. Crown-shields nine. Rostral broader than high. Frontal about as broad as long. Nasal in two parts, nostril between. A loreal. Oculars 1—2, anterior large. Labials seven, anterior smaller, eye over the fourth and fifth, fourth supporting the postorbital, fifth small triangular, sixth very large, extending forward over the fifth, seventh large. Scales lustrous, imbricate, smooth, in 17 rows. Ventrals 184—190. Anal entire. Subcaudals 60, not divided.

Lustrous black. Belly more or less plumbeous; yellowish under the throat and neck. The young are probably more yellow on the anterior portions. Georgia, Texas.

SPILOTES OBSOLETUS.

COLUBER OBSOLETUS *Holbrook*, 1842, *N. A. Herp.* III, p. 61, pl. XII.

Body long, stout, compressed; head distinct from the neck; tail near one sixth of the total. Eye medium, over the fourth and fifth labials. Labials eight, fifth and seventh not in contact, sixth small, wedge-shaped. Infralabials nine, fourth and fifth largest. A loreal, sometimes two. Orbitals 1—2, posterior resting on the fifth labial. Scales in 17 rows. Ventrals 193. Subcaudals 56—60.

Black, plumbeous below; anterior half of the body with reddish-white spots on the bases of some of the dorsal scales, and on the bases of ventrals. On the sides and under the head yellowish-red. Posterior margins of labials and lower plates of the head black. Texas.

SPILOTES CORAIS.

COLUBER CORAIS *Daudin*, 1803, *Hist. Rept.* VII, 23.SPILOTES CORAIS *Dun. & Bibr.*, 1854, *Erp. Gén.* VII, 223.

Elongate, stout, somewhat compressed; head large, subquadrangular in cross section at the eye, crown flat; tail rather less than one fifth of the total, tapering. Eye moderate, pupil round. Mouth-cleft deep, curved. Teeth gradually enlarging backward, smooth. Snout prominent, broad, rounded. Crown-shields normal. Internasals subquadrate. Frontal and prefrontals short; the latter bent down on the sides of the face to the loreal. Rostral broader than high. Nasal divided. One preocular, broader above.

Postoculars two, small. Temporals 2+2, upper anterior small. Labials eight, anterior four small, fourth and fifth under the eye, sixth triangular, seventh and eighth very large, the last separated from the parietals by a pair of large temporals. Infralabials nine, fourth and fifth larger. Submentals two pairs, subequal. Scales large, smooth, in 17 (15 to 17) very oblique rows, outer broad. Ventrals 206, large, broad, bent up on the sides. Anal entire. Subcaudals 72 pairs.

Brown, lighter posteriorly, with faint indications of zigzag transverse bands of darker. These bands occupy the margins of the scales and are more distinct on the removal of the epiderm. Belly, chin, and throat little lighter than the upper portions. Brazil to Venezuela, possibly to Mexico.

SPILOTES VARIABILIS.

Daneril & Bibron, 1854, Erp. Gén. VII, 220.

Long, slender, compressed; head large, long, subquadrangular in section across the eye, snout broad, crown slightly convex over frontal and parietals; tail moderately slender, less than one third of the total, tapering. Eye large, pupil round. Nostril lateral. Mouth-cleft deep, curved. Teeth equal, smooth. Crown-shields normal, broad. Rostral broader than high, reaching the top of the snout, convex. Prefrontals bent down on the side of the face. Nasal divided. Loreal small. One preocular, large, reaching the top of the head. Postoculars two, equal. Temporals 1+2. Labials nine, the anterior three with the fifth and seventh small, eye over the fourth, fifth, and sixth, seventh triangular, allowing the sixth and eighth to meet above it, eighth very large. Infralabials ten, sixth larger. Submentals two pairs, equal. Scales large, much imbricate on the sides, faintly keeled, in 15 rows (irregular, 15 to 17, if small scales irregularly interspersed with the larger are counted). Ventrals 220, broad, bent up on the sides. Anal entire. Subcaudals 121 pairs.

Anteriorly white (yellow in life), marked with chevron bands of brown or black, posteriorly more dark color. Belly whitish with prolongations from the dorsal bands extending upon it. Head whitish, marked with black (a spot on each parietal and supraorbital margined with black). A large black spot on each side of the neck immediately behind the head. Total length $23\frac{1}{2}$ inches; tail $5\frac{1}{2}$ inches. Quite variable in colors and squamation. South and Central America to Mexico.

SPILOTES MELANURUS.

Dunéril & Bilron, 1854, Exp. Gén. VII, 224.

Specimens belonging to this species differ from those of *S. variabilis* in that the posterior two thirds of the body are entirely black. Temporals 1+1+2. Labials eight. Keels on the scales obsolete, faintly indicated near the middle of the body. Rows of scales 16. Ventrals 226. Anal entire. Subcaudals 123 pairs. Total length $58\frac{1}{2}$ inches; tail $15\frac{1}{4}$ inches.

SPILOTES AURIBUNDUS.

Cope, 1861, Pr. Ac. N. Sc., Phil., 300.

Labials eight, ultimate as high or higher than the penultimate, fifth large, approaching or reaching the postocular. Scales in 18 or 19 rows, weakly keeled, first four or five smooth.

Crown of head yellow, crossed by four more or less irregular cross-bands of black. Posterior third of body crossed by numerous narrow, chevron-formed cross-bands of yellow; the tail annulate with the same. Mirador, Vera Cruz.

SPILOTES POECILONOTUS.

Günther, 1858, Cat. Col. Scales in Brit. Mus., 100.

Elongate, compressed; head distinct from the neck, rounded in front; tail slender. Rostral broader than high, just reaching the top of the head. Internasals short, broader behind; prefrontals much larger. Frontal broad in front, obtuse-angled behind. Supraciliaries broad behind. Parietals short. One preocular, not reaching the frontal. Postoculars two. Loreal present. Nasal divided. Labials nine, fourth, fifth, and sixth in orbit. Scales elongate, much imbricate, in 21—23 rows, dorsal keeled, the three vertebral most strongly. Ventrals raised on the sides. Anal entire. Subcaudals in two rows.

Uniform brown above, with the exception of some scales in the medial row, which are yellow, with black tip. Head yellow and brown variegated. Belly yellow anteriorly. Ventrals black-edged about the middle of the body, becoming entirely black posteriorly. Total length 63 inches; tail 18 inches. Honduras, Mexico.

PITYOPHIS.

Holbrook, 1842, N. A. Herp. IV, 7.

Long, moderately stout, compressed; head distinct from the neck, subconical, broad posteriorly, pointed at the muzzle; tail rather short. Eye moderate, pupil round. Teeth equal, smooth. Crown-shields nine to eleven. Prefrontals one to two pairs. Nasal divided. A loreal, rarely suppressed. One to two preoculars. Postoculars two to five. Scales keeled, in 27 to 35 rows. The individual variation is very great in specimens belonging to species of this genus, particularly so in those of the Rocky Mountains.

PITYOPHIS MELANOLEUCUS.

COLUBER MELANOLEUCUS Daudin, 1799, Hist. Nat. Rept. VI, 409.

PITYOPHIS MELANOLEUCUS Holbrook, 1842, N. A. Herp. IV, p. 7, pl. I.

Long, moderately stout, slightly compressed; head moderate, distinct from the neck, appearing subquadrangular when viewed from the front, narrowing forward, conical or pointed, crown slightly convex, very broad behind; snout prominent; tail short, rather less than one seventh of the total, stout, tapering. Eye moderate, pupil round. Mouth-cleft deep, slightly curving. Crown-shields more or less irregular, commonly two pairs of prefrontals. Rostral very prominent, narrow, swollen, extending on the upper surface of the head, between the internasals. Internasals short, wide. Outer prefrontals extending on the side of the face to the loreal, inner narrow posteriorly. Frontal broad anteriorly. Supraciliaries prominent, broadening backward. Parietals large, nearly as wide as long. Nasal in two parts, nostril between. Loreal small, sometimes united with adjacent plates. One preocular. Postoculars three to four. Two or more temporals in contact with the postoculars, sometimes broken into small scales. Labials eight, fourth below the eye, fifth under the postorbital, seventh largest. Infralabials thirteen to fourteen, fifth and seventh large, hinder very small. A pair of large submentals, followed by a pair of smaller ones, which are separated by a pair of small plates. Scales lozenge-shaped, usually in 27 or 29 rows (ranging in the different varieties from 27 to 35), outer broad, lateral smooth, vertebral keeled. Ventrals 212—216, broad. Anal entire. Subcaudals 58 to 60 pairs.

Color whitish, tinged with red on the back, with a dorsal series of large spots of dark brown or black—becoming bands posteriorly—thirty or more

in number. Flanks with four series of small spots, alternating, more or less irregular faded and broken; the lower series on the edge of the abdomen. Posteriorly the lateral spots unite into a short vertical band, and upon the tail this unites with the dorsal spot, forming a transverse band. Head-shields spotted with brown. A band across the posterior extremity of the prefrontals through the eye, across the cheek above the angle of the mouth, a vertical wedge-shaped bar below the eye, and the margins of the labials black. On old specimens these bands are often obsolete, as is also the case to some extent with spots of the back and flank. The scales generally have a darker line along the middle. A specimen from Florida has three series of alternating quadrate brown spots on the belly, back nearly uniform brown posteriorly, clouded and spotted anteriorly, and the dorsal spots rather indistinct.

PITYOPHIS CATENIFER.

Loreal one, sometimes two. Preocular one, often two, lower small. Postoculars two to five, commonly three. Rows of scales 31 to 33. Ventrals 222 to 243. Subcaudals 58 to 72 pairs.

Spots 90 to 98 (19 to 23 on the tail). The band across the head through the eyes is distinct to obsolete. The vertical bar under the eye is usually distinct. Dorsal and upper lateral series of spots more or less confluent so as to form a chain on the back; in cases where the spots are separate the effect is much the same. The colors vary much in intensity; the spots are sometimes black, and occasionally quite indistinct. The ground color varies from white to reddish brown. Ventrals 230—247. Subcaudals 60—71. Oregon to Lower California.

var. SAYI.

Closely resembling *P. melanoleucus*. Spots rather more numerous, though not so many as in *P. bellona*. Sometimes two loreals. Preoculars 1 (1—2). Postoculars 3 (2—4). Scales in 27—29 rows. Ventrals 220—240. Subcaudals 52—55 pairs.

Belly more or less blotched in the middle. Mississippi Valley to Rocky Mountains, and southward to Mexico.

var. MEXICANUS.

Prefrontals two pairs, sometimes soldered into a single pair. Preoculars two, lower small. Postoculars four. Labials nine, fifth in the

orbit. Scales in 33—35 rows. Ventrals 229—239. Subcaudals 57 to 65 pairs. Reaches a length of seven feet.

With the band across the frontal and behind the eye. More or less spotted. Southwestern part of Mexico. (From descr.)

VAR. BELLONA.

Ground color lighter than in *P. melanoleucus*. Body more slender; apparently not reaching the same dimensions. In large specimens the bar below the eye and that across the head to the angle of the mouth are conspicuous. Dorsal series of spots from 70 to 80. Some specimens have the colors very much faded, others have the spots very black and distinct, particularly on the tail. The young have four alternating series of spots on each flank, the lower on the outer edge of the abdomen. Scales in 31 to 35 series. Ventrals 220—231. Subcaudals 43—63. I have taken this species at great distances from water on the open prairies. Rocky Mountain region.

ELAPHIS.

ELAPHIS *Aldrovandus*, 1640 and 1765; *Bonaparte*, 1831 and 1840;

Dunéril & Bibron, 1852 and 1854; *Günther*, 1858.

ELAPHE *Fitzinger*, 1833 and 1843.

Of moderate slenderness, slightly compressed, belly flat; head distinct from the neck, crown flat, muzzle rounded; tail moderate, tapering. Eye medium, pupil round. Teeth equal, smooth. A loreal. Nasal in two parts, nostril between. Orbitals 2—2 to 3. Scales keeled, in 19 to 25 rows. Anal bifid or entire. Subcaudals in two rows. In particular cases the carination of the scales becomes obsolete.

(*Scotophilis*.)

Anal entire;

Light greyish-brown (reddish in life), with dorsal blotches of brown or black, varying to uniform black *obsoletus*.

Pale red, with darker black-bordered blotches on the back, varying to black, with a few white markings under the chin and neck *guttatus*.

Anal bifid;

with four longitudinal brownish bands *quadrivittatus*.

ELAPHIS OBSOLETUS, *pl. IV., fig. 2.*COLUBER OBSOLETUS *Say, 1823, James' Acct. of Louj's Exp. I, 140.*

Long, slender, tapering to head and tail, compressed; belly flat, angled at the edges; neck small; head large, broad behind, narrow in front of the eyes, depressed, with flat crown; tail short, near one fifth of the total, slender. Eye moderately large, over the fourth and fifth labials, pupil round. Mouth-cleft deep, curved. Head-shields nine, broad. Frontal broad anteriorly. Prefrontals bent down on the side of the face. Snout prominent. Rostral large, a little broader than high, convex. Nasal in two parts, nostril between. Loreal low. One large anteorbital. Post-orbitals two. Temporals 2+3. Labials eight, seventh largest. Infralabials twelve, sixth largest. Submentals two pairs, the hinder separated by small scales. Scales lozenge-shaped, extremity rounded, in 27 rows, two or more of the outer smooth, medial keeled, those on the flanks more faintly. Ventrals about 233. Anal entire. Subcaudals about 85 pairs.

Light greyish-brown, reddish in life, varying to black. In light colored specimens of small size there is a dorsal series of about forty large spots, with darker edges, separated by narrow spaces of light color. On the flank three series of alternating spots, the spots in each half as large as the next above. On the extremities of the abdominals a series of twice as many quadrate spots. On each side of the median line of the belly a series of quadrangular blotches. In large specimens only the upper and lower of the series on the flank can be distinguished. A dark-edged band across the postfrontals, through the eye, over the angle of the mouth. A pair of diverging bands from the parietals to the neck. Anterior spots more or less confluent. Darkening with age until nearly or quite black. East of the Rocky Mountains.

VAR. ALLEGHANIENSIS.

Dark brown to lustrous black, mottled with white under the throat. Series of scales 27. Alleghany region and eastward.

VAR. LINDHEIMERII.

Dark plumbeous, with brown or black blotches on back and flanks. Series of scales 29. Texas.

Through its varieties this species is so closely related to the following that the separation is a matter of considerable difficulty. It is quite

likely that a larger series of specimens may render the degradation of *E. obsoletus* to the rank of a variety of *E. guttatus* unavoidable.

ELAPHIS GUTTATUS, *pl. IV, fig. 1.*

COLUBER GUTTATUS Linné, 1766, *Syst. Nat. ed. XII*, 385.

ELAPHIS GUTTATUS Dum. & Bibr., 1854, *Erp. Gén. VII*, 273.

Elongate, compressed, belly flat, angled at the sides; head elongate, distinct from the neck, tapering regularly to the moderately broad snout, crown flat; tail short, stout, nearly one sixth of the total, tapering. Eye moderate, over the fourth and fifth labials, pupil round. Mouth-cleft deep, curved. Head-shields nine, broad. Parietals broad. Frontal broad anteriorly. Prefrontals bent down on the side of the face. Internasals small. Rostral large, convex. Nasal in two parts, nostril between. A loreal. One large anteorbital. Two postorbitals. Temporals 2+3. Labials eight, seventh largest. Infralabials eleven (10—12), second small, sixth largest. Submentals two pairs, hinder separated by small scales. Scales lozenge-shaped, fore extremity rounded, in 27 rows (with the varieties 25—29), dorsal faintly keeled, keels obsolete on the lateral, two or more of the outer slightly broader and smooth. Ventrals broad, 217 to 236. Anal bifid. Subcaudals 63 pairs.

Pale red in life (brownish-yellow in alcohol), with about forty dorsal blotches of darker red, with a narrow border of black, anteriorly longer than broad, posteriorly broader than long, reaching to a point near the middle of the flank, separated by spaces of two or three scales in width. On the flank several (3 to 4) alternating very irregular and indistinct series of spots, the lower reaching the keel of the edge of the abdomen, and having twice as many spots as the others. Below yellowish with quadrangular blotches of black, alternating on each side of the median line or confluent. Head reddish, with a darker black-edged band across the hinder portion of the postfrontals through the eye across the angle of the mouth; a similar diverging band across the parietals to the neck, inclosing a light-colored space on the top of the neck and back of head. Vertical margins of labials and infralabials more or less black. The pattern of markings is usually confused, except in young specimens.

VAR. VULPINUS.

Light brown, more or less red or gray, with broad quadrate blotches of chocolate or brown. Series of scales 25. Michigan and adjacent States.

ELAPHIS QUADRIVITTATUS.

COLUBER QUADRIVITTATUS *Holbrook*, 1842, *N. A. Herp.* III, 89.

ELAPHIS QUADRIVITTATUS *Dun. & Bibr.*, 1854, *Erp. Gén.* VII, 265.

Elongate, compressed, angled at the sides; neck small; head distinct elongate, tapering regularly to the moderately broad snout; crown flat; tail short, near one fifth of the total, tapering. Eye moderate, over the fourth and fifth labials, pupil round. Mouth-cleft deep, curved. Head-shields nine, broad. Frontal broader anteriorly. Prefrontals bent down on the face. Internasals much smaller than prefrontals. Rostral large, broad. Nasal in two parts, nostril between. A loreal. Anteorbital large. Two postorbitals. Temporals 2+3. Labials eight, seventh largest. Infralabials eleven, second small, sixth largest. Submentals two pairs, hinder smaller and separated by small scales. Scales lozenge-shaped. Extremity rounded, in 27 rows, dorsal faintly keeled, lateral with faint keels, two or more of the outer broader and smooth. Ventrals 238. Anal bifid. Subcaudals 95.

Light yellowish-red (tinged with green anteriorly), with four longitudinal bands of brownish-red. The bands occupy the third to the fifth and the tenth to the twelfth rows on each side. Belly mottled with brownish-red. These colors occupy only the epiderm; that taken off, the markings appear as in the species *guttatus*. The head is uniform yellowish-red, with a faint tinge of green, but on close observation the outlines of the head-bands can be readily traced. A specimen before me, secured by Prof. J. A. Allen in Florida, has the colors of *quadrivittatus*, bands and all, and at the same time the black-margined spots and shorter tail of *C. guttatus*.

DROMICUS.

Bibron, 1843.

Body somewhat slender, rounded; head distinct from the neck, long, ovoid, crown flat; tail tapering. Eye medium, pupil round. Posterior maxillary teeth longer, smooth. A loreal, in one species united with prefrontal. Crown-shields nine. Nasal in two parts. Orbitals 1—2. Scales

smooth, commonly in 17 or 19 rows, rarely reaching 15 or 23. Common in the West Indies and South America to Mexico. One species found on the southeastern coast of the United States.

A loreal;

scales in 17 rows; labials seven;

ventrals about 163; a band on three vertebral series

laureatus.

ventrals about 126; dorsal band on eleven series

flavilatus.

scales in 17 to 19 series;

labials nine

margaritiferus.

No loreal;

scales in 19 series;

labials eight

putnami.

DROMICUS LAUREATUS.

Günther, 1868, *Ann. & Mag. Nat. Hist.*, 419.

Body and tail moderately slender; form of the head as in *Coronella laevis*. Eye of moderate size. Maxillary teeth smooth, becoming longer, stronger and further apart backward. Internasals scarcely half as large as prefrontals. Frontal longer than the snout. Parietals longer than the frontal. Rostral not extending to the upper surface of the head. Loreal square. One preocular reaching to the upper surface of the head, but not touching the frontal. Postoculars two. Labials seven, third and fourth in orbit. Temporals 1+2+3, the anterior touching both oculars. Submentals two pairs, nearly equal. Scales smooth, in 17 rows, many with a small apical groove. Ventrals 163. Anal bifid. Subcaudals 95 pairs.

A lead-colored band three scales broad from the nape along the median line of the back to the end of the tail. Flanks reddish, with a very indistinct greyish streak along the fourth outer series of scales. Two yellow lines across the rostral; the lower runs along the upper labials and across the neck, thus entirely encircling the head; the upper runs along the canthus rostralis, and stops or terminates in the temporal region. Lower parts uniform yellowish. Length 21 inches; tail 7 inches. City of Mexico.

DROMICUS FLAVILATUS.

Cope, 1860, Pr. Ac. N. Sc., Phil., 222.

Habit moderately slender; head distinct from the neck, elongate, oval; tail 3.2 times in the total. Crown-shields nine. Internasals nearly quadrate. Prefrontals longer than wide. Frontal three fourths as long as wide. Labials seven, third and fourth in the orbit, fifth higher than long, with the sixth separated by a narrow temporal from the parietal, seventh longer than high. Infralabials nine, four bounding the submentals. Nostril in the anterior portion of the divided nasal. Loreal very small, high as long. Oculars 1—2. Scales thin, without pores, in 17 series. Ventrals 126. Subcaudals 77 pairs.

A rich golden brown above, the scales of the two inferior rows on each side broadly gold-edged, the color of the back commencing on the third row. Ends of the scales of the vertebral row sometimes darker tipped. Head dark brown, darkest behind, with numerous but obscure paler vermiculations. Side of head paler, with a reddish-brown band from the rostral plate through the eye to the middle of the last labial. Labials whitish, with black dots on the posterior in oblique rows. Below white, lower labials sparsely black dotted. A pair of pale dots on the common occipital suture. Fort Macon, N. C.

DROMICUS MARGARITIFERUS.

HERPETODRYAS MARGARITIFERUS Schlegel, 1837, Ess. Phys. Serp., 184 pt. II; and Abbild. t. 44, fig. 19—20.

DROMICUS MARGARITIFERUS Günther, 1858, Cat. Col. Snakes, 126.

Elongate, moderately slender; head distinct from the neck, long, crown flattened, muzzle prominent; tail slender, near one half of the total length. Eye rather large, pupil round. Mouth deeply cleft, curved. Upper post-maxillary teeth large, isolated, smooth. Crown-shields nine. Prefrontals bent down to the loreal. Nasal divided. Loreal longer than high. One anteocular. Postoculars two, followed by two temporals. Labials nine, fourth, fifth, and sixth in the orbit, the three preceding the last large. Infralabials ten, sixth large. Submentals two pairs, posterior larger. Scales lozenge-shaped, in 17—19 rows, feebly keeled, outer two or three broader and smooth. Ventrals 154 (Schlegel), 115 (Duméril). Anal bifid.

Greenish-yellow on flanks and below. All the scales bordered with brown. The medial rows are described as black, with a whitish or yellowish center. Top of head yellowish-brown. Mexico and Central America.

DROMICUS PUTNAMI.

Jan., 1863, *Eleuc. Sist. Opid.* 67; 1867, *Icon. Ophid. Livr.*, 24, pl. 6, fig. 3.

Moderately slender, belly broad; head distinct from neck, depressed, narrow anteriorly; tail near one fourth of the total, tapering, pointed. Eye moderate, pupil round. Mouth-cleft deep, slightly curved. Crown-shields nine. Rostral erect, broader than high. Internasals moderately large. Prefrontals bent down on the side of the face. No loreal. Frontal narrow, hexangular. Supraciliaries large. Parietals large. Nasal in two parts. Oculars 1—2. Temporals 1+2+3. Labials eight, third, fourth, and fifth in orbit. Infralabials ten, fifth and sixth large, third not in contact with the submentals. Submentals two pairs, nearly equal. Scales smooth, glossy, in 19 rows, dorsal narrow, lateral broader. Ventrals 171. Anal bifid. Subcaudals 81 pairs.

Light brownish, with a dorsal band of brown three and two half scales wide. Outer row of scales a shade darker than the flanks. Belly lighter. Dorsal band continuing forward on the head. A narrow light-edged brown line from the eye toward the angle of the mouth. Total length $21\frac{3}{4}$ inches; tail $5\frac{1}{2}$ inches. San Blas, Jalisco, Mexico.

CORONELLINAE.

Subcylindrical, moderate to stout; head more or less distinct from the neck, which is rather stout; tail tapering, thick or slender. Snout moderate. Teeth equal and smooth; or posterior larger, separated by an interspace, smooth or grooved. Eyes small to moderate, pupil round (except in *Mesotes* and possibly *Hypsiglena*). Scales smooth, or, in exceptional cases, keeled.

Posterior maxillary teeth longer, grooved;

scales in 17 to 21 series

scales in 15 series

Maxillary teeth compressed, posterior longer, smooth;

anal entire

Postmaxillary teeth longer, generally separated by an interspace;

anal usually bifid

Teeth equal, smooth;

anal bifid

rostral swollen, bent back on the snout

TACHYMENIS.

ERYTHROLAMPRUS.

OPHIBOLUS.

LIOPHIS.

DIADOPHIS.

RHINOCHEILUS.

Postmaxillary teeth larger, separated by an interspace;	
rostral trihedral, sharp-angled	HETERODON.
Postmaxillary teeth longer, smooth;	
rostral swollen, rounded	CEMOPHORA.
eye large, pupil round	XENODON.
eye small, pupil subelliptical	HYP SIGLENA.

TACHYMENIS.

Wiegmann, 1834.

Moderately stout to slender; belly broad, rounded; head distinct from the neck, narrow in front; tail moderate, tapering. Eye moderate, pupil round or subcircular. Posterior maxillary teeth longer, grooved; anterior equal or slightly longer than those immediately following. Loreal present or united with other shields. Crown-shields nine. Internasals somewhat narrow in front. Oculars 1—2. Scales smooth, in 17 to 21 rows. Anal bifid.

Labials seven; scales in 19 series;

longitudinal bands, continued on the head	<i>lineata.</i>
ending at the back of the head	<i>imperialis.</i>
vermilion, shading through orange to the golden beneath	
	<i>lateritia.</i>
a dark band with white borders on each side of the neck	
	<i>proterops.</i>

Labials eight; scales in 21 (19—21) rows;

a white band on each side of the anterior part of the trunk	
	<i>fissidens.</i>
brown, with an indistinct dorsal streak	
	<i>bipunctata.</i>

head and neck brown; behind this a yellowish neck band of about four scales in width

melanocephala.

Several of these species can hardly be regarded as firmly established. The descriptions are insufficient for comparison.

TACHYMENIS LINEATA.

TOMODON LINEATUS Duméril & Bibron, 1854, *Exp. Gén.* VII, 936, pl. 73.

Head distinct from the neck; tail near one fourth of the total. Posterior upper maxillary teeth longer, grooved. Crown-shields normal. Rostral

convex, bent backward. Nasal in two parts. Loreal trapezoidal. One anteorcular. Postoculars two, subequal. Labials seven (7—8), gradually increasing in height backward to the penultimate, third and fourth in orbit (3d to 5th). Infralabials eight. Submentals two pairs, subequal, posterior diverging. Scales smooth, in 19 rows. Ventrals 159—169. Anal divided. Subcaudals 62—70 pairs.

Gray above, white below. A band of brown from the forehead to the tail. A brown line from each nostril along the side of the head and middle of the flanks, black-edged and becoming black backward. Labials and infralabials marked with brown. Sometimes a brown or black line near the edge of the abdomen. The dark edges of the subcaudals form a line under the middle of the tail.

TACHYMENIS IMPERIALIS.

TAENIOPHIS IMPERIALIS Baird (*Conn.*), 1859, *U. S. and Mex. Boundary Survey Rept.*, 23, *pl. XIX, f. 1.*

Form tapering anteriorly and posteriorly; tail continuous with the body; head distinct from the neck; snout rounded. Rostral gibbous, twice as broad as high. Loreal as high as long. Internasals one third as large as prefrontals, larger than the loreal. Frontal elongate, narrow, pentagonal, scarcely wider anteriorly. Anteorcular one, large, wider above. Postoculars two, small, upper largest. Scales smooth, in 19 rows, central narrow and acute, outer broader.

Deep purplish black above, with two dorsal stripes of yellowish-brown from head to tip of tail, and separated by a narrow vertebral line of the ground color. Head black above, with two narrow yellow lines from the nostrils to the sides of the occiput, crossing the upper angle of the orbit. Labials and under part of head yellowish, minutely mottled with black. Ground color of the back extending on to the ends of the ventrals. Middle of the abdomen uniform light yellowish in the alcoholic specimen, said to be bright red in life. (From descr.) Brownsville, Texas.

TACHYMENIS LATERITIA.

CONIOPHANES LATERITIUS Cope, 1861, *Pr. Ac. N. Sc., Phil.*, 524.

Head broad, pointed. Muzzle prominent, acute. Loreal square. Oculars 1—2. Labials seven, eye over third and fourth, fifth very large. Infralabials ten. Scales in 19 rows. Frontal nearly as broad as long. Anal bifid. Total length 24 inches; tail 7 inches.

Bright vermilion, punctulate with brown, passing through orange to golden on the belly. Head and neck for ten scales backward black. Labials bordered and traversed by yellow lines. Parietals dotted with yellow. (From descr.) Guadalupe, Mexico.

TACHYMENIS PROTEROPS.

CONIOPHANES PROTEROPS Cope, 1860, *Pr. Ac. N. Sc., Phil.*, 249.

Small; head scarcely distinct from the neck, muzzle not elevated. Anterior plates of the head small. Loreal a little longer than high. One pre, two postoculars. Labials seven, third and fourth in orbit. Frontal elongate. Parietals long. Infralabials nine. Submentals two pairs, subequal. Ventrals 130. Anal bifid.

Light brown above, every scale densely punctulated with darker, especially near the margins. From the first to the fourth rows this is deeper, giving the sides a darker shade. The vertebral rows of scales from the occipitals to the end of the tail are also darker. Top of the head densely and obscurely vermiculated and punctulated. The dark shade on the fourth row of scales becomes a band anteriorly, and is bordered above and below with white on the neck. The lower white border is continued to the eye, and is bordered above on the labials with black. The upper white border is discontinued on the neck, but reappears as a spot three scales back of the occipitals. Inferior half of rostral, upper and lower labials, chin, throat, and belly, light brownish-yellow, densely punctulated with brown. Each labial with a darker spot in the center. Fewer punctulations on the urosteges. (From descr.) Jalapa, Mexico.

TACHYMENIS FISSIDENS.

CORONELLA FISSIDENS Günther, 1858, *Cat. Col. Snakes*, 26.

Scales in 19—21 rows. Anal bifid. Labials eight, fourth and fifth in orbit.

Back greyish-olive, the darker coloration of the sides well defined towards the back. A black streak, sometimes white-edged beneath, through the eye. On both sides of the anterior part of the trunk a white band, beginning at the occiput, and soon disappearing. Belly white, on the sides some irregular small black spots. Hinder maxillary tooth longest, grooved. Mexico.

TACHYMENIS BIPUNCTATA.

CORONELLA BIPUNCTATA Günther, 1858, *Cat. Col. Snakes*, 36.

Moderate; tail rather elongate; head distinct from the neck, rounded, rather pointed in front, crown flat. Rostral just reaching the top of the head. Internasals small. Prefrontals much larger. Parietals moderate, rounded behind. Frontal nearly twice as long as broad, with somewhat concave lateral edges. Nasal in two parts, nostril between. A square loreal. One anteorcular. Postoculars two. Labials eight, fourth and fifth in orbit. One elongate temporal shield in front, four or five smaller ones behind. Scales rhomboid, rather small, in twenty-one rows. Anal bifid.

Brown, with a very indistinct darker dorsal streak. Darker coloration of the sides well defined towards the back. Labials yellowish above, black-edged. Beneath yellowish; from the chin to the tip of the tail two punctated lines, each ventral plate being provided with two black spots toward the middle; there is another indistinct punctated line on each edge of the abdomen. Total length 17 inches; tail $5\frac{1}{2}$ inches. (From Gthr.)

TACHYMENIS MELANOCEPHALA.

Peters, 1869, *Mb. Berl. Akad.*, 876.

Resembles somewhat *T. bipunctata* in squamation and infralabials. The entire head and neck for ten scales has a blackish-brown ground color, behind which there is a yellowish neck band of four scales in width, while the balance of the body is yellowish-brown above and uniform yellowish beneath. (From descr.) Puebla, Mexico.

ERYTHROLAMPRUS.

Boie, 1826.

Elongate, cylindrical; head not distinct from the neck, rounded, muzzle short; tail short to moderate. Crown-shields normal. Eye moderate, pupil round. Posterior upper maxillary teeth longer, grooved. Nasal in two parts, nostril between. Oculars 1—2. A loreal. Scales in 15 rows. Anal divided.

E. VENUSTISSIMUS. var. D.

Günther, 1858, *Cat. Col. Snakes*, 48.

Colors black, red, and yellow in rings around the body. Rings complete, but not arranged in pairs, broad, alternating with white rings of the same breadths. Muzzle black in front.

OPHIBOLUS.

Baird & Girard, 1853.

Body moderate to stout, rounded; head medium to small, little larger than the neck, muzzle short, rounded, crown flat. Eye medium, pupil round. Maxillary teeth compressed, posterior larger, smooth, not isolated. Nasal in two parts, nostril between, rarely but one nasal. Loreal generally present. One anteorcular. Two, sometimes three postoculars. Scales smooth, short, broad, in 17 to 23 rows. Anal entire. Subcaudals in two rows.

Loreal present or absent;

scales in 19 rows; red, with pairs of black rings separated by white or yellowish;

ventrals 169 to 176

doliatus.

Loreal present;

scales in 21 to 25 rows;

colors red, brown, or greyish, to very dark and yellowish;

ventrals 180 to 224

triangulus.

black or brown, with yellowish markings, to whitish or yellowish, marked with dark

getulus.

OPHIBOLUS DOLIATUS, *pl. V, fig. 2.*

COLUBER DOLIATUS Linn., 1766, Syst. I, 379.

Long, slender, subcylindrical; head little larger than the neck, crown slightly convex, narrow forward, snout prominent; tail short, about one seventh of the total, thick, tapering to a point. Eye small, over the third and fourth labials, pupil round. Mouth-cleft medium, curved. Crown-shields nine, broad. Frontal broad, short. Supraoculars and internasals small. Prefrontals bent down on the side, frequently united with the loreal. Nasal in two parts, nostril between. Loreal present or absent, sometimes present on one side and absent on the other as in the specimen figured. Oculars 1—2, lower small. Temporals 1+2. Labials seven, hinder three large and wedged by the temporals. Infralabials eight, fifth large, second and posterior small. Submentals two pairs, hinder smaller. Scales smooth, subhexangular, rounded at the extremity in 19 (17—19) rows, broadening toward the outer, which is as broad as long. Ventrals large, 169—176. Anal entire. Subcaudals 31—43 pairs.

Scarlet in life, reddish-yellow in alcohol, with 20—25 pairs of narrow black rings, those of each pair separated by a yellow band. The space between the pairs is nearly equal to that occupied by the pair and the included yellow. The black rings diverge and become narrow toward the belly, which in cases they do not cross. Occasionally the rings of different pairs unite on the abdomen. The first pair is incomplete, and does not encircle the neck, the anterior ring crosses the hinder extremity of the parietals, and commonly sends forward on the frontal an oblong or square extension, it stops at the angle of the mouth. The yellow rings widen on the ventral surface. Southeastern States.

VAR. ELAPSOIDEUS.

Loreal shield absent. Southeastern States.

OPHIBOLUS TRIANGULUS, *pl. V, fig. 1.*

COLUBER TRIANGULUM *Boie*, 1827, *Isis*, 537.

Moderately stout, subcylindrical; head small, little broader than the neck, depressed, crown slightly convex, snout comparatively broad; tail short, one sixth to one seventh of the total. Eye small, over the third and fourth labials, pupil round. Mouth-cleft deep, not much curved. Crown-shields nine, broad. Frontal broader anteriorly. Internasals small. Pre-frontals bent down on the sides of the head. Rostral broader than high, convex. Nasal in two parts, nostril between. Loreal low. Oculars 1—2, lower small. Temporals 2+2 to 3. Labials seven, posterior three larger. Infralabials nine (9—10), fifth large. Submentals two pairs. Scales smooth, lozenge-shaped, ends rounded, in 21 rows, outer as broad as long. Ventrals 200—215 (185—215). Anal entire. Subcaudals 45—55 pairs.

Brownish-yellow (more or less red in life), greyish on the flanks, with broad black-edged blotches of reddish-brown or brownish-red. The yellow separating the dorsal series is narrower than the blotches, and widens toward the flank, where in most varieties it bifurcates, and with the opposite band surrounds a medium-sized, light-centered blotch. Occasionally there are two series of alternating blotches on the flank. Belly yellowish, with quadrate black blotches, often more black than yellow. Back of

head with a triangular spot of yellow, bilobed in front. A band of brown across the hinder part of prefrontals through the eye to the angle of the mouth. East of the Mississippi Valley.

var. CALLIGASTER.

Colors somewhat lighter than in specimens from the Eastern States. Scales in 25 rows. Northern portion of Mississippi Valley.

var. MEXICANUS.

Body as in specimens from the Eastern States; head much swollen at the temples. Scales in 23 rows. Ventrals 193—199. Anal entire. Subcaudals 56—58 pairs.

Brownish-grey to greyish-brown, with 38—40 transverse blotches of red, inclosed by black, light-edged rings. These rings may be described as pairs of transverse bands, which unite on the flank just above a large light-edged black spot. The space occupied by the red and black together about equals the dark one separating the rings. Alternating with the spots on the flank there is a series of spots on each edge of the abdomen, which are more or less confluent with the blotches of the middle of the belly. Belly yellow, blotched with black, sometimes much more black than yellow, more yellow forward. Chin and throat yellow. A large black spot behind the eye, V-shaped marks of black on the top of the head, opening forward. First blotch behind the head irregular, emarginate arrow-shaped or divided longitudinally. Mexico, near San Luis Potosi.

var. DOLIATUS, *pl. V, fig. 2.*

Red, dorsal blotches more or less ring-like, 20 to 30, with or without the spot on the flank. The triangular spot on the back of the head is replaced by a transverse band, in front of which there is either a narrow black band or the whole top of the head is black, excepting the labials and rostral. Ventrals 181 to 208. Subject to great variation. Hab. southern part of Mississippi Valley.

var. GENTILIS.

Crown-shields, oculars, and rostral black. Prefrontals and loreal mottled with yellow, with twenty to forty yellow rings encircling the body. Between the yellow there are pairs of black rings inclosing a narrow space of red, across which the black is sometimes confluent on belly and back,

thus reducing the red color to vertical bars on the flank. (Ventrals 183 in a Utah specimen before me.) Hab. Colorado and Utah.

var. ZONATUS.

Fusiform, round; head small, scarcely distinct from the neck, depressed, rounded, crown slightly convex; tail short, stout to near the end. Eye small, pupil round. Muzzle narrow, rounded. Mouth-cleft nearly straight. Teeth equal, smooth. Head-shields short, broad. Rostral broad, low, convex. Internasals and prefrontals much broader than long. Frontal nearly as broad as long, posterior angle obtuse. Nasal in two parts, nostril between. Loreal very small. One anteocular, nearly as long as high. Postoculars two, lower small. Temporals 2+2. Labials seven, eye over third and fourth, fifth and sixth large. Infralabials eight, fifth large. Submentals two pairs, posterior half as large as the anterior. Scales short, broad, in 21 rows, outer broader than long. Ventrals 213, broad. Anal entire. Subcaudals 51 pairs.

Red. Each scale black-tipped. Head black to the temple. Surrounded by a yellow ring across the hinder portion of the parietals; behind this a complete black ring of equal width—about four scales—touches the angle of the mouth. Body with seventeen pairs of narrow black rings—14th on the vent—confluent on the belly, but including a yellow space of their own width three to four scales on the back, and separated from each other pair by a red space about as broad as that occupied by the pair and included space. Acapulco, Mexico.

O. GETULUS var. PYROMELANUS.

Head distinct from the neck, muzzle contracted; tail five and one half times in the total. Crown-shields nine. Frontal broad. Posterior submentals half as long as the anterior. Dorsal scales in 23 rows, rather broad, outer not abruptly enlarged. Ventrals 224. Anal one. Subcaudals 66 pairs.

Ground color ochraceous white. Fifty to fifty-eight black annuli, on the anterior portion of the body each is split by a vermilion annulus, posteriorly the division is incomplete, all extending with irregularities on the belly. In one specimen all the black annuli to the middle of the tail are divided by the red, leaving the black as a margin to it; they are four scales wide behind the middle of the body. In another only four anterior

rings are completely divided, those on the following third of the body being divided by red on the sides; the remaining annuli black, three scales wide; white annuli one and one half scales; anterior or nuchal red annulus widest, its anterior black margin attaining the parietals. An ochraceous band from the yular region, not quite completed, across the parietals. Muzzle, prefrontals, and labial margin ochraceous, remainder of top and sides of head black. Total length 30.5 inches. (From descr.) Sonora.

OPHIBOLUS GETULUS, *pl. V, fig. 3.*

COLUBER GETULUS *Linne*, 1766, *Syst. Nat. ed. XII*, 382.

OPHIBOLUS GETULUS *Baird & Girard*, 1853, *Cat. N. A. Serp.*, 85.

Stout, subcylindrical, slightly compressed, belly broad, rounded; neck large; head not very distinct, depressed, narrow in front; tail short, less than one sixth of total, tapering to a point. Eyes medium, over the third and fourth labials, pupil round. Mouth-cleft deep, outline curved. Crown-shields nine, broad. Frontal broader anteriorly. Rostral low, broader than high, convex. Nasal in two parts, nostril between. Loreal small. One large preocular. Postoculars two. Temporals 2+3. Labials seven, increasing in size to the sixth. Infralabials nine, fifth largest. Submentals two pairs. Scales rhomboid, smooth, in 21 rows (21—23), outer broad. Ventrals 210—225 (with the varieties 210—240). Anal entire. Subcaudals 48—55 pairs (48—65).

Lustrous brown to black, exceedingly variable. Back crossed by 25 to 35 narrow yellow bands, which bifurcate on the flanks to meet short longitudinal bars on the outer rows with which they form more or less complete unions to the bands immediately preceding or following, thus inclosing large spaces of black on the dorsal rows. Belly uniform black, black with yellow markings, yellow with quadrangular blotches of black, or nearly to quite uniform yellow. Head spotted with yellow, vertical margins of black. Some specimens are more yellow than black, others have very little yellow. Southeastern United States.

var. *SAYI*, *pl. V, fig. 4.*

Yellow bands more numerous, sometimes indistinct. All or nearly all of the scales yellow in the center. Rows of scales commonly 21. Mississippi Valley and Southwestward.

VAR. BOYLIJ.

Broad white transverse bands or spots on the sides. Dorsal rows usually 23. California to Mexico.

LIOPHIS.

Wagler, 1830.

Moderately slender; belly broad, rounded; head little larger than the neck, depressed; muzzle rather short; tail moderate, tapering. Eye moderate, pupil round. Crown-shields nine. A loreal. Oculars 1—2. Posterior maxillary teeth longer, smooth, generally separated by an interspace. Scales smooth in 17 to 19 rows.

LIOPHIS ELAPOIDES.

PLIOCERCUS ELAPOIDES Cope, 1860, *Pr. Ac. N. Sc., Phil.*, 253.

Body cylindrical; head scarcely distinct; tail two fifths of the total. Posterior upper maxillary teeth longer, smooth. Crown-shields normal. Rostral just visible from above. Length and breadth of the frontal equal to the length of the suture of the parietals. Two preoculars; upper large, not reaching the frontal; lower small, partially between the third and fourth labials. Postoculars two. Height and length of the loreal equal. Nasal divided. Labials eight, fourth and fifth in orbit. Infralabials eight, the last three times as long as the seventh, sixth largest; these three plates border within a large shield which diverges from the outer posterior extremity of the hinder submental. Submentals two pairs, equal. Scales smooth, in 17 rows. Ventrals 131 (to 143). Anal bifid. Subcaudals 89 pairs. Total length $19\frac{3}{4}$ inches; tail $7\frac{1}{2}$ inches.

Brilliant red bands from four to six scales in width encircle the body; these are separated by black rings, in sets of three, separated by yellow intervals, ten or eleven on the body—one at the anus and six or seven on the tail. The outer ring of each three is one and a half scales wide, and is not continued on the belly; the yellow interval is of the same width, and the central black is three and a half or four scales wide. The first three is on the head and neck; the central black is seven or eight scales wide, and does not extend upon the neck, but involves the ends of the parietals and the last upper labial. The anterior yellow ring crosses the parietals and involves one and a half temporals, the sixth, seventh, and half the eighth upper labials. All the head anterior to this is deep lus-

trous black, except a narrow oral border of yellow. Chin immaculaté. Many of the scales of the body are tipped with brown, many with black. (From descr.) Jalapa, Mexico.

DIADOPHIS.

Baird & Girard, 1853.

Small. Body subcylindrical, slightly depressed, slender; head distinct from the neck, depressed; tail short, tapering. Teeth equal, smooth. Crown-shields normal. Loreal present. Nasal divided. Eyes medium, pupil round. Mouth-cleft deep. Scales smooth, in 15—17 rows. Anal bifid. Subcaudals in two rows.

With pairs of transverse bands;	
scales in 17 rows	<i>annulatus.</i>
with a white band along each side of head and neck	<i>decoratus.</i>
with three longitudinal bands of brown	<i>fulvicittis.</i>
with an occipital band;	
scales in 15 rows;	
ventrals 148—160	<i>punctatus.</i>
ventrals about 190	<i>docilis.</i>
scales in 17 rows; occipital band rarely absent;	
ventrals more than 200	<i>regalis.</i>

DIADOPHIS ANNULATUS.

ENICOGNATHUS ANNULATUS Duméril & Bibron, 1854, Exp. Gén. VII, 335.

Head little larger than the neck, slightly convex, muzzle rounded; tail stout, long, near half as long as the body. Eye moderately large, pupil round. Crown-shields nine, broad. Angle of rostral between internasals nearly a right one. Rostral convex. Loreal small, lower behind than in front. Prefrontals decurved to the loreal. One large preocular. Postoculars two. Labials eight, fourth, fifth, and sixth in orbit, latter large. Scales smooth, lozenge-shaped, in 17 rows. Ventrals 142. Anal divided. Tail broken.

Brown, tinted with green. Head varied with black on the supraciliaries, temporal region, and in three bars beneath the orbit. Neck tinged with reddish, with pairs of transverse bands on the dorsal rows, followed by three series of vittae or spots of black on the median and third or fourth lateral rows. Flanks and ends of ventrals punctulate with brown. (From descr.) Coban (Haute Vera Paz).

DIADOPHIS DECORATUS.

CORONELLA DECORATA Günther, 1858, *Cat. Col. Snakes*, 55.

Body moderate, slightly depressed; head distinct, depressed; tail slender, near one third of the total. Eye moderate, pupil round. Crown-shields regular. Internasals and prefrontals broad. Frontal broad, acute-angled posteriorly. Parietals large, in length equal to that of the head in front of them. Nasal small, in two parts, nostril between. Loreal an irregular polygon, in contact with six plates. Anteorbitals three, lower very small, in notch between third and fourth labials. Postoculars two, lower smaller. Temporals 1+2, or fused with labial. Labials eight, fourth and fifth in orbit, seventh large and abruptly higher than the sixth. Infralabials nine, sixth large. Submentals two pairs, subequal, posterior diverging. Scales smooth, flat, rounded at the extremity, in 17 rows, outer broader. Ventrals 160. Anal bifid. Subcaudals 107 pairs, sometimes a few simple.

Greyish-brown, head darker, lighter toward the tail. A narrow white dark-edged band from the muzzle along the facial angle through the upper postocular slightly downward to the neck, then rising and occupying the fifth row of scales on the body, where it becomes brownish, and continues as a light line above a narrow line of dark on the fourth row. A narrow black line on the vertebral row. Labials and ventral surface whitish, the anterior labials and each extremity of the ventrals with a spot of dark. Length $9\frac{1}{8}$ inches; tail $3\frac{1}{8}$ inches. Mountains of Alvarez, Mexico.

DIADOPHIS FULVIVITTIS.

RHADINEA FULVIVITTIS Cope, 1875, *Batr. and Rept. Costa Rica*, 139.

Very likely to prove a variety of the preceding. "Head small, not very distinct from the body. Frontal a little longer than the suture from it to the nasals, and a little shorter than the common suture of the occipitals, two-thirds as wide as long. Rostral small, low; postnasal higher than long; loreal as high as long. Superior labials eight, seventh highest; temporals 1—1. Inferior labials ten, sixth largest, in contact with middle of post genecials. Scales poreless, in 17 rows. Gastrosteges 177. Anal divided; urosteges 91.

"Color above fulvous, below fulvous yellow. The three brown bands extend from the end of the nose to near the end of the tail; the lateral involves the fourth and half of each adjacent row of scales, and is black-

edged; the dorsal is three and two half scales wide, and is also black-edged. The brown is paler on top of the head, and the ground color is a narrow yellow band to the eye. Lips yellow, like the lower surfaces, unspotted." Orizaba, Vera Cruz.

DIADOPHIS PUNCTATUS, *pl. II, fig. 2.*

COLUBER PUNCTATUS *Linné*, 1766, *Syst. Nat.*, ed. XII, I, 376.

DIADOPHIS PUNCTATUS *Baird & Girard*, 1853, *Cat. N. A. Serp.*, 112.

Body slender, subcylindrical, slightly depressed; head distinct from the neck, depressed, crown flat; tail near one fifth of the total, tapering to a point. Crown-shields normal, broad, anterior short. Rostral broad, low, not reaching back on the head. Snout prominent, rounded. Frontal pentagonal. Nasal divided, nostril mainly in the anterior portion. Loreal nearly square. Oculars 2—2. Temporals 1+1. Labials seven, sixth largest, third and fourth in orbit, or labials eight, seventh larger, fourth and fifth in orbit. Infralabials eight, fifth very large. Posterior submentals half as large as the anterior. Eye moderate, pupil round. Mouth-cleft deep, more than half behind the eye. Scales smooth, in 15 rows, dorsal narrow, outer as broad as long. Ventrals 148—160. Anal bifid. Subcaudals 36—56 pairs.

Uniform bronzed, ashy or greenish brown above, with a narrow yellowish, dark-edged band across the occiput (sometimes obsolete). Hinder edge of the extremities of the ventrals with a black spot, giving the appearance of a zigzag line. Belly yellow or orange, purplish or reddish, uniform or with black spots, irregularly scattered or in a central longitudinal series. Chin yellow to upper margins of the upper labials, which, with that of the rostral, are black. New England to Mississippi.

VAR. ARNYI.

DIADOPHIS ARNYI *Kennicott*, 1859, *Pr. Ac. N. Sc., Phil.*, 99.

Abdomen more spotted with black, ventrals 160. Subcaudals 50 pairs. Kansas.

VAR. DOCILIS.

Baird & Girard, 1853, *Cat. N. A. Serp.*, 114.

Body above uniform ash-gray; yellowish white beneath, spotted with black. A proportionally broad yellowish white occipital ring, margined

with a narrow black line. Dorsal scales in 15 rows. Ventrals 193. Anal entire. Subcaudals 57 pairs. (From descr.) Rio San Pedro of Rio Grande.

DIADOPHIS REGALIS.

Baird & Girard, 1853, Cat. N. A. Serp., 115.

"Body above uniform greenish-ash; beneath light yellow, scattered all over with small black spots. No occipital ring. Dorsal scales in 17 rows. Ventrals 237+1+58 pairs." Sonora, Mexico.

A specimen of this species from San Luis Potosi has 210 ventrals. Anal bifid, and 73 pairs of subcaudals. The lower surface of the tail and hinder abdomen are a bright orange red, shading to yellow forward. Belly spotted irregularly with black. Labials and chin-shields with black spots on their hinder margins. Scales in the outer row dark with yellow bases, medials dark, finely sprinkled with ashy.

The three preceding species of *Diadophis* are very closely related. I am inclined to consider them as one, which, through several varieties, gradually increases in length, number of ventral plates and brilliancy of coloration from New England to Mexico.

RHINOCHEILUS.

Baird & Girard, 1853.

Body elongate, tapering little to head and tail, slightly compressed; head small, rather indistinct, narrow, crown convex, snout produced, pointed; tail short, tapering regularly. Maxillary teeth equal, smooth, without interspaces between anterior and posterior. Crown-shields nine. Rostral prominent, bent back on the top of the snout. Nasal in two parts, nostril between. Loreal present. Oculars 1—2. Scales smooth, short, broad, in 23 rows. Anal entire. Subcaudals entire or divided (the specimen before me has nineteen of the posterior divided).

RHINOCHEILUS LECONTEI.

Baird & Girard, 1853, Cat. N. A. Serp., 120.

Long, slender, subcylindrical, belly flat; head rather indistinct from the neck, little larger, crown convex, narrow and pointed in front; tail short, near one eighth of the total, stout, tapering. Posterior upper maxillary teeth stouter. Mouth-cleft medium, slightly curved. Eye moderate, pupil subcircular (apparently a little oblong, erect). Rostral prominent,

vertically compressed, so as to appear pointed in a side view, crowding backward between the oblique internasals. Nasal in two parts, nostril between. A loreal. Oculars 1—2. Temporals 2+3. Labials eight, fourth and fifth in orbit, seventh largest. Infralabials nine, fifth largest. Submentals one pair, followed by four small shields. Scales lozenge-shaped, smooth, flat, in 23 rows, outer broader. Ventrals broad, 191—206. Anal entire. Subcaudals 40—49 (on one specimen half of them are divided).

Black, red and yellow in life. In alcoholic specimens uniform yellowish white, with about thirty-three transverse black bands, separated by narrower spaces—26 to the vent. Each black band is composed of a narrow, elongate, subquadrangular spot on the middle of the back—about three scales long by five in width—and a wedge-shaped extension of black scales, with yellow spots, down each flank to the edge of the abdomen. On the outer two rows and the extremity of the ventrals on each side there is a series of irregular black spots, twice as many as there are of the bands. The spaces between the black bands were red. Each scale has a black spot in the center. Head black from frontal to neck. On each side of this the scales are black, with light centers. Upper labials margined with black. Muzzle yellowish.

The pattern of coloration is about as below. A dorsal series of elongate black spots—about 26 to the vent—separated by spaces equal or smaller, a series of twice as many small black spots on the flank—placed opposite the extremities of the former—and a third series on the edge of flank and abdomen alternating with the series above it. The increase of the amount of black on the scales tends to form bands and obliterate the original pattern. Southern California to Mexico.

VAR. TESSELLATUS.

Labials eight. Infralabials ten. Ventrals 178. Subcaudals 37 entire, plus 14 pairs. Coahuila, Mexico.

HETERODON.

Beauvois, 1799; Latreille, 1800, Hist. Rep., IV, 32.

Of moderate size. Body stout, belly flat. Head short, broad, little larger than the neck. Snout very prominent, shovel-shaped. Rostral a produced trihedral. Posterior maxillary teeth larger, separated from the

others by an interspace. Head, neck, and body very distensible. Tail short, pointed. Scales keeled or smooth. Subcaudals bifid. United States to the Argentine Republic.

Azygos shield in contact with the prefrontals	<i>platyrhinus</i> .
Azygos surrounded by small plates	<i>simus</i> .

HETERODON PLATYRHINUS, *pl. VI., fig. 5.*

Latreille, 1802, Hist. Nat. Rept., IV, 32, pl. XXVIII, figs. 1-3.

Stout, tapering backward, body, neck and head very distensible; head broad, short, slightly convex on the crown; tail short, near one sixth of the total, tapering. Eye large, surrounded on sides and below by small scales, pupil round. Mouth-cleft moderate, forming a single curve with the anterior face of the rostral. Posterior maxillary teeth larger, smooth, separated by an interspace from the others. Crown-shields broad. Supraciliaries large. An azygos shield behind the rostral, in contact with the internasals. Internasals subtriangular, separated by the rostral and azygos. Rostral very prominent, curving upward into an acute-edged trihedral or shovel, with a sharp ridge between the internasals. Nasal in two parts, the valvular nostril between. Loricals two, lower larger. Oculars ten (9-14). Temporals varying, three to four. Labials eight, sixth and seventh large. Infralabials eleven, third and fourth large. Submentals one pair, followed by small shields, in contact with three labials. Scales keeled, broad at the extremity, in twenty-five rows (23-25), dorsal narrow, caudal and outer row broad, first row smooth. Ventrals 120-150. Anal bifid. Subcaudals 45-60 pairs.

Reddish or yellowish-brown to black (*var. niger*). A series of twenty to thirty dorsal spots, usually surrounded by a margin of very light color. Tail with about nine transverse bands. On the upper portion of the flank there is a series of medium-sized spots alternating with the dorsal; below this there are three or four alternating series of small more or less indistinct spots, the lower encroaching upon the ventral scales. Yellow under the chin and throat, darkening posteriorly and becoming olivaceous mottled or blotched with brown. A dark light-edged band across the prefrontals through the eye to the angle of the mouth. A similar band on the frontal and supraciliaries just behind the eye, more or less joined with the broad band from each parietal to the sides of the neck. Immediately behind the

head, between the bands from the occiput, which spread around it, there is a median short longitudinal band of half their length. Mississippi Valley to the Atlantic.

VAR. NIGER.

SCYTALE NIGER *Daudin*, 1803, *Hist. Rept.*, V, 342.

HETERODON PLATYRHINOS, var. NIGER, *Jan.*, 1863, *Elenc. Sist.*

Dark brown or olivaceous to uniform black.

H. catesbyi of Dr. Günther is not separated from *H. nasicus* by the description. Southeastern States.

HETERODON SIMUS, *pl. VII., fig. 4.*

COLTHER SIMUS *Linné*, 1766, *Syst. Nat.*, ed. XII, 375.

HETERODON SIMUS *Hollbrook*, 1842, *N. A. Herp. IV*, 57, *pl. XV.*

Stout, tapering backward, belly flat, body, neck, and head very distensible. Head broad, short, slightly convex on the crown; tail short, near one sixth of the total, thick, tapering. Eye large, pupil round. Mouth-left moderate, forming with the face of the rostral a continuous curve. Posterior maxillary teeth isolated, larger, smooth. Crown-shields broad. Parietals, frontal, and prefrontals frequently cut into smaller plates. Supraciliaries broad. Azygos shield (behind the rostral) surrounded, and the internasals separated from the rostral, by small plates. Rostral broad, very prominent, produced obliquely upward and expanded into an acute-edged spade-shaped trihedral, sharp-ridged on the top of the snout. Nasal in two parts, sometimes three; nostril valvular. Loreals two, upper smaller. Oculars 10—13. Temporals 4+5, varying much. Labials eight, sixth and seventh large. Infralabials ten (9—11), fourth largest, four in contact with the submentals. Submentals one pair, short. Scales keeled in 25 (23—27) rows, medial narrow, lateral and caudal broad and rounded. The outer row is keeled in some, smooth in others, and in cases the keels are very indistinct on two or three rows above it.

Yellowish-brown to brownish-yellow. Pattern of coloration similar to that of *H. platyrhinus*, with a dorsal series (30—50) of large blotches; alternating with this on the upper part of the flank a series hardly half as large; below the latter, on the side, are three or four alternating series—more or less irregular and confluent—of small spots, the lower being upon the ends of the ventrals. Belly yellow, more or less clouded with brown,

to more black than yellow. Chin and throat yellowish. Tail generally lighter below. A brown band across the prefrontals through the eye to the angle of the mouth; a similar band behind the eye on the frontal and supraciliaries, separated from the former by a narrow band of light color. A broader band of brown from each parietal backward to the neck, and between them on the median line of head and neck a short longitudinal bar. Mississippi Valley to the Atlantic.

var. *NASICUS*, *pl. VI, fig. 6.*

HETERODON NASICUS Baird & Girard, 1853, *Cat. N. A. Serp.*, pp. 61, 157.

Rostral more produced and bent; prefrontals more dissected; head bands shorter and broader; belly with large quadrate blotches of dark color to nearly uniform black. Scales in 23 rows. Ventrals 135—150. Anal bifid. Subcaudals 34—50. Arkansas, Texas, New Mexico, California.

var. *KENNERLYI.*

HETERODON KENNERLYI Kennicott, 1860, *Pr. Ac. N. Sc., Phil.*, 336.

Head broad, short anteriorly.

Pattern of colors as in *simus*, spots smaller, more numerous, dorsal sometimes divided into two alternating series a portion of the length. Belly black, irregularly spotted with white. Oculars, loreal, internasals, azygos, prefrontals, and sometimes the frontal, more or less cut into small plates. In a specimen before me there are 21 plates in front of supraciliaries and frontal on the top of the head. Scales in 23 rows, outer broader, smooth, balance keeled. Ventrals 148. Anal bifid. Subcaudals 35 pairs. Mexico.

CEMOPHORA.

Cope, 1860.

Body elongate, subcylindrical; head rather indistinct; tail short, conical. Eye small, pupil round. Posterior maxillary teeth longer, smooth. Rostral swollen, not keeled. Nasal in two parts, nostril between. Loreal rarely absent. Scales smooth, in 17—19 rows. Anal entire. Subcaudals in two rows.

CEMOPHORA COCCINEA, *pl. VI, fig. 1.*COLUBER COCCINEUS *Blum., 1788, Licht. & Voigt., Mag., V.*

Rather small. Long, slender, subcylindrical; head little larger than the neck, subconical, snout turned upward, with prominent rostral, but not compressed into a sharp edge as in *Heterodon*. Tail short, near one seventh of the total, tapering to a point. Eyes small, pupil round. Mouth-cleft not very deep, somewhat curved. Posterior maxillary teeth larger. Crown-shields normal. Frontal broad, hexagonal. Supraciliaries very small. Prefrontals broader than long. Rostral swollen, extending back between the internasals. Nasal entire, sometimes grooved or half divided, occasionally divided. A small loreal. Oculars 1—2, lower smaller. Temporals 1+2. Labials six, second commonly entering the orbit, third beneath the eye and postorbitals. (A specimen shows fusion of the second and third on one side, and the second, third, and fourth on the other.) Infralabials eight, fourth largest. Submentals one pair. Scales smooth, in 19 rows; dorsal lozenge-shaped, pointed, outer little broader. Ventrals broad, 157—174. Anal entire. Subcaudals 34—45 pairs.

Crimson in life, yellowish in alcohol; white below. Crossed by pairs of transverse bands of dark brown or black, each pair inclosing a narrow band of yellow. Sixteen to nineteen yellow bands on the body, four to five on the tail; they are limited on the outer rows by a series of spots of dark, which alternate with the dark bands. Black rings about equal to yellow, two or three scales. The red space between the black rings has a width about equal to that occupied by the pair and the included yellow. The red spaces are more or less completely inclosed by the union of the black bands beneath them into a ring. The first black band crosses the head, behind the eye, to the angle of the mouth; in front of this the head is red, yellowish on the snout, behind it on the occiput the first yellow band is followed by the second black. Southern States east of the Mississippi.

VAR. COPEI.

The specimen described and figured by Prof. Jan under this name has the red blotches on the back much shorter and more numerous than is usual in the species. It had thirty-two black rings on body and tail. The loreal enters the orbit below the anteorbital. Tennessee.

XENODON.

Boie, 1827.

Stout, subcylindrical, belly flat; head distinct from the neck, short, broad, crown flat; tail short, stout, tapering. Posterior maxillary teeth larger, smooth. Eye large. Crown-shields nine. A loreal. Rostral moderate, reaching the top of the snout. Nasal divided. Scales smooth, in oblique series, much imbricate. Anal entire or divided. Subcaudals in two rows. Brazil to Mexico; East Indies.

XENODON BERTHOLDI.

Jan, 1863, *Arch. per la Zool. II, fasc. II*, 108.

Stout, belly broad; head large, broad, crown flattened; tail short, tapering. Eye moderate, pupil round. Posterior maxillary tooth long, smooth. Crown-shields nine, short, broad. Prefrontals bent down to the loreal. Frontal broad, concave or truncate in front. Parietals as broad as long. Nasal in two parts, nostril between. Loreal small. One anteorcular, short, high. Postoculars two or three, varying much in size and shape in different individuals. A large temporal in contact with the oculars. Labials eight, fourth and fifth in orbit, seventh largest. Infralabials eleven, sixth largest. Submentals two pairs, anterior larger. Scales smooth, lustrous, in 19 rows, lateral very oblique, much imbricate. Ventrals 149. Anal entire. Subcaudals 47 pairs (42—47).

Brownish, with twelve to fourteen dark light-edged transverse bands, which widen and bifurcate on the flanks, thus inclosing on the middle of the back a light space, somewhat of a lozenge-shape, which frequently has a darker patch in its center. Lower part of flank spotted with dark. Abdomen yellowish to olive, mottled with darker. This species resembles *X. rhabdocephalus* very much; the most prominent difference is in the entire anal shield. Closely allied to *X. colubrinus* Gthr. Mexico.

HYP SIGLENA.

Cope, 1860, *Pr. Ac. N. Sc., Phil.*, 246.

Cylindrical; head distinct from the neck, depressed, broad posteriorly, conic anteriorly; tail short, less than one fourth of the total. Upper posterior maxillary tooth longer, smooth, separated by an interspace. Pupil elliptic, erect. Crown-shields normal. Nasal divided. A loreal. Oculars 2—2. Scales smooth. Anal bifid. Subcaudals in two rows. Arizona; Lower California.

HYPSIGLENA OCHRORHYNCHA.

Cope, 1860, Pr. Ac. N. Sc., Phil., 246.

Elongate, subcylindrical, tapering little; head distinct from the neck, short, subconical, crown flattened, snout prominent; tail short, near one sixth of the total, tapering. Eyes small, pupil vertical. Post maxillary teeth longer, smooth, isolated. Nasal in two parts, nostril between. A loreal. Oculars 2—2. Temporals 1+2+3. Labials eight, fourth and fifth in orbit, sixth and seventh largest. Infralabials nine or ten, fifth or sixth largest. Submentals two pairs, posterior smaller. Scales smooth, in 21 rows, outer as broad as long. Ventrals 167. Anal bifid. Subcaudals 48 pairs.

Scales of body and head minutely punctulate with brown, giving a light brown or greyish appearance. With four series of alternating brown spots on each side, about 33 to vent. The two dorsal rows are the larger, and alternate or unite. The spots decrease in size down the flank; the lower are obsolescent. Belly light, immaculate. Head light; a brown band from the eye above the angle of the mouth to the neck, spreading posteriorly. Cape St. Lucas; Arizona.

CALAMARINAE.

Body cylindrical, stout to elongate; head seldom larger than the neck, comparatively short. Eye small to medium, pupil round. Usually some of the head-shields united. One anteorbital, sometimes united with the loreal. Not more than two postorbitals. Scales smooth or keeled, in 13 to 19 rows. Teeth equal and smooth, or the posterior maxillary larger and grooved or smooth.

- One preocular; loreal sometimes fused with other shields;
 nasal usually grooved; loreal rarely present; rostral produced;
 anal bifid; scales smooth, in 17 rows FICIMIA.
 nasal entire; scales smooth, in 19 rows;
 anal entire; loreal present CHEILORHINA.
 nasal divided; scales smooth, in 17 rows; no loreal;
 anal bifid STENORHINA.
 nasal divided; no loreal; scales smooth, in 15 rows;
 anal divided or entire TANTILLA.
 nasal entire; loreal rarely present; scales smooth, in 15 rows;
 anal entire or divided; post maxillary teeth larger, grooved
 ELAPOMORPHUS.
 nasal entire or divided; scales smooth or keeled, in 15 rows;
 anal bifid CONTIA.
 Loreal and preocular united;
 nasal divided; scales keeled;
 anal entire NINIA.
 nasal divided; prefrontal entering the orbit;
 anal bifid; scales smooth or keeled VIRGINIA.
 nasal entire; rostral produced; scales smooth;
 anal bifid CARPHOPHIS.
 nasal divided; scales smooth;
 anal entire GEOPHIS.

FICIMIA.

Gray.

Moderately stout, subcylindrical; head moderate, slightly depressed; tail short to medium. Teeth small, equal, smooth. Eyes small to medium, pupil round, subcircular in *variegata*. Head-shields seven to nine,

short, broad; prefrontals and internasals often united. Rostral large, prominent, produced backward between the internasals or prefrontals, convex, sometimes resembling that of *Heterodon*. Nasal usually entire and grooved behind the nostril, sometimes divided or without the groove. Oculars 1—2. Scales smooth, broad, in 17 rows. Anal divided. Subcaudals in two rows. (Anterior labials, rostral, and nasal sometimes united.)

Internasals and prefrontals not united;	
rostral separating internasals but slightly;	
prefrontal and labials in contact	<i>frontalis.</i>
rostral separating internasals, but not reaching frontal	<i>cana.</i>
Internasals and prefrontals united;	
nasal and anteorbital in contact	<i>nasus.</i>
rostral not reaching the frontal;	
loreal generally distinct	<i>maculata.</i>
rostral reaching the frontal;	
loreals fused with prefrontals;	
back with black cross bars	<i>variegata.</i>
back blackish-olive	<i>olivacea.</i>

FICIMIA FRONTALIS.

TOLUCA FRONTALIS Cope, 1864, *Pr. Ac. N. Sc., Phil.*, 167.

Muzzle prominent, pointed, slightly narrowed. Prefrontals slightly separated by the rostral. Nasal long. Frontals in contact with labials. Preocular low, narrow. Two postoculars, lower barely in contact with one temporal. Seven upper labials; eye over third and fourth; seven lower. Parietals longer than frontal. Anterior suture of frontal a little longer than straight lateral. Scales in 17 rows, equal, thin. One pair of short submentals. Ventrals 144. Anal bifid. Subcaudals 44 pairs. Total length 10 $\frac{3}{8}$ inches; tail, 1 $\frac{1}{2}$ inches.

Greyish-brown, more rufous medially, with about thirty-six rhombic dark-edged spots of brown, six scales wide and four long, the angles of which are produced as vertical lateral bars. Spots becoming similar to cross-bands posteriorly, separated by a pale spot on the vertebral line. A brown cross-band across prefrontals and frontals; a longitudinal band on each occipital and side of nape. (Cope.) From Colima, Mexico. A doubtful species.

FICIMIA CANA.

GYALOPION CANUM Cope, 1860, *Pr. Ac. N. Sc., Phil.*, 243.

"Form stout; tail one eighth of total length. Head slightly distinct, large, depressed. Rostral plate acute; its anterior border elevated; its upper surface concave. It is produced backwards, separating the prefrontals, not reaching the vertical. Frontals two pairs. Nasal confounded with the first labial, a groove from the nostril to the suture of the second labial. No loreal; its place supplied by the post frontal. One pre two post oculars. Scales smooth; anal and subcaudal scutellae divided. Teeth small, of equal lengths. Pupil round."

Internasals small, triangular. Anterior border of frontal not angulated. Labials seven, third and fourth in orbit. Infralabials seven, fourth largest. Submentals short. Scales broad, in 17 rows. Ventrals 130. Subcaudals 28 pairs. Total length $7\frac{1}{2}$ inches, tail 11 lines.

Back brownish-grey, with irregular transverse bands of brown (thirty-one in specimen described), which extend to the ventrals, and anteriorly exhibit a tendency to divide in three series of spots. Eight spots on tail. First spot on neck large. A brown band from one angle of the mouth to the other, another through the eye across the frontals. Dirty yellowish beneath, including the first row of scales. Arizona.

FICIMIA NASUS.

CONOPSIS NASUS Günther, 1858, *Cat. Col. Snakes*, p. 6.

Body cylindrical; head not distinct from neck, rounded, conical, sharply pointed in front; tail stout, tapering. Eye moderate, pupil round. Mouth moderate. Teeth equal, smooth. Upper jaw much the longer. Rostral a protruding four-sided pyramid, with rounded edges, resembling *Rhinostoma*, but without sharp ridge. Frontals and internasals fused, bent on the sides. Frontal very large, longer than broad, six-sided. Supraciliaries elongate. Nasal entire, in direct contact with the anteorbital. No loreal. One antecular, not raised on the crown. Postoculars two. Labials seven, third and fourth in orbit. Temporal one, large, elongate. Submentals two pairs, anterior larger. Scales smooth, short, rounded at the tip, in 17 rows. Ventrals 118. Anal bifid. Subcaudals 38 pairs.

Nearly uniform dark olive, anteriorly some obsolete black spots; sides lighter; belly yellowish-white, edges of each ventral plate with two or three black spots. Total length 10 inches; tail 2 inches. (Günther.) California.

FICIMIA MACULATA.

OXYRHINA MACULATA Jan, 1862, *Prodr. Icon. Gèn. Ophid.*, pt. II, 61.

Subcylindrical; head not distinct from neck, subconical, slightly depressed, muzzle produced. Snout long, rostral conical. Eye moderate, pupil round. Mouth-cleft deep, curved. Prefrontals and internasals united. Frontal broader forward, hexangular. Nasal entire. A loreal, sometimes absent. One anteorbital. Postorbitals 1 to 2. Temporals 1+2. Labials 6 to 7, the two preceding the last more or less wedged. Infralabials 6 to 7, fourth large, anterior four in contact with the submentals. Two pairs submentals, anterior larger. Scales smooth, short, broad, in 17 rows. Ventrals 125—131. Anal bifid. Subcaudals 28 to 36 pairs.

Brownish, more or less punctulate, with a dorsal series of irregular spots, the anterior of which are somewhat elongate. Flanks marked with brown, as if the dorsal spots had continued downward and broken up. Belly light, with spots of dark irregularly placed in a series on each side. (Jan.) Mexico.

FICIMIA VARIEGATA.

AMBLYMETOPON VARIEGATUM Günther, 1858, *Cat. Col. Snakes*, 7.

Moderate; head not distinct, rounded, muzzle not elongate; tail rather short, tapering. Eye moderate, pupil sub-elliptical. Rostral bent upward, with a convex ridge, not tapering behind, separating the prefrontals. Internasals and prefrontals united. Frontal large, seven-sided. Oculars 1—2. No loreal. Labials seven, third and fourth in orbit. Anterior temporal elongate. Scales smooth, nearly square, in 17 rows. Ventrals 151. Anal bifid. Subcaudals 37 pairs.

Back reddish-white, with 56 black narrow cross-bars, two on the neck larger than the remainder. Head with symmetrical black markings. Sides black spotted. Belly uniform whitish. (Günther.) Mexico.

FICIMIA OLIVACEA.

Gray, 1849, *Cat. Snakes Brit. Mus.*, 80.

Head moderate, rather depressed; body subcylindrical; tail tapering. Rostral large. Internasals and prefrontals fused, separated by the rostral. Loreal and prefrontals united. Supraciliary small, triangular. Frontal and parietals large. Nasal triangular, grooved behind the nostril. Eyes small, pupil round. Scales smooth, lustrous, broad, rounded at the tip.

Blackish-olive; lips, chin, and beneath pale yellowish. (Gray.) Mexico.

CHEILORHINA.

Shape and general appearance that of *Elapomorphus*, from which it is distinguished by the greater number of rows of scales, and by the smooth teeth, which gradually increase in size backward.

CHEILORHINA VILLARSII.

(*De Fil.*) *Jun*, 1862, *Prodr. Icon. Ophid.*, pp. 55 and 57; *Icon. Lirr.*, 48, pl. I, fig. 5.

Head not distinct from the neck; tail short, thick, blunt; snout prominent, broad, rounded. Internasals united with the prefrontals. Frontal hexangular, anterior angle produced. Supraciliaries short, broad. Nasals entire, united with the anterior labials. Loreal present. Preocular one. One postocular. One temporal, in contact with the ocular. Five labials, including that united with the nasal, third below eye and orbitals. Infralabials six. Submentals one pair. Scales broad, smooth, in 19 rows. Ventrals 213. Anal entire. Subcaudals 15 pairs.

Crown, labials, and infralabials black to the temporals. With transverse rings of black, more or less incomplete below, separated by spaces of about half their width. The first black ring crosses the neck, and occupies about seven scales. Two rings surround the tail; all grow narrower on the flank. Western Mexico.

STENORHINA.

Duméril & Bibron, 1854, *Erp. Gén. VII*, 865.

Moderate, subcylindrical, belly flattened; head little larger than the neck, depressed, narrow forward; tail short, stout, conical. Eye small to moderate, pupil round. Posterior maxillary teeth a little longer, grooved. Internasals and nasals usually fused. Nasals divided. No loreal. Scales smooth, in 17 rows. Generally one ante and two post oculars. Anal bifid. Subcaudals in two rows. Mexico to Central America.

Uniform brownish or slightly blotched;

bases of scales darker

with longitudinal bands of darker.

fremenvillei.

quinquelineatus.

STENORHINA FREMENVILLEI.

Duméril & Bibron, 1854, *Erp. Gén. VII*, 868.

Cylindrical, belly broad, tapering slightly to neck and tail; head moderate, little, if any, larger than the neck, depressed, crown flattened, slightly arched in front of the eye; tail short, stout, conical. Eye small,

pupil round. Internasal of each side united with the anterior portion of the nasal. Rostral broad, low, blunt-angled between the internasals. Prefrontals broad, in contact with the posterior portion of the second labials. Frontal hexangular, broader forward, hinder angle more acute. Parietals short, broad. No loreal. One anteorbital, beneath the supra-ciliary, not in contact with the nasal. Two post orbitals, lower small. Temporals 1+2. Labials seven, third and fourth in orbit, fourth to sixth large, and nearly equal in size. Infralabials seven, second smallest, fifth largest. Submentals two pairs, posterior small. Scales smooth, broad, rounded at apex, lateral as broad or broader than long. Ventrals 171—175 (165—175). Anal bifid. Subcaudals 35—40 pairs (35—43).

Light yellowish, to dark olivaceous, brown, base and sides of each scale darker. Belly yellowish to olivaceous, mottled with olive, bases of ventrals darker. Throat, chin, and lips lighter to white or yellow. Central America.

STENORHINA QUINQUELINEATA.

MICROPHIS QUINQUELINEATUS Hallowell, 1854, *Pr. Ac. N. Sc., Phil.*, 97.

“Head of moderate size, short, depressed in the middle, covered with nine plates; snout rounded; nostrils between the nasal and anterior frontal plates; a frenal; one ante ocular; two post oculars; seven superior labials, eye over third and fourth; eyes of moderate size, pupil round; supra-ciliary plate not projecting over the eye; tail of moderate length. . . . Head, body, and tail above ash color; five dark blue stripes extending from the occiput to the base of the tail, the three intermediate ones broader than the lateral; the other stripes prolonged upon the tail, the middle one to near its extremity; 17 rows of smooth quadrangular scales; 170 abdominal scuta; two bifid preabdom; 33 subcaudal.” Honduras.

TANTILLA.

Baird & Girard, 1853, *Cat. N. A. Sep.*, 131.

Body slender, subcylindrical; head indistinct, depressed; tail rather short, tapering. Eye small, pupil round. Nine head-shields. Nasal divided, nostril in anterior part, sometimes entire. No loreal. Orbitals 1—1 to 2. Scales smooth, lustrous, in 15 rows. Anal divided or entire. Subcaudals in two rows.

No band across the occiput; oculars 1—1; labials six with three longitudinal bands of darker color	<i>gracilis.</i> var. <i>calamarina.</i>
oculars 1—2; labials seven; head black	<i>nigriceps.</i>
an occipital band; oculars 1—2; labials seven	<i>coronata.</i>
a black spot on the occiput; oculars 1—1	<i>planiceps.</i>

TANTILLA GRACILIS, *pl. VI., fig. 3.*

Baird & Girard, 1853, Cat. N. A. Serp., 132.

Small, wormlike, slender, slightly depressed; head small, not distinct from neck, depressed, crown convex; tail short, slender, tapering to a point. Eye small, over third and fourth labials, pupil round. Mouth-cleft deep, nearly straight. Snout broad, very prominent. Head-shields nine, broad. Frontal hexagonal, or rounded in front. Rostral broad, subtriangular, convex. No loreal. Nasal in two parts, nostril in anterior. Orbitals 1—1. Temporals 1+1. Labials six, posterior two larger, the temporals wedging between them. Infralabials six, fourth larger, anterior pair slightly separated by the submentals in the specimen described. Submentals two pairs, posterior smaller. Scales smooth, in 15 rows, dorsal lozenge-shaped, outer and caudals somewhat broader. Ventrals 124—129. Anal bifid. Subcaudals 41—45 pairs.

Light reddish or olivaceous-brown, head darker. Beneath much lighter, yellowish. Posterior labials white; the white extending upward to the middle of the temporal. Individuals vary in regard to contact of mentals and submentals. Texas.

VAR. CALAMARINA.

Cope, 1866, Pr. Ac. N. Sc., Phil., 320.

Head flat, not distinct from neck; tail near one sixth of total length. Head-shields nine. Internasals narrow. Prefrontals in contact with second labial. Frontal longer than broad, angled in front. Supracilia-ries small. Nasal large. Orbitals 1—1, small. No loreal. Labials six, third and fourth in orbit, and supporting the orbitals. Infralabials seven,

fourth largest, the anterior widely separated from each other by contact of mental and submentals: Anterior submentals longer than broad, posterior minute. Total length $7\frac{1}{2}$ inches; tail $1\frac{1}{2}$ inches.

Brown, end of muzzle yellow, lower surfaces and occipital region pale. Sides and top of head and three longitudinal bands blackish; the latter extend on to the common line of the third and fourth, and on the vertebral series of scales. Guadalaxara, Mexico. (Cope.)

TANTILLA NIGRICEPS.

Kennicott, 1860, Pr. Ac. N. Sc., Phil., 328.

More slender than *T. gracilis*, head narrower. Frontal more elongate posteriorly, parietals narrower. Orbitals 1—2. Labials seven.

Uniform brownish white (in Ale.), light below. Crown as far as the parietals deep black; no indication of a post-occipital black ring as in *T. coronata*. (Kennicott.) Texas and New Mexico.

TANTILLA CORONATA.

Baird & Girard, 1853, Cat. N. Amer. Serp., 131.

Small, worm-like, slender, slightly depressed; head small, indistinct, depressed, crown convex; tail short, slender, tapering to a point. Head-shields nine, broad. Prefrontals short. Frontal hexangular, broad anteriorly. Parietals long, much separated in front by the frontal. Snout broad, prominent. Rostral broad, bent back on the snout. Nasal divided, nostril in anterior portion. No loreal. Orbitals 1—2. Eye small, over the third and fourth labials, pupil round. Labials seven, posterior largest. Infralabials six (6—7), fourth largest. Posterior pair of submentals much shorter. Temporals 1+1. Scales smooth, subrhomboidal, in 15 rows, outer broader. Ventrals 144. Anal bifid, rarely entire. Subcaudals 40 pairs.

Reddish-brown, head darker. A band of black from one half to six scales wide across the neck on the dorsal rows, sometimes absent. A narrow band of light color, a scale or two in width, crosses the occiput on the extremities of the parietals. Frequently a light spot on the fifth labial, reaching toward the eye. Infralabials more or less black. Belly and throat uniform whitish. South Carolina to Mississippi.

T. wagneri (Jan), from Florida; has an entire anal-shield.

TANTILLA PLANICEPS.

COLUBER PLANICEPS *Blainville*, 1835, *Nouv. Ann. du Muséum*, p. 294, pl. 27, f. 3—3b.

TANTILLA PLANICEPS *Cope*, 1875, *Check list*, p. 35.

Slender, elongate, cylindrical, head small, depressed, little larger than the neck; snout short; tail short, pointed. Nostril lateral, in the middle of the elongate nasal. Eyes medium. Mouth broad, deeply cleft. Head-shields nine. No loreal. Orbitals 1—1. Ventrals 134. Subcaudals 56 pairs. Scales broad, lustrous, smooth.

Uniform russet (light reddish-brown) above, whitish below, with a black spot on the occiput and commencement of the neck.

Duméril & Bibron say the nasals are divided by a suture below the nostril. *Erp. Gén.* VII, 858. (From descr.) California.

ELAPOMORPHUS.

(*Wiegmann*) *Fitzinger*, 1843, *Syst. Rept.*, 25.

Form resembling that of *Elaps*. Long, cylindrical; head rather indistinct from the neck; crown convex; tail short, thick, conical. Eyes small, pupil round. Muzzle broad, rounded. Posterior upper maxillary teeth generally somewhat larger and grooved. Nasal entire, rarely bifid. Loreal exceptionally present. Internasals two, or fused with prefrontals, or each other. Prefrontals two, sometimes united. Preocular one. One postocular, sometimes two. Scales smooth, in 15 rows. Anal entire or divided. Subcaudals in two rows.

ELAPOMORPHUS MEXICANUS.

Günther, 1862, *Ann. & Mag. Nat. Hist.*

Moderately slender; tail near one fourth of total. Internasals small, short, broad. Prefrontals short, broad. Frontal six-sided, anterior a little greater than a right-angle. Oculars 1—2, anterior not in contact with the frontal. Labials seven, third and fourth in orbit, seventh largest. Anterior temporal large. Submentals two pairs, anterior rather larger. Scales smooth, in 15 rows. Ventrals 158. Anal bifid. Subcaudals 52.

Brownish olive, with three blackish longitudinal bands, viz: one, almost linear, along the vertebral series of scales; the two others along the sides, each composed of two blackish, one line running along the middle of the second outer series, the other along that of the third. Neck with a pair

of small yellowish spots; front part of the snout crossed by a yellowish band; a black spot on the lips below the eye; uniform yellowish below. (From descr.) Mexico.

CONTIA.

Baird & Girard, 1853.

Moderately elongate, sub-depressed; head little larger than the neck, depressed; tail short, tapering to a point. Eyes small, pupil round. Crown-shields normal. Muzzle protruding. Rostral convex, broad. Nasal entire or divided. A loreal, sometimes united with other shields. One anteorbital. Generally two postorbitals, sometimes one or three. Temporals commonly 1+2. Labials seven (5—7). Infralabials seven (6—8). Scales smooth, rarely keeled, in 15 series. Anal bifid. Subcaudals in two rows.

(*Sonora.*)

Colors in transverse bands;

ventrals 149; bands 31

semiannulata.

ventrals 158; bands 41

occipitalis.

ventrals 158—167; bands 31

isozona.

scales keeled on the tail

aemula.

(*Contia.*)

Colors not in transverse bands;

oculars 1—2;

labials eight; nasal grooved;

scales slightly keeled at the tail

pygaea.

labials seven; flank with longitudinal band;

scales smooth;

entrals about 161, with black bases

mitis.

ventrals about 145

episcopus.

CONTIA SEMIANNULATA.

SONORA SEMIANNULATA Baird & Girard, 1853, Cat. N. A. Serp., 117.

Body subcylindrical, moderate; head not very distinct from neck, narrower on the snout, muzzle produced, elongated; tail short, about one

fifth of total, tapering. Eye moderate, pupil round. Head-shields nine. Frontal narrow anteriorly. Rostral large. Internasals one third smaller than prefrontals. Nasal divided. Loreal elongated, angular, horizontal. Orbitals 1—3. Posterior 2—3. Temporals 1+2. Labials seven, nearly equal, first and last smallest. Infralabials eight (7—8), fourth largest. Scales smooth, broad, in 15 rows, outer broader. Ventrals 149. Anal bifid. Subcaudals 39 pairs.

Body crossed with transverse jet black bars, 25 from head to anus, extending between and involving the exterior rows, becoming narrower on the flanks; along the back three to four scales long. Space between the bars above orange red, one scale wider than the black bars; on the sides greenish, with the base of the scales blackish. On the tail six black rings, continuing all around, covering two to eight scales; intermediate space red orange, four to five scales wide. Beneath uniform dull green, the black bars of the body not touching the scutellae. (From deser.) Sonora.

CONTIA OCCIPITALIS.

RHINOSTOMA OCCIPITALE Hallowell, 1854, *Pr. Ac. N. Sc., Phil.*, 95.

"Teeth of equal length, posterior ones not channeled; head depressed, eyes small; a rostral, two anterior and two posterior frontals; one anterior and two posterior oculars; a ventical; two occipitals; a small frenal; nostrils in a single plate; seven superior labials, the eye resting on the third and fourth. Abdominals 158; subcaudals 34.

"Milk white above, with 41 transverse black bands, including seven upon tail and one upon posterior part of head; six complete rings upon tail; throat and abdomen white, with transverse bands continuous with those upon back of tail; 15 rows of smooth quadrangular scales." Mohave Desert.

It is possible that this species and *C. isozona* will yet be united with *C. semiannulata*.

var. ANNULATA.

Completely encircled by black rings. Colorado Desert.

CONTIA ISOZONA.

Cope, 1866, *Proc. Ac. N. Sc., Phil.*, 304.

Tail four and two fifths times in the total length. Eye small, diameter twice in length of muzzle. Orbitals 1—2. Rostral rounded, slightly produced backward. Prefrontals and internasals much broader than long. Loreal twice as long as high. Labials seven, all higher than long. Eye over third and fourth. Scales in 15 rows, all broader than long. Ventrals 158—167. Anal divided. Subcaudals 52 pairs.

Twenty black half rings, separated by equal spaces of pinkish ground color. Belly immaculate. Tail completely six-annulate. The second specimen was vermilion between the bars and on top of the head. Probably a variety of *C. semiannulata*. (From descr.) Utah.

(Procinura.)

CONTIA AEMULA.

PROGINURA AEMULA Cope, 1879, *Pr. Ac. N. Sc., Phil.*, 262.

Post maxillary teeth longer, smooth, not isolated. Subcylindrical (elapomorphic); head flattened on crown, muzzle projecting, rounded; tail short, near one sixth of total, covered with roughly keeled scales. Head-shields nine. Rostral obtuse-angled on top of snout. Prefrontals wider than long. Frontal wide, sending a long angle backward. Parietals short, wide. Nasal in two parts, nostril between. Loreal quadrangular. Orbitals 1—2. Temporals 1+2. Labials seven, first longer than high, third and fourth in orbit. Infralabials eight, fourth largest. Submentals two pairs, posterior small. Scales broad, rounded, in 15 rows, outer broader, smooth on anterior portion of body, medial rows keeled a short distance in front of the vent, caudals all keeled. Ventrals 148. Subcaudals 41 pairs.

Colors similar to those of *Elaps fulvius*. Broad black rings surround the body; these are broadly bordered with yellow, and separated by red twice as wide as the yellow. Red scales with black centers. A black spot on the crown covers frontal, supraciliaries, and parietals. Total length 0^m, 364; tail 0^m, 061. (From descr.) Mexico.

(Contia.)

CONTIA PYGAEA, *pl. VII, fig. 5.**Cope, 1871, Proc. Ac. N. Sc., Phil., 223.*

Small, moderately stout; head little broader than the neck, subconical, slightly depressed; tail short, near one sixth of the total length, distinct, tapering to a point. Eye moderately large, pupil round. Head-shields nine. Rostral broad, low. Internasals narrow in front. Nostril small, round, nearer the upper edge of the nasal, which is grooved to the anterior labial. Lower angle of the loreal produced toward the eye. One anteorbital, narrow below. Postorbitals two. Labials eight, third, fourth, and fifth touching the eye. Infralabials 9. One temporal in contact with the orbitals, sometimes divided. Scales keeled, roof-shaped, or smooth, in 17 rows, apparently subhexangular, medial narrow, outer wider than long. Ventrals 120—133. Anal bifid. Subcaudals 35—54 pairs.

Back uniform brown; in some young specimens each scale bears a light line, which has the effect of strong keels. Bases of the ventrals dark toward the flanks, more uniform yellow medially. The dark color of the ventrals along the outer row forms a zigzag longitudinal line. Largest specimen $6\frac{1}{2}$ inches; tail one inch. North Carolina to Florida.

*CONTIA MITIS.**Baird & Girard, 1853, Cat. N. A. Serp., 110.*

Elongate, depressed; head depressed, little larger than the neck; snout short, broad, rounded in front; tail short, rather thick, tapering gradually to a short distance from the tip, whence more abruptly to the spine-like point. Eyes small, over the third and fourth labials. Head-shields nine, anterior short, broad. Rostral broader than high. Nasal divided, often partly or entirely united. Loreal present, sometimes fused with the prefrontals. Orbitals 1—2, rarely 1—1. Temporals 1+2. Labials seven, sixth largest. Infralabials seven, fourth largest. Anterior submentals much the larger. Scales smooth, in 15 rows, outer broader. Ventrals 161. Anal divided. Subcaudals 33—39 pairs.

Light reddish-brown, punctulate with black. A yellowish line on each flank, on the fourth and fifth rows of scales. Bases of all the scales darker, more distinctly so on the third and fourth rows, which presents the appearance of a darker line continued forward to the nostrils, darkening on the sides of the head and neck. Base of each ventral broadly

banded with black. The subcaudals are much lighter, which suggests the possible existence of a habit of carrying the tail raised from the ground. California.

CONTIA EPISCOPA, *pl. VI, fig. 2.*

LAMPROSOMA EPISCOPEUM Kennicott, 1859, *Mex. Boundary Survey Rept.*, p. 22, *pl. VIII, f. 2.*

CONTIA EPISCOPA Cope, 1875, *Check list*, 36.

Body stout, subcylindrical, tapering posteriorly; head not distinct from neck, short, depressed, crown flat; tail short, thick, conical. Eye moderate, pupil round. Nostril small, lateral. Nasal entire. Head-shields normal. Frontals short, broad. The upper angle of the rostral bent back between the internasals. Loreal small, sometimes united with the postfrontals. Orbitals 1—2. Labials seven, third and fourth in orbit. Infralabials seven, fourth largest. Temporals 1+2. Anterior submentals very large, posterior very small. Scales smooth, nearly square, in 15 rows, outer broader. Ventrals 145. Anal bifid. Subcaudals 45 pairs.

Uniform yellowish or greenish-brown, each scale with a lighter margin. Belly yellowish. Total length $8\frac{3}{4}$ inches; tail $1\frac{3}{4}$ inches. Texas.

NINIA.

Baird & Girard, 1853.

Moderately elongate; tail rather short. Eye small, pupil round. Teeth equal, smooth. Internasals and supraoculars small. Nasal divided. Loreal and anteorbital fused. Postorbitals two. Scales keeled, in 17—19 rows. Anal entire. Subcaudals bifid.

Prefrontals united	<i>liebmanni.</i>
tail short;	
labials seven; brownish-white to brown;	
a black collar	<i>sebae.</i>
labials six; brown to black;	
no collar	<i>sieboldi.</i>
tail long, slender;	
back black; belly white;	
a light collar	<i>diademata.</i>

NINIA LIEBMANNI.

CHÆSODROMUS LIEBMANNI Reinhardt, 1860, *Vidensk. Meddel. Kjøbenhavn.*, 242.

Head little larger than the neck; muzzle broad. Habit similar to other species of this genus. Prefrontals united, entering orbit. Internasals separate. Nasal divided. Loreal and anteorbital united. Postorbitals and supraocular united. Frontal broad. Temporals 1+2. Labials seven, third and fourth in orbit. Infralabials eight, fifth largest. Scales keeled, in 17 rows. Ventrals 130. Subcaudals 42 pairs.

Brown, with a transverse band of light color across the parietals. Mexico.

NINIA SEBAE.

STREPTOPHORUS SEBAE Dum. & Bibr., 1854, *Exp. Gén.* VII, 515.

NINIA SEBAE Cope, 1860, *Pr. Ac. N. Sc., Phil.*, 340 (name).

Of moderate size, subcylindrical, belly broad; head rather distinct from the neck, crown convex, muzzle broad and rounded; tail short, tapering, slender posteriorly. Eye small, pupil round. Head-shields nine. Rostral small. Internasals and supraoculars very small. Frontal short, broad. Prefrontals nearly as large as frontal, entering the orbit. Parietals large. Nasal in two parts, nostril between. Loreal large, fused with anteorbital. Postorbitals two, lower small. Temporals 1+2. Labials seven, third and fourth in orbit, the former extended backward to reach it. Infralabials seven, fourth large. Submentals two, anterior twice as large as posterior, not in contact with the mental. Scales keeled, in 17 or 19 rows, outer broad. Ventrals 146 to 151. Anal entire. Subcaudals 50 to 64 pairs.

Brownish white, scales tipped with dark. A row of small black spots on each side of the back. Head-shields black; labials more or less margined with the same color. Separated from the black of the head by about three scales there is a black collar on the neck, varying in length, extending to the outer rows of scales. Belly and labials glossy yellowish-white. Total length $11\frac{3}{4}$ inches; tail $2\frac{1}{4}$ inches. Mexico and Central America.

NINIA COLLARIS.

STREPTOPHORUS SEBAE VAR. COLLARIS Jan., 1865, *Icon. Ophid. Livr.* 12, pl. III, fig 6.

Differs from *N. sebae* in that it has no spots behind the broad collar on the neck, and in the punctulations on the hinder part of the abdomen. A variety of *N. sebae*.

NINIA SIEBOLDI.

ELAPIODES SIEBOLDI Jan., 1862, *Prodr. Icon. Gén. Ophid.*, II, 21.

Elongate, subcylindrical, belly flattened; head not distinct from neck, subconical, slightly depressed, narrowed in front; tail short, conical. Eye small, pupil round. Head-shields nine. Internasals very small. Pre-frontals large. Frontal broader than long, supraciliaries much shorter. Rostral small, little broader than high. Nasal divided. Anteorbital and loreal fused. One postorbital. Labials six, third and fourth in orbit, fifth largest; in the specimen figured it is in contact with the parietal. Infralabials seven, fourth largest. A pair of submentals, followed by large shields. Scales lustrous, iridescent, carinate in the dorsal rows to the tail, in 17 series, outer broader and smooth. Ventrals 146—154. Anal entire. Subcaudals 34—38 pairs.

Dark brown to black. Belly and throat white. Subcaudals darker at the bases. Outer row of scales marked with lighter. Labials and infralabials little lighter than top of head. (Jan.) Mexico.

NINIA DIADEMATA.

Baird & Girard, 1853, *Cat. N. A. Serp.*, 49.

All the upper portion of the trunk deep black; with a white collar; all the ventrals black in their central portion and margined with white, forming thus two white lateral bands. Labials six, third and fourth in orbit. Rather more slender than other *Ninia*e. Scales in 19 rows. Ventrals 145. Anal entire. Subcaudals 89 pairs.

The essential colors are the deep black and the pure white. Thus the under side of the head and the neck, which seems to be encircled by a collar, are of a beautiful white, as also the sides of the belly and tail where it forms a fine lateral band. The black covers the remainder of the body. The head is twice as long as wide. Total length 0^m, 347; tail 0^m, 067. Mexico. (B. & G.)

VIRGINIA.

Baird & Girard, 1853.

Rather small; body stout; belly broad; head distinct from the neck, subconical, depressed; tail short, thick, tapering to a point. Eye medium, pupil round. Crown-shields eight or nine. Internasals two or one. Nostril anterior, in the anterior portion of the divided nasal. Loreal and anteor-

bital united. Prefrontal entering the orbit. Postorbitals two. Anal and subcaudals bifid. Scales smooth or keeled, in 15 to 17 rows.

Scales in 17 rows,

keeled;

labials five;

one internasal

striatula.

two internasals

inornata.

smooth or keeled;

labials six

elegans.

scales in 15 rows, smooth or keeled;

internasals two;

postorbitals two

valeriae.

postorbitals three

harperti.

VIRGINIA STRIATULA, *pl. VII, fig. 2.*

COLUBER STRIATULUS Linné, 1766, *Syst. ed. XII, I, 375.*

Small, moderately slender, subcylindrical, belly broad; head small, not distinct from the neck, subconical, narrow in front; tail short, stout, tapering. Snout moderate, narrow. Eye small, pupil round. Mouth-cleft medium, slightly curved. Nostrils anterior, directed outward. Crown-shields eight. Prefrontals long and broad, entering the orbit. One internasal. Rostral subtriangular, very small. Nasal in two parts, nostril in anterior. No loreal, fused with anteorbital. Orbitals 1—1. Temporals 1+1. Labials five, third and fourth in orbit, fifth largest. Infralabials six, fifth largest. Posterior pair of submentals half as long as the anterior. Scales keeled, in 17 rows, dorsal narrow, outer broad and faintly carinate. Ventrals 119—130. Anal bifid. Subcaudals 25—46 pairs.

Uniform greyish or reddish-brown; light yellowish or reddish beneath. With or without a band of light color across the occipitals which spreads upon the posterior labials. Massachusetts to Mississippi.

VIRGINIA INORNATA *n. sp.*

Moderately stout, slightly depressed; head small, not distinct from the neck, subconical; tail short, tapering to a point. Snout short, blunt. Eye moderate, pupil round. Nostrils anterior, directed horizontally in the

anterior portion of the divided nasal. Nine head-shields. Rostral very small, not reaching the top of the head. Two internasals (left smaller in each specimen). Loreal elongate, with the prefrontal forming the anterior border of the orbit. Prefrontals as broad as long. One postorbital. Temporals 1+1. Labials five, third and fourth in orbit, fifth largest. Infralabials six, fifth largest. Posterior submentals half as large as the anterior. Scales carinate, lustrous, in 17 rows, dorsal narrow, outer broad and faintly keeled. Ventrals 125—129. Anal divided. Subcaudals 36 pairs.

Uniform brownish olive on the dorsal rows. No band on the occiput. Ventrals whitish, tinged with olive on the bases. Largest specimen total length $10\frac{1}{2}$ inches; tail $1\frac{1}{2}$ inches. Texas.

VIRGINIA ELEGANS.

Kennicott, 1859, Pr. Ac. N. Sc., Phil., 99.

“Resembles *V. valeriae*; vertical and occipital plates narrower. Dorsal scales very narrow and elongated, much more so than in *V. valeriae*, disposed in 17 rows.

“Color uniform light olivaceous brown above; dull yellowish white beneath. Readily distinguished from the nearly allied *V. valeriae* by the narrower dorsal scales in 17 rows instead of 15 as in that species.” Southern Illinois and southward.

VIRGINIA VALERIAE, *pl. VII, fig. 3.*

Baird & Girard, 1853, Cat. N. A. Scp., 127.

Moderate, subcylindrical, belly broad; head small, elongate, little larger than the neck, subelliptical; tail short, tapering to a point. Eye medium, over third and fourth labials, pupil round. Mouth-cleft deep, nearly straight. Snout narrow. Rostral erect, as broad as high. Nasal divided, nostril in anterior portion. No loreal. Internasals two. Prefrontals broad, entering the orbit. One elongate preocular, loreal and ocular united. Postoculars two (1 to 3). Temporals 1—2, anterior in a notch between the fifth and sixth labials. Frontal broad, subhexangular. Parietals elongate. Labials six, fifth and sixth larger. Infralabials six, fourth largest. Submentals two pairs, posterior rather larger. Scales smooth, or with a weak keel on those of hinder portion of the body, in 15 rows, dorsal broad, rhomboid, outer broader. Ventrals broad, 117—128. Anal bifid. Subcaudals 24 to 37 pairs.

Greyish-brown, with two to four irregular longitudinal series of black dots, sometimes absent. Belly uniform yellowish. Maryland to Georgia and Illinois.

VIRGINIA HARPERTI.

CARPOPHIS HARPERTI *Duncker & Bibrón, 1854, Exp. Gén. VII, 135.*

VIRGINIA HARPERTI *Cope, 1875, Check list, p. 35.*

Two triangular internasals. Prefrontals large, entering the orbit. Frontal pentagonal. Nasal bifid. Loreal and anteorbital fused, forming a long subrectangular plate. Postorbitals three, second largest. Temporals 1+2, anterior large, not in contact with orbitals. Labials six, third and fourth in orbit, fifth largest, in contact with the parietal by its upper angle. Infralabials six. Two pairs of elongate submentals, posterior divergent. Scales smooth, in 15 rows. Ventrals 111. Anal bifid. Subcaudals 32 pairs.

Yellowish or olivaceous gray, punctulate with blackish. Lower portions white. South Carolina. (D. & B).

CARPPOPHIS.

Gervais, 1843.

Body comparatively stout, subcylindrical; head small, not distinct from the neck, depressed; tail stout, conical. Crown-shields normal, short, broad, often fused or subdivided. Snout prominent, rostral bent back on the top. Nasal entire. Preocular united with the loreal, or very small. Scales broad, smooth, glossy, in 13 to 15 rows. Anal bifid. Subcaudals in two rows.

A small preocular;

 a pair of internasals

straminea.

 with transverse bands of black

cincta.

preocular united with the loreal;

 prefrontals and internasals fused

helenae.

 prefrontals and internasals separate;

 back reddish-brown

amoena.

 back black

vernix.

CARPPOPHIS STRAMINEA.

CHILOMENISCUS STRAMINEUS *Cope, 1860, Pr. Ac. N. Sc., Phil., 339.*

Internasals partly or entirely separated by the rostral. Frontal obtuse-angled in front. Labials seven, third and fourth in orbit. Infralabials

eight, fifth larger. Submentals two pairs, hinder half as long as the anterior. Temporals 3+3. Scales hexagonal on the flanks, elongate on the back, in 13 rows. Ventrals 117. Subcaudals 22 pairs. Length 9 inches; tail $1\frac{1}{2}$ inches. Lower California. (Cope.)

CARPHOPHIS CINCTA.

CHILOMENISCUS CINCTUS Cope, 1861, *Pr. Ac. N. Sc., Phil.*, 303.

Internasals separated by the rostral. Nasal grooved behind the nostril. One small anteorcular. Two postoculars. Labials 7, first long, remainder high, except the last two, which are nearly equilateral. Scales very broad, smooth, in 13 rows. Anal divided. Tail very short.

Reddish-white. Encircled by sixteen black rings upon the body and three upon the tail. Rings and spaces about equal, three to five scales each. Parietals and frontal black to the second labials. Chin shaded with black. Guaymas, Gulf of California. Possibly the same as the preceding. (Cope.)

CARPHOPHIS HELENÆ.

CELUTA HELENÆ Kennicott, 1859, *Pr. Ac. N. Sc., Phil.*, 100.

CARPHOPIUS AMOENA var. *HELENÆ*, Jan, 1862, *Arch. per la Zool.*

"Snout shorter and narrower than in *C. amoena*. A single pair of frontal plates. Color above lustrous chestnut brown; beneath pale yellowish (flesh-color in life), color of the abdomen extending to second lateral row of dorsal scales. Readily distinguishable by the suppression of the anterior frontals." Mississippi, Illinois.

CARPHOPIUS AMOENA, pl. VII, fig. 1.

COLUBER AMOENA Say, 1825, *Jour. Ac. N. Sc., Phil.*, 237.

CARPHOPIUS AMOENA Gerrais, 1843, *Diet. d'Hist. Nat.*, III, 191.

Small, moderately stout, subcylindrical, belly broad; head not distinct from the neck, crown convex, muzzle broad; tail short, near one fifth of the total, thick, conical. Eye small, pupil round. Mouth moderate, nearly straight. Head-shields nine, broad, excepting the supraciliaries, which are very small. Frontal hexagonal. Internasals small. Rostral about as broad as high, very convex. Nasal entire, the small nostril in its anterior half. No loreal. Orbitals 1—1, anterior elongate. One temporal. Labials five, third and fourth under the eye, fifth largest. Infralabials six, fourth

largest. Submentals two pairs, posterior less than half as large as the anterior. Scales smooth, in thirteen rows, medial a little longer than broad, outer and caudal as broad as or broader than long. Ventrals 112—131. Anal divided. Subcaudals 24—36 pairs.

Uniform lustrous reddish-brown above (more red in life); light yellowish beneath to second row of scales (flesh-color or red in life). Mississippi Valley to Illinois and to Massachusetts.

VAR VERMIS.

CELUTA VERMIS *Kem.*, 1859, *Pr. Ac. N. Sc., Phil.*, 99.

Larger than *C. amoena*. Prefrontals and internasals separate. Black color of the back reaching only to the third rows of scales. Missouri and Kansas. A variety of the preceding.

GEOPHIS.

Wagler, 1830.

Body elongate, subcylindrical; head scarcely distinct from the neck; tail short, subconical. Eyes small. Teeth equal, smooth. Crown-shields nine. Internasals much smaller than the prefrontals, rarely suppressed or fused. Nasal divided. Lorcal and anteorbital united. Postorbitals one to two. Scales smooth, in 15 to 19 rows. Anal entire. Subcaudals in two rows.

Scales in 15 rows;

back transversely banded with black;

parietal and labials in contact

semidoliatus.

scales in 17 rows;

back black;

frontal longer than broad;

parietal and fifth labial in contact

bicolor.

back plumbeous;

frontal broader than long;

parietal and labials not in contact

latifrontalis.

scales in 19 rows;

back crossed by bands of black;

parietals and labials not in contact

lippiens.

GEOPHIS SEMIDOLIATUS.

RABDOSOMA SEMIDOLIATUM *Dun. & Bibr.*, 1854, *Exp. Gén.* VII, 93.

Body slender; tail short, stout. Scales smooth, in 15 rows. Ventrals 172. Anal entire. Subcaudals 24 pairs. Whitish inferiorly; the white also serves as a ground color for large quadrilateral black or brown spots or transverse bands on the back, which are separated by spaces narrower than themselves, and about thirty in number. The black cap on the head is separated from the first spot on the back by a narrow collar of white. Trunk 0^m, 275; tail 0^m, 027.

“Cinq plaques sus-labiales; pas de squamme temporale. Dessus du corps coupé en travers par de grandes taches noires, sur un fond blanc. Plaque rostrale fort grande; bord antérieur de la frontale brisé sous un angle très-ouvert; susoculaires extrêmement courtes; première nasale ne descend pas plus bas que la seconde; point de préoculaire; une seule post-oculaire; cinq supéro-labiales, dont la 3^e seule touche à l’œil; deux paires de plaques inter-sous-maxillaires. Queue robuste. . . . Il n’y a qu’une seule squamme temporale, mais elle est extrêmement développée; elle est oblongue pentagone inéquilatérale et située le long de la plaque pariétale au dessus de la dernière supéro-labiale.” Mexico. (D. & B.)

GEOPHIS BICOLOR.

Günther, 1868, *Ann. and Mag. N. Hist.*, 416.

“Head rather broad, short, and depressed; body and tail of moderate length. Eye small. Anterior frontals about one fourth the size of posterior; vertical rather longer than broad, with the anterior angle very open; occipitals as long as postfrontals and vertical together, rounded behind; six upper labials, the third and fourth entering the orbit; the fifth is the largest, and forms a long suture with the occipital. The remainder of the temple is covered by scale-like temporals, 1+2. Two postoculars. Anterior chin-shields twice as long as posterior, in contact with four labials. Scales in seventeen rows, smooth, without apical groove. Ventrals 160—168; anal entire; subcaudals 39—48.

“Upper parts uniform black; below white; on the two or three outer rows of scales the white color appears in more or less distinct small spots, whilst the black of the upper parts extends to the angles of the ventral shields, each subcaudal black in front.” Largest 14½ inches long; tail 3 inches. City of Mexico.

GEOPHIS LATIFRONTALIS n. sp.

Moderately stout, tapering slightly toward the head, belly broad; head not larger than the neck, depressed, snout broad, rounded, crown a little convex; tail stout, conical, short, near one eighth of the total. Eye very small. Head-shields nine. Rostral broader than high, bent back in a blunt angle on the top of the snout. Internasals small. Prefrontals large, as broad as long, entering the orbit. Frontal short, broader than long. Supraoculars small, nearly as broad as long. Parietals large, about twice as long as broad. Nasal divided. Loreal and anteorbital united. One postorbital (two on one side, lower very small). Temporals 1+2, anterior large. Labials six, third and fourth in orbit, fifth largest, and separated from the parietals by a long temporal. Infralabials seven, fourth largest. Submentals two pairs, short, broad; posterior half as long as anterior, scale-like. Scales smooth, lustrous, in 17 rows, medial little longer than broad, outer broader than long. Ventrals 179, broad. Anal entire. Subcaudals in two rows, 32 pairs.

Back uniform dark plumbeous, tinged with purple; margins of scales lighter. Ventrals white, mottled with leaden to uniform leaden. Total length $15\frac{1}{4}$ inches; tail $1\frac{3}{8}$ inches. Found fifty miles south of San Luis Potosi, Mexico, by Dr. Edw. Palmer.

GEOPHIS LIPPIENS.

SYMPHOLIS LIPPIENS Cope, 1861, *Pr. Ac. N. Sc., Phil.*, 524.

Body stout, tail blunt, hardly twice as long as the head, ending in a convex shield. Muzzle prominent, obtuse. Teeth equal, smooth. Eye very small. Rostral large. Prefrontals as broad as long. Frontal long, right-angled at each end. Supraciliary and upper postocular fused, the former as broad as the frontal. Loreal in orbit. One small preocular over the loreal. Nasal united with the labial. Labials five, last not so high as long. One temporal. Scales poreless, broader than long, in 19 rows. Anal entire. Subcaudals bifid.

Yellow, with eighteen black bands. A band covers the muzzle to behind the eyes; two are on the tail. Total length $20\frac{3}{4}$ inches. Guadalajara, Mexico. (Cope.)

TOXICOPHIDIA.

Snakes with fixed grooved or movable tubular fangs, connected by ducts with special glands for the secretion of venom.

Fangs grooved, erect, immovable.

PROTEROGLYPHA.

Tail conical

CONOCERCA.

Tail compressed

PLATYCERCA.

Fangs tubular, reclining, erectile.

SOLENOGLYPHA.

No pit between the eye and nostril

ABOTHROPHERA.

A pit on the side of the face

BOTHROPHERA.

CONOCERCA.

ELAPIDAE.

Body elongate; head moderate, crown flattened; muzzle short, broad, rounded; tail stout, short to medium. Loreal generally absent. Fangs erect, grooved, without or with smaller smooth teeth behind them. Eyes small to moderate, pupil round in most genera. Scales smooth.

Elaps, the only genus of the family of which species are known in North America, is also found in South America, Africa, and the East Indies.

ELAPS.

Schneider, 1801.

Cylindrical, moderately to very slender; head indistinct, rounded, depressed, crown flattened, muzzle short and broad; tail short, stout. Eye small, pupil round. No other tooth behind the fang. Nasal in two parts. No loreal. Anteorbital one, sometimes fused with prefrontal. Postorbitals two, sometimes one. Scales smooth, in 13—15 rows. Anal entire or divided. The American species are distributed from the Southern United States to the Argentine Republic. The species mentioned in the synopsis are those regarded as tolerably well established.

- Black rings not in groups;
 first broad ring behind the occiput black *fulvius*.
 first broad ring behind the occiput red *euryxanthus*.
 black rings in threes;
 occipital band red; a yellow band in front of the eyes; middle
 band of group wider *decoratus*.
 occipital band black;
 black bands subequal, narrow; 13—14 groups *elegans*.
 occipital band yellow;
 middle band of each group much wider; 7—8 groups
laticollaris.

ELAPS FULVIUS, *pl. VIII, fig. 3.*

COLUBER FULVIUS Linné, 1766, *Syst. Nat.*, ed. XII, p. 381.

ELAPS FULVIUS Cuvier, 1817, *Repn. Anim.*, ed. I, p. 84.

Slender, cylindrical; head little broader than the neck, depressed, rounded; tail short, thick, conical, near one seventh of the total. Mouth-cleft medium, nearly straight. Head-shields nine, short, broad. Rostral low, broad, subtriangular. Nasal in two parts, nostril between, anterior larger. Orbitals 1+2, exceptionally one postorbital. Labials 7 (6—8), third and fourth in orbit, fifth, sixth, and seventh larger. Infralabials 7, fourth large, in contact with the posterior pair of submentals. Scales smooth, broad, in 15 rows, outer broader. Ventrals 202—236 (*tenere*, *B. et G.*, 224—236). Anal divided, sometimes entire. Subcaudals 25—44 pairs. Head and tail ringed with black and yellow, body with black, red, and yellow. Head black, with a yellow ring crossing the occipitals, its width equal to their length, widening downward. Body encircled by 13—20 rings of each of the black and the red, in some of the varieties the number is greater. The yellow are twice as many and serve as borders to the black, in some cases they are nearly obsolete. All the rings vary much in width, commonly the yellow ones are very narrow, and with the included red one, occupy about as much space as one of the black; frequently the black is the narrower. The red is generally spotted or dotted with black; specimens from Alabama and Florida show in the midst of the red band an oblong spot on each side of the vertebral line and a large rounded spot on the middle of the belly. Largest specimen two feet in total length; tail three inches. Hab. Southern United States to Mexico, and southward through varieties.

VAR. NIGROCINCTUS.

ELAPS NIGROCINCTUS Girard, 1854, *Pr. Ac. N. Sc., Phil.*, 226.

Head black to the middle of the parietals. Body encircled by 19—21 white-bordered black rings, separated by red spaces. Some scales and some of the ventral plates in the red spaces marked with black. Tail with about six rings, black and white. Ventrals in one specimen 222+55. The white color was probably yellowish in life. Mexico and Central America.

VAR. AFFINIS.

ELAPS AFFINIS Jan., 1859, *Rev. & Mag. Zool. (Prodr. Ophid., pp. 6 and 14, pl. B)*.

Head black from muzzle to postorbitals. A yellow space extending to the third or fourth scales on the neck is followed by bands of black three or four scales wide separated by red spaces of ten to fourteen scales each. In the red spaces the scales are tipped with black, and there are also black spots of greater size irregular shape and position. Ventrals 216; anal divided; subcaudals 40 pairs. Mexico.

VAR. DISTANS.

ELAPS DISTANS Kennicott, 1860, *Pr. Ac. N. Sc., Phil.*, 338.

“Body slender, with very narrow black rings, four or five scales in width, separated by intervals three or four times as wide, of brownish or reddish, entirely unspotted. No light rings separating the red and black ones. Under lip and jaw wholly without black, and the tip of the nose light.” Chihuahua, Mexico (Kenn.) Florida (Cope.)

VAR. APIATUS.

ELAPS APIATUS Jan., 1859, *Rev. & Mag. Zool. (Prodr. Ophid., p. 11)*.

Snout tipped with black. A spot of yellow in front of the eyes. A black band across the head through the eyes to the labials. A black collar behind the parietals four scales in width. Thirty-one black bands on the body; eight on the tail. The scales in the red spaces are not spotted with black, but here and there are scales entirely of that color. Ventrals 202; anal divided; eight entire subcaudals and 33 pairs. (Jan.)

VAR. EPISTEMA.

ELAPS EPISTEMA Dumeril & Bibron, 1854, *Exp. Gén. VII*, 1222.

Back with large spots of black, without rings. Black of muzzle extending to postorbitals. Occiput and throat yellow. Behind this a black collar

interrupted at the first abdominal shields. Large rounded light-bordered black spots, ten in number, widely separated on the body. Tail with three wide rings of black. Scales of back black-tipped; belly probably uniform red or yellow. Mexico. (D. & B.)

VAR. DIASTEMA.

ELAPS DIASTEMA *D. & B.*, 1854, *Erp. Gén. VII*, p. 1222.

Muzzle and vertex black. Occiput crossed by a band of white or red (yellow), behind which there is a black one extending under the throat. Fourteen or fifteen black white-bordered rings on the body, separated by white (red) spaces ten or twelve times as wide, in which the scales are black-tipped. A second specimen had reddish intervals and twenty-one rings. Mexico. (D. & B.)

VAR. CEREBRIPUNCTATUS.

ELAPS CORALLINUS var. *Peters*, 1869, *M. B. Berl. Akad.*, 877.

Snout to parietals black. A yellow band across the parietals. From the parietals a black yellow-edged band covers the first seven rows of scales. Ten narrow yellow-edged black rings on the body. Tail black and yellow. The scales in the broad red spaces are tipped with black. Pueblo, Mexico.

ELAPS EURYXANTHUS.

Kennicott, 1860, *Pr. Ac. N. Sc., Phil.*, p. 337.

"Head very small, narrower than the neck; entirely black as far back as the angle of the mouth. Banded alternately with black and light brick-red, separated by narrow rings of creamy white, all the bands immaculate. First broad ring behind the occiput red instead of black as in the other species." Sonoran region.

ELAPS LATICOLLARIS.

Peters, 1869, *Monatsh. Berl. Akad.*, 877.

Head black to the parietals. A yellow band across the parietals covers the first two rows of scales. Behind this a black ring of twelve or thirteen scales in width, then a yellow of three or four scales, and then a black of four or five precede the first red one, which occupies from six to nine scales. Similar red rings separate the eight or nine triads of black ones on the body. The middle ring of each triad is nearly twice as broad as the others, from which it is separated by narrow yellow spaces. The

red scales are black-tipped. Tail, with three broad black, separated by narrow yellow rings. Pueblo, Mexico.

ELAPS DECORATUS.

Jan., 1859, Rev. & Mag. Zool., Prodr. Icon. Ophid., pp. 7, 14, pl. B.

Rostral and internasals black. A black band on the frontals reaches the labials. A collar of black behind the parietals; behind this a narrower ring separated by four scales of yellow. A band in front of the eyes yellow; that on the occiput red. Seven scales behind the second black band on the neck are red, then commence the triplets of black. In each triad the outer bands are about half as wide as the middle, which occupies three or four scales, separated from the others by narrow spaces of yellow. Number of triads fourteen, separated by as many spaces of red, in which the scales are black-tipped. Ventrals 204—218. Subcaudals 14—20. Mexico.

ELAPS ELEGANS.

Jan., 1859, Rev. & Mag. Zool. Prodr. d'une Icon. Ophid., pp. 6, 13, pl. B.

Muzzle to postoculars black. A white parietal band is interrupted on the middle of the head. Behind this to the neck the head is black. A couple of series of white scales on the neck precede a large triangular black spot posterior to which there is a white (red) space of three or four black-tipped scales. From this point there are thirteen or fourteen triads of black rings, more or less confluent on the abdomen, separated by single series of white scales. The intervals between the groups is about the width of the rings themselves, three or four scales each. Tail black, with six white bands on single series of scales. Ventrals 199—213+29—37.

PLATYCERCA.

HYDROPHIDAE.

Sea serpents. Venomous. Elongate, subcylindrical anteriorly, compressed posteriorly; head rather indistinct, generally depressed; tail short, compressed into a paddle. Eye small, pupil round. Fang small, erect, grooved in front, followed by other teeth. Nostril valvular. Head-shields generally irregular. No loreal. Scales small, with or without keels, or tubercular. No claw at the side of the vent. Hab. tropical portions of the Indian and Pacific oceans, sometimes entering fresh waters.

PELAMYS.

Daudin, 1803.

Moderately long; neck rather large; head depressed; snout long, broad. No internasals. Nasals in contact, nostrils near the hinder margins. No loreal. Scales small, not imbricate. Ventrals indistinct or small. Lower jaw not notched in front. One species of this genus is frequently met with off the western coasts of Southern Mexico.

PELAMYS PLATURA.

Much compressed posteriorly; neck stout; head little larger than the neck, long, depressed; snout long, broad, and rounded in front; tail short, of moderate width. Head-shields abnormal. No internasals. Nasals longer than broad, nostril near the outer posterior angle. Prefrontals hexagonal, receiving the anterior angle of the frontal between them. No loreal. One anteorbital, sometimes two. Two to three postorbitals. Labials 7—9, second larger. Scales polygonal, commonly hexangular, not imbricate; lower with a convexity or tubercular keel; upper flat. In the specimen before me there are 53 rows around the middle of the body. Ventral scales similar.

Back (23 rows of scales) and top of head black. Upper lip and lower half of body yellow. Lower lip marked with black. Tail with three transverse bands of black, the rounded ends of which extend near the lower margin, where they alternate with four spots of black. Vent in a spot of black. Tip of tail black. The coloration varies greatly in different individuals. The specimen described was taken on the western coast of Nicaragua.

SOLENOGLYPHA.

BOTHROPHERA.

CROTALIDAE.

Stout, fusiform; head large, distinct from the neck; subtriangular, crown flat; tail short, with a rattle formed of horny rings, or tapering to a point as in the Colubers. Pupil oblong, erect. A pit between the eye and the nostril. Fangs tubular. Venom glands behind the eye at the side of the skull. Scales keeled. Anal entire.

With a rattle;

parietal shields scale-like; frontal absent or divided

CROTALUS.

frontal and parietals present

SISTRURUS.

without a rattle;

frontal and parietals present

ANCISTRODON.

crown-shields scale-like;

subcaudals in two rows except near extremity

LACHESIS.

subcaudals in two rows, rarely entire;

supraciliaries normal

TRIGONOCEPHALUS.

supraciliary small, commonly with a row of small prominent scales along its edge

TELEURASPIS.

CROTALUS.

Linneé, 1754.

Tail terminating in a rattle. A deep pit in front of the eye. Top of the head covered with scales, with several larger plates in front, or, in the subgenus *Crotalophorus*, having parietals and frontals as in the *Colubridæ*. Anal entire. Subcaudals simple, a few of the posterior sometimes divided.

Parietals and frontals scale-like;

nasal divided;

scales in 29 to 31 rows; keels swollen or tubercular;

a dorsal series of diamond-shaped spots

durissus.

scales in 25 to 29 rows; keels not tubercular;

a dorsal series of diamonds, varying to oblong subquad-rangular spots or to transverse bands

adamanteus.

scales in 23 to 25 rows;	
a dorsal series of more or less irregular and broken	
transverse bands	<i>horridus.</i>
nasals undivided;	
supraciliary produced as a sort of horn;	
scales in 21 to 23 rows	<i>cerastes.</i>
“supraciliaries and other large plates smooth”	<i>lepilus.</i>
nasals divided;	
scales in 21 to 23 series, slightly carinate	<i>tigris.</i>
scales in 23 to 25 rows	<i>lugubris.</i>
parietals and frontals like those of the colubers;	
scales in 23 to 25 rows	<i>catenatus.</i>
scales in 21 to 23 rows	<i>miliaris.</i>

CROTALUS DURISSUS.

Linné, 1758, Syst. Nat., ed. X, I, 214.

Stout, elongate, fusiform; belly broad; head large, triangular, tail short, thick, with more or less acuminate rattle. Eye small. Internasals triangular. Frontals two pairs. Scales of occiput and temples keeled. Rostral touched by six plates. Two anteorbitals; upper large, quadrangular; lower elongate, narrow, sometimes crowded from the orbit. Two loreals. Five small orbitals beneath and behind the eye. Three (2—5) rows of scales between suborbitals and labials. Labials 13—16, first and fifth rather larger; infralabials 14—18. Submentals one pair, large, or, through division of the anterior pair of infralabials, two pairs, anterior small. Scales lozenge-shaped, with strong swollen keels, in 29—31 rows, outer broader, outer row smooth. Ventrals broad, 175—183+19—33 (175—199+19—33.)

Yellowish-brown. A light-edged dark band across the head, through the eyes to the angle of the mouth. A similar band of dark brown from the head behind the supraciliaries on each side of the neck. These are followed by a dorsal series of 25—27 (25—30) diamond-shaped spots of brown, with lighter center and yellow borders occupying single rows of scales. The borders continue upon the flanks there inclosing rhombs of the ground color, and below them a series of half-rhombs which alter-

nate with a series of smaller spots on the lower edge of the side. Belly yellowish, clouded or mottled. Hab. Mexico to Brazil.

CROTALUS TRISERIATUS.

CROTALUS TRISERIATUS *Wiegman, Mus. Berol.*

CROTOPHISUS TRISERIATUS *Wagler, 1830, Syst. Amph., p. 176.*

"Notaco olivaceo serie macularum in spina dorsi irregularium rhombearum, rufuscenti-fuscarum, margine antico nigro-limbatarum, serie macularum minorum, ejusdem formæ ac coloris in utroque latere, posteriorum cum mediis confluentium, anteriorum a mediis taeniola pallida diremtarum; gastraco nigricante versus collum albescente; vitta pone oculos rufo-fusca." Mexico. (Wagler.)

CROTALUS ADAMANTEUS.

Bourcuis, 1799, Trans. Am. Phil. Soc. IV, 368.

Stout, fusiform; head large, triangular, broad behind, covered with small scales; tail short, thick. Eye moderate. Two or more pairs of rugose small internasals and prefrontals. Head covered with small keeled scales. Supraciliaries large, rugose, separate from the rostral by three (3—4) plates. Rostral higher than broad. Anterior portion of nasal larger, quadrangular, posterior narrow. Anteorbitals two; lower small, angular. Two loreals, or upper united with a small prefrontal. Sub and postorbitals five to six, small. Three (3—4) rows of scales between suborbitals and labials. Labials 14—16, first and fourth larger. Infralabials 15—19. Submentals one pair, large. Scales pointed, with strong keels, in 27—29 rows, outer broad, with keels obsolete. Ventrals broad, 169—179+1+21—32 (169—188+19—32.)

Yellowish-brown, with a dorsal series of 27—36 rhomboidal yellow-margined, light-centered, dark-brown spots. Single series of yellow scales form the margins; these meet on the fifth or sixth rows on the flank, where they inclose an alternating series of less distinct rhombs and an opposite series of half-rhombs on the outer rows. Posteriorly the spots fuse and become transverse bands. Belly yellowish, more or less blotched or punctulate with brown on the bases of the scutes. Tail black or barred with bands of black. A dark band included between narrow white lines from the forehead through the eye to the angle of the mouth. A narrow white line on each side of the pit, nostril, and on each edge of the rostral shield;

Hab. North Carolina to Texas, and through varieties to California and Mexico.

VAR. SCUTULATUS.

CAUDISONA SCUTULATA Kenn., 1861, *Pr. Ac. N. Sc., Phil.*, 207.

Rattle and rostral plates as in *C. lucifer*. Dorsal rows 25; superior labials 16; 3—4 rows of interorbital scales, bounded in front by two shields. Yellow stripe from eyebrow above rictus oris. Yellowish-brown, with a dorsal series of truncate, brown, yellow-edged rhombs; tail black-ringed. (*Kenn.*) Arizona.

Colors resembling somewhat those of *C. adamanteus*. About thirty dorsal rhombs. Rostral higher than wide, upper angle acute. Internasals and prefrontals regular, without small scales or plates around them. A pair of plates between the front portions of the supraciliaries and a pair of smaller ones behind them. Anterior portion of the nasal twice the size of the posterior. Two small loreals, one above the other. One or two rows of scales between the labials and suborbitals. Scales in 25 series. Ventrals 172+21. Taken by Dr. Edw. Palmer near San Luis Potosi, Mexico.

Var. ATROX.

CROTALUS ATROX Baird & Girard, 1853, *Cat. Serp.*, pp. 5 and 156.

Marked by bleached colors, shortening of the rhombs in the dorsal series and increase in their number and tendency posteriorly to fuse with the lateral spots. Sometimes the anterior infralabials are divided, forming a small pair of submentals behind the mental. Labials 14—19. Scales in 25—27 rows (25—29), outer two or three smooth. Ventrals 179—186+20—24, one to several of the subcaudals paired. Hab. Texas to Mexico.

CROTALUS MOLOSSUS.

CROTALUS MOLOSSUS Baird & Girard, 1853, *Cat. Serp.*, 10.

Two internasals. Four prefrontals. A pair between the supraciliaries in contact with each other and the prefrontals. Labials 18. Infralabials 17. Scales in 29 rows.

Yellowish. With a dorsal series of rhombs similar to those of *C. adamanteus*, with the outer angle produced down the side as a vertical bar, centers light, confluent anteriorly. Ventrals 187+25. New Mexico.

CROTALUS CONFLUENTUS.

CROTALUS CONFLUENTUS Say, 1823, *Long's Exp.* II, 48.

Distinguished from *atrox* by the subdivision of the head-shields, a greater shortening of the spots in the dorsal series, and the ashy colors. The number of spots and bands is increased. Internasals divided so as to place small plates between them and the nasals. Loreals two or one—the upper being united with the small prefrontal. A single scale between the anterior suborbital and the labial. Labials 14—17. Scales in 25—29 rows, outer two or three smooth. Ventrals 178—186+19—27, posterior subcaudals bifid or entire. Spots in dorsal series 41—54. Young with the white lines very distinct on the head. A lot of specimens from the same locality have the rows of scales in 25, in 27, and in 29 rows.

Specimen 2455 (*Mus. Comp. Zoöl.*), taken near Ft. Hays, Kansas, by Mr. J. A. Allen, has a supraciliary produced as in *C. cerastes*, two loreals on one side and one on the other, and 29 rows of scales. *C. pyrrius*, I am inclined to think, was founded upon an individual variation.

CROTALUS LUCIFER.

CROTALUS LUCIFER Baird & Girard, 1852, *Proc. Ac. N. Sc., Phil.*, 177.

Spots or rhombs broader than in *atrox* or *confluentus*. Colors generally darker. The black band from the eye backward passes above the posterior labials. Nasal and anterior labial frequently separated by small scales. One or two scales between the anterior suborbital and the labials. Loreal one, sometimes two. Internasals subdivided, separated by a row of small scales from the nasals (sometimes from rostral also). Labials 13—16. Scales in 25 (25—27?) rows. Ventrals 169—179+19—24. Spots and bands 30—42. Hab.

CROTALUS EXSUL *n. sp.*

Comparatively slender; neck small; head large, broad behind, outline as seen from above a suboval; snout short, broad; tail short. Eye moderately large. Head covered with small imbricate striate scales; a pair of larger ones in contact with the rostral and each other, another pair in contact with the nasals, but separated by a pair of small prefrontals. A larger scale on each side between supraciliary and postnasal. Supraciliaries large, striate. Rostral higher than wide, subtriangular, rather pointed above. Anterior

nasals large, narrowing toward the rostral; posterior higher and shorter. Loreals two, or upper fused with prefrontal. Anteorbitals two, upper large, lower elongate, narrow. Three plates surround the pit, in front of which is a group of smaller scales (7—11). Sub and postorbitals 6—7. Anterior orbital separated from the sixth labial by two scales. Cheek scales larger than labials, smooth. Labials 16—17, sixth largest. Anterior pair of infralabials small, not in contact. A pair of moderately large and broad submentals, separated in the anterior half of their length by a pair of wedge-shaped plates. The mental is separated from the submentals, and the anterior infralabials from each other by a pair of polygonal plates somewhat larger than the labials. Scales in 27 rows, striate and keeled, except the outer two. Ventrals 188+24. Rattle resembling that of *adamanteus*. In the specimen described, which is quite small, from the hindmost ring of the rattle to the fifth there are eight which show a very slight increase in size; in front of these the next has suddenly enlarged, and from this to the twelfth and most recent the size remains the same.

Light greyish-brown. Shapes of markings similar to those of *C. lucifer*. About thirty-three rounded, dark-edged spots on the back, separated by whitish spaces of a scale in width, and twice as many small spots on the lower edge of the flank, which also bears faint indications of a series alternating with the dorsal. Tail, with a dorsal series of black spots opposed to others on each side, with which they are more or less confluent. Belly uniform white or clouded lightly with olive near the flanks. Head nearly uniform brownish-grey, with faint indications of a light band from the frontal region to the central and hinder labials. A very faint band of light behind the eye toward the angle of the mouth. Total length 20 inches; tail $1\frac{1}{2}$ inches. Hab. Cedros Island, Lower California. Prof. Alex. Agassiz, two specimens.

CROTALUS HORRIDUS, *pl. IX, fig. 1.*

Linné, 1758, Systema Nat., ed. X, I, 214.

Stout, elongate fusiform; head distinct, triangular, very broad behind; neck small; tail short, thick, slightly compressed, outlines of the rattle approaching the parallelogramic. Head with one or more pairs of rugose plates in front. A pair of internasals, generally present, separated from the large elongate supraciliary by a single plate. With or without a pair or more of small frontals. Rostral higher than broad. Nasal in two por-

tions, anterior large, posterior narrow. Generally two small loreals, sometimes more. Eye rather large. Two anteorbitals; lower small, triangular, touching the orbit in a sharp angle. Sub and postorbitals five or six, small, anterior separated from the labials by one or two scales. Cheek scales larger, smooth. Labials 12—16, first and fifth larger. Infralabials 13—18. A pair of large submentals. Scales keeled, in 23—25 rows, outer broad, nearly or quite smooth. Ventrals 165—175+19—25.

Brownish yellow to yellowish-brown or black. A dorsal and two lateral series of light-edged black spots, which become confluent into transverse bands posteriorly. Anteriorly the spots appear as if formed of sections of a zigzag band. Tail banded in young, black in adults. The young have the spots lighter in the center; a pair of elongate spots extend from the back of the head on the neck; a band extends from the eye across the angle of the mouth, a series of small oblong spots on the flank alternates with those of the other series, and occasionally the anterior spots of the dorsal series are bifid. There is great variation in shape and size of spots and in depth of ground color; some are almost black, others are of a sulphur yellow. The sides of the rattle are nearly parallel in old specimens, and commonly there is a secondary groove or series of indentations between the median groove and the upper edge. In a specimen of a total length of fifty-four inches the tail is less than four. Hab. Massachusetts to Mississippi.

CROTALUS CERASTES.

Hallowell, 1854, *Pr. Ac. N. Sc., Phil.*, 95.

Small; head small, not angulate, crown tubercular. A single large nasal. Lateral edge of supraciliary produced above the eye so as to resemble a horn. Labials 11—13. Rows of scales 21—23. Ventrals 146+17.

Yellowish, with a dorsal series of indistinct brown blotches, below which on the flanks are irregular series of brown dots. A brown stripe from the orbit over the angle of the mouth. (From descr.) Hab. desert regions about the Gila and Colorado rivers.

The supraciliary is sometimes produced upward in specimens of *C. confluentus*, which see.

CROTALUS LEPIDUS.

CAUDISONA LEPIDA Kenn., 1861, *Pr. Ac. N. Sc., Phil.*, 206.

"Head ovoid, tapering to the nose, which is very narrow, pointed, and much depressed. Nostril very small, circular, and placed near the point of the nose in about the middle of a single nasal. Two elongated frontals (internasals) in contact, extending behind the nostrils. Supraciliaries and other large plates smooth. Rostral subtriangular, broader than high, the apex turned back slightly upon the crown. Upper preorbital small, and separated from the postnasal by the width of two larger plates. Labials rather large, 12 above, 10—12 below. Color of head yellowish ash."

"By the smoothness and size of the plates and absence of the horn, it will at once be distinguished from *C. cerastes*." Two heads from Presidio del Morte and Eagle Pass were described. Species of doubtful position.

CROTALUS TIGRIS.

(Kenn.) Baird, 1858, *U. S. and Mex. Bound. Survey Rept.*, 14, pl. IV.

Slender; head small, depressed. Supraciliaries and frontals smooth. Four frontals, six postfrontals (does this mean a pair of internasals, a pair of prefrontals and dissected frontal and parietals?) Two rows of scales between suborbitals and labials. Labials 14 above, 13—14 below. Series of scales 21—23, very slightly carinated.

Color yellowish ash above, with rather small, indistinct dorsal brown blotches anteriorly; two posterior thirds of body banded with brown. (Kennicott.) Deserts of Gila and Colorado.

Two internasals. Two prefrontals. Two shields between the supraciliaries, separated somewhat in front by a triangular shield; behind these irregular scales or small shields. Labials 10, a small shield separating the fourth and fifth from the eye. Series of scales 21. Ventrals 160+21.

Brownish, with seven series of spots, median larger, confluent posteriorly. A dark band from the lower border of the orbit above the angle of the mouth. (*C. intermedius*.) Mexico.

CROTALUS TRISERIATUS.

Wagler, 1830, *Syst. Amph.*, 176.

A pair of internasals. Four or more prefrontals. Frontal region covered with scales or small polygonal plates. Labials 12—14. One to two scales

between suborbitals and labials. Series of scales 23—25, outer two smooth. Ventrals 137 to 154—166⁺ 21—31.

A pair of oval dark-brown spots on the occiput. An elongate spot on each side of these from the supraocular above the hinder labials. A bar from each internasal through the eye to the angle of the mouth. A bar from the eye to the labials. A light line across the head on the supraoculars. A pair of large elongate spots on the neck between the hinder extremities of the bars from the supraoculars. Body with three series of spots on each side, upper two more or less confluent; the upper is larger and darker, and has somewhat the appearance of a medial series that has been divided by a vertebral line. Infralabials with three or four spots on each side. Belly irregularly spotted with brown, darker backward. Mexico.

(*Sistrurus*.)

CROTALUS CATENATUS, *pl. IX, fig. 2.*

CROTALINUS CATENATUS *Reif*, 1818, *Am. Month. Mag.*, IV, 41.

Larger than *C. miliaris*, with which it agrees in plan of coloration, except in the possession of a dorsal band of red, which, however, is not always present in that species. It is readily distinguished by a large anteorbital in contact with the nasal and prefrontal, a small subtriangular loreal, which does not reach the prefrontal, and a greater number of dorsal rows and of ventral scutes. Rostral high, with seven sides, all concave. Nasals large. Loreal small, subtriangular, acute-angled between the anteorbital and nasal. Anteorbitals two; upper large, much longer than the lower, bent upward to meet the prefrontal, in contact with the nasal; lower narrow. Sub and postorbitals 3—5. One to five small scales in front of the pit. The large suborbital in contact with the fourth and fifth labials. Labials 11—14. Scales in 23—25 rows, one to three of the outer smooth. Ventrals 136—153+20—29. Spots in the dorsal row, 35—48.

Varies from light ashy brown to black on the back. Beneath the range is from yellowish, spotted or punctulate with olivaceous, to dark olive or slaty brown, flecked with lighter. In cases the spots on the flank are separated by very narrow lines of light color. Specimens from the more exposed prairies are apt to be ashy and faded above and very dark beneath. Seven series of spots, dorsal large and more or less emarginate in front and

behind, along the middle of the body; frequently one or two of the lateral series are obsolete.

Top of head brown, with a light band on the fore parts of supraciliaries and frontal, a brown band from each parietal shield to the first blotch on the neck, a brown band from the eye to the neck, separated from the parietal by a light space, a light band from the nostril around the angle of the mouth, and a vertical bar of light color on each side of the pit. Labials dark. Lower lip with two light spaces on each side. Marks of head sometimes obsolete. Hab. Mississippi Valley from Ohio.

CROTALUS MILIARIUS.

Linneé, 1766, Syst. Nat., ed. XII, I, 372.

Small. Body fusiform; head distinct, broad behind, narrow and angular forward, with flat crown; tail short, rattle small. Eye small. Rostral high, broad near its upper extremity. Anterior nasal much larger. Nostril small. Loreal four-sided, largely in contact with the prefrontal. Two anteorbitals, upper much broader. One to four small scales in front of the pit. Sub and postorbitals four to six, anterior large. Check scales large, smooth. Labials 10—12. Infralabials 8—13. Scales in 21—23 rows, median pointed, all keeled, or the outer smooth. Ventrals 130—139+27—35, posterior subcaudals often bifid.

Greyish or ashy brown, varying from quite light to very dark. A dorsal series of 30—43, large light-edged more or less irregular black spots, band-like or divided toward the tail. Three series of alternating smaller spots on the flank, the upper often indistinct—sometimes obsolete, the lower on the edge of the abdomen. The majority of specimens have a narrow reddish band along the middle of the back, more distinct between the spots, though apparently crossing them. Three series of alternating spots on the belly, blending posteriorly, commonly irregular forward. Tail with six to twelve transverse bands. Head marked as in *C. catenatus*. Usually the parietal bands do not meet on the neck in the first spot of the dorsal series. The light band across the frontal and supraciliaries is sometimes indistinct, as are also the vertical bars on each side of the pit. The light band from the nostril around the angle of the mouth and the two spots on each side of the lower jaw are more distinct. The dark band from the eye to the neck varies much in length. Hab. Southern United States to Mexico.

VAR. RAVUS.

CROTALUS RAVUS Cope, 1865, *Pr. Ac. N. Sc., Phil.*, 191.CAUDISONA RAVA Cope, 1875, *Check List*, p. 33.

Scales in 23 rows. Ventrals 147+26.

Color yellowish, with from 26 to 31 elongate narrow parallelogramic deep brown spots, four scales long to five wide, and a series of as many short, transverse bars on the sides opposite them; a series of thrice the number of small spots on the interior rows of scales. Head pale, without spots or marks, except a minute punctulation. A brown anteriorly furcate nuchal spot. Table land of Mexico. (From descr.)

ANCISTRODON.

Beauvois, 1799.

Head triangular; snout prominent, its sides forming an acute angle with the top; tail without a rattle, tapering to a point. Head-shields nine. Frontal and parietals large. Loral present or absent. Scales in 23—25 rows.

A loral; scales in 23 rows

contortrix.

no loral; scales in 25 rows

piscivorus.ANCISTRODON CONTORTRIX. (*Copperhead*.) *Pl. VIII, fig. 1.*BOA CONTORTRIX *Linnaé*, 1766, *Syst. Nat.*, ed. XII, I, 373.ANCISTRODON CONTORTRIX *Baird & Girard*, 1853, *Cat. Serp. N. Amer.*, 17.

Body moderately stout, elongate fusiform, belly broad, back slightly compressed; head distinct, subtriangular, broad behind, crown flat; tail short, pointed. Head-shields nine. Parietals more or less broken posteriorly. Many specimens have small scales at one or both ends of the frontal. Rostral broad to the upper extremity. Nasal in two parts, anterior larger. Loral moderate, in contact with prefrontal. Anteorbitals three, upper large, middle narrow, lower very small. Sub and post-orbitals 4—6, separated by one scale from the third and fourth labials. Supralabials eight, first two and last smaller. Infralabials ten, second and last smaller. One pair of submentals large, followed by two pairs of smaller ones. Scales in 23 rows, outer broad, smooth, or faintly keeled. Ventrals 146 to 154. Subcaudals 42—49, commonly some of the posterior bifid.

Light reddish-brown, lighter below, with fifteen to twenty inverted Y-shaped darker spots, which vary in position from opposite to alternate, on each flank. About twice as many dark spots appear on each edge of the abdomen, sometimes spreading and confluent. All the scales are punctulate with brown. Head brighter, more copper-colored, sides with a band of cream color, bordered by a narrow line of light, inclosed by another of dark, passing from the upper postorbital above the labials around the angle of the mouth, and forward through the middles of the infralabials. A small round, light-edged spot of dark brown near the inner edge of each parietal. Tail darker. Terrestrial.

ANCISTRODON PISCIVORUS. (*Moccasin.*) *Pl. VIII, fig. 2.*

CROTALUS PISCIVORUS *Lacépède*, 1789, *Hist. Serp. II*, pp. 130 and 424.

ANCISTRODON PISCIVORUS *Cope*, 1859, *Pr. Ac. N. Sc., Phil.*, 336 (*name*).

Body stout, fusiform, belly broad, back slightly compressed; head distinct, subtriangular, broad behind, crown flat; tail short, pointed. Head-shields nine. Occipitals more or less broken posteriorly. Rostral broad to the upper extremity. Nasal in two parts, subequal. No loreal. Anteorbitals three, upper large, middle small, not reaching the nasal, lower very small. Postorbitals four (3—5.) Supralabials eight (7—9), third larger, reaching the orbit, first and last small. Infralabials ten (10—11.) Submentals one pair large, followed by two pairs of small. Scales in 25 rows, outer faintly keeled. Ventrals 136 to 145. Subcaudals 42 to 54, some of the posterior bifid.

Brown, reddish or olive. Eleven to fifteen more or less irregular and badly defined vertical bars or pairs of bars of dark brown, with lighter centers, on each flank. Tail dark brown or banded. Brownish-yellow beneath, with blotches of dark, which sometimes spread over the entire abdominal surface. Head uniform above, light-colored specimens showing a small round spot of dark near the inner edge of each parietal. A yellow-edged, dark band as wide as the eye passes from the postorbitals above the angle of the mouth to the neck. Three similar bands on the infralabials of each side. The head-markings are sometimes obsolete. In the variety *pugnax* from Texas the second labial is narrowed or crowded up. The total length of a large specimen is 44 $\frac{1}{4}$ inches; tail 6 $\frac{1}{2}$ inches. Aquatic. Southern States, from the Carolinas to Texas.

The following is taken from a Mexican specimen: Markings similar to those of *A. piscivorus*, posterior part of body almost black, general appearance very dark. Labials eight, third and fourth below the eye, separated from it by the sub and postoculars (5). Two antecoculars. No loreal. The specimen has five pairs of submentals, the anterior nearly twice the size of those next following. Scales in 24 rows. Ventrals 135. Subcaudals one pair, plus eleven entire, plus thirty-one pairs. Mexico.

LACHESIS.

Daudin, 1803.

Body long, back compressed, belly broad; head large, broad behind, crown flat; tail short, pointed. Fangs and pit as in *Crotalus*. Pupil vertical. Crown covered with small keeled scales. Scales with tubercular or swollen keels. Tail without a rattle, pointed. Anal entire. Subcaudals in two rows, except near the extremity, where they are small and keeled like those of the upper surface. Brazils to Mexico.

LACHESIS MUTUS.

CROTALUS MUTUS *Linne*, 1766, *Syst. Nat.*, ed. XII, 373.

LACHESIS MUTUS *Daudin*, 1803, *Hist. Rept.* V, 351.

Resembles *Crotalus durissus* in form and squamation. Elongate, tapering to neck and tail, flanks compressed, belly broad; head large, broad behind, crown flat; tail short, near one twelfth of the total, thick, pointed. Eye moderate, pupil erect. Crown-shields small, keeled. Supraciliaries large. A pair of internasals, separated by small scales. A few large scales about the pit. Labials ten (9—10), third large and below the pit, sometimes a small plate wedged between the second and third. Infralabials 15—16, anterior largest. Submentals one pair of large, followed by several pairs of small ones. Scales lozenge-shaped, broad, keeled, in 33—35 rows, outer broader and faintly keeled, inner with tubercular or swollen keels. Ventrals broad, 221—230. Anal entire. Subcaudals 33—35 pairs, followed by small keeled scales under about one sixth of the length of the tail.

Yellowish-brown, with about twenty-eight (27—29) light-centered brown lozenges on the back, more or less united along the vertebral rows, and produced toward or upon the ventrals. Flanks between the dorsal lozenges with or without spots or blotches of brown. Tail with 4—7 transverse

blotches of brown or black, separated by narrow spaces of light color. Lower surface yellow, somewhat clouded posteriorly. Head spotted with dark on the parietal region, with a transverse band of dark in front of the supraciliaries through the eye to the hinder labial. Description from Brazilian specimens. Species said to occur in Southern Mexico.

VAR. STENOPHRYS.

LACHESIS STENOPHRYS Cope, 1875, *Batr. and Rept. Costa Rica*, 152.

Muzzle short, depressed; rostral an equilateral triangle. Crown-scales flat, hexagonal, faintly keeled. A trapezoid plate in front of the large superior preocular. Four rows of scales between labials and orbits. Labials nine, third largest, second low and not bounding the pit. Pit bordered by three plates; upper narrow, in contact with both preoculars; lower wider, above the third labial; anterior above the second labial. Supraciliaries narrow, separated by twelve series of scales. Infralabials thirteen, first large. Submentals one pair, truncate in front, narrowly rounded behind. Scales in 36 rows, ten on each side of the median line with a central tubercle. Ventrals 200. Subcaudals 32 pairs, plus 17 transverse series of small ones under the extremity of the tail, terminal spine well developed.

Brown, with twenty-three reddish-brown rhombs on the median rows; the lateral angles are dark spots on the flank, sometimes isolated. On the middle of the body the rhombs have pale centers, posteriorly they are darker, irregular, confluent into a zigzag band. Tail dark brown, with narrow light cross-bands. Lower surfaces greenish-yellow, except the throat and chin, which are white (in spirits). A black band from the eye above the labials, broken upon the neck into a series of spots. Top of head uniform brown. Total length 0^m, 495; tail 0^m, 50. (Cope.) Sipurio.

TRIGONOCEPHALUS.

Oppl, 1811.

Elongate fusiform, back slightly compressed; head large, distinct from the neck, depressed, triangular, broad behind, pointed in front; muzzle prominent, angular; tail short, tapering to a point. Crown-shields small, scale-like. Eye moderate, pupil vertical. Scales keeled. Anal entire. Subcaudals two-rowed, rarely simple. Mexico to Patagonia.

Subcaudals bifid:	
scales in 27—29 rows	<i>asper.</i>
scales in 23—25 rows	<i>atrox.</i>
subcaudals entire:	
scales in 27 rows	<i>affinis.</i>

TRIGONOCEPHALUS ASPER.

var. n. C. LANCEOLATI.

Body long, tapering to head and tail; head large, broad behind, pointed and angular in front; crown flat, covered with small keeled scales. A pair of internasals, in contact behind the rostral; behind these on each side a larger plate rests against nasal, loreal and anteoocular, but is separated from the large supraciliary by a small scale. Rostral subquadrangular. Nasal in two parts, anterior larger. Loreal quadrilateral. Anteooculars two; upper much larger, sometimes divided; lower half as long, narrow. Suboculars narrow, fused, separated from the labials by a scale. Postoculars 2—3. Labials seven, posterior four large. Infralabials 10—11. Submentals one pair, not in contact with the mental. Scales keeled, in 27 (27—29) rows. In large specimens the keel of the median rows is swollen, and does not reach the end of the scale.

Reddish-brown, with Y-shaped marks of light color on the flanks, which, when opposed, inclose rhombs of the ground color or lighter. When regularly opposed the pattern resembles that of *Crotalus horridus* or *Lachesis mutus*. Flanks lighter with one or more series of spots near the abdomen and upon its edge. Belly and sides of head yellowish, the former mottled posteriorly, the latter with a brown line from the eye to a point near the angle of the mouth. The specimens described were taken at Obispo, on the Isthmus of Darien, by Dr. Maack. Compared with specimens from Martinique and St. Lucia, they are distinguished by brighter colors and swollen keels. The pattern of markings is similar. The numbers and shapes of labials and other head-plates are the same.

TRIGONOCEPHALUS ATROX.

COLUBER ATROX Linné, 1754, *Mus. Ad. Fridr.*, Tab. 22, fig. 2, and Gmelin, 1788, *Syst. Nat.*, 1107.COPIAS ATROX Merrem, 1820, *Tent. Syst.*, 154.

Body long, moderately slender, tapering to head and tip of tail; head broad behind, pointed in front, crown flat, covered with small keeled scales;

rostral angles sharp; tail short, near one seventh of the total, tapering to a slender extremity. Eye moderate, pupil erect. Anterior portion of the nasal larger. Rostral higher than wide. A pair of small internasals, commonly in contact; a similar shield over each loreal in contact with the former. Labials eight (7—8), separated from the orbit by one or two scales, second entering the pit, posterior small. Infralabials ten, anterior pair large, in contact. Submentals two pairs, posterior smaller, and separated by smaller plates. Scales elongate, keeled, in 25 (23—25) rows, outer a little broader. Ventrals broad, 199—205. Anal entire. Subcaudals 60—71 pairs.

Olivaceous, greenish or yellowish; dark-clouded to uniform yellowish below. With scattered transverse dark-edged bands of brown, sometimes divided along the vertebral row of scales, and tending toward alternation. The variation in respect to marking is very great in this species. Instead of transverse bands some specimens have triangles on the flanks. Others have the bands much broken, sometimes forming dorsal and lateral series of spots more or less irregular. Generally the bands are separated by spaces much wider than themselves. A dark band from the eye to the neck. Light-colored specimens show a lighter border to the spaces between the bands; often the center of the space is dark. Brazil to Central America, possibly to Mexico.

TRIGONOCEPHALUS AFFINIS.

BOTHROPS AFFINIS Bocourt, 1868, *Ann. Sci. Nat.*, 201.

Nasal divided; subcaudals simple; second labial not forming the anterior border of the lachrymal fosse; rostral triangular, apex rounded, not separated from the nasal by a row of scales; supraciliary long and narrow. Nine or ten labials, fifth largest, and separated from the orbit by four rows of scales. Back with twenty-three to twenty-five rhomboidal spots.

Resembling *Teleuraspis nummifer*, but distinguished by a more elongate form, the rostral in contact with the nasals, instead of being separate by a single row, the longitudinal development of the supraciliaries, and the greater number of rhomboidal spots. The spot beneath the eye is characteristic. Scales in 27 rows, Guatemala.

TELEURASPIS.

Cope, 1859.

Moderately stout; head large, distinct from the neck, depressed, triangular, broad behind, narrowed in front; muzzle prominent, angular; tail short, tapering to a point. Crown-shields small, scale-like. Eye moderate, pupil vertical. Supraciliary replaced by small scales or with a row at the orbital edge. Scales keeled. Anal entire. Subcaudals entire or bifid.

Subcaudals paired;

scales in 21 rows; an undulating band along the back

undulatus.

subcaudals entire;

no band behind the eye; with scattered spots of dark color

nigroadspersus.

a band from the eye; with a dorsal series of lozenge-shaped spots

nummifer.

TELEURASPIS UNDULATUS.

TRIGONOCEPHALUS (ATROPOS) UNDULATUS Jan, 1859, *Prodr. Icon. Ophid.*, 32.

Supraciliaries replaced by prominent scales, forming a sort of crest; one in particular is elongate, and constitutes an obtuse little horn. Crown covered by small carinate scales, a series of larger around the rostral angle. Labials twelve, none touching the pit; separated from the orbit by two scales. Infralabials thirteen. Submentals two pairs, posterior small; anterior in contact with three infralabials. Scales keeled, in 21 rows. Ventrals 177. Anal entire. Subcaudals 42 pairs.

Greyish, mixed with brown, an undulating brown band along the back. Flanks with small black spots. Belly greenish-yellow, with small black dots. (Jan.) Mexico.

TELEURASPIS NIGROADSPERSUS.

BOTHROPS (TELEURASPIS) NIGROADSPERSUS *Steindachner*, 1870, *Sitz. Akad. Wien.*, p. 348, *Taf. VIII.*

Head broad, depressed. Body compressed; tail short. Nasal divided, over the first labial. Two prominent shields at the outer edge of the large, smooth supraciliary. Rostral angles sharp. Keeled scales on crown and side of head; labials, pit-shields, nasals, and supraciliaries larger, smooth.

Labials nine, third largest, separated from the orbit by two scales, second forming a border to the pit. Scales keeled, in 23 rows. Ventrals 162. Anal entire. Subcaudals 55, entire.

Yellow, with scattered spots or dots on head and body; belly light yellow. Central America to Mexico.

TELEURASPIS NUMMIFER.

ATROPOS NUMMIFER *Ruppell*, 1845, *Museum Senckenbergianum*, 313.

BOTHIROPS NUMMIFER *Jan*, 1863, *Elenc. Sist.*, 126.

Stout; head short, triangular, broad behind; tail short, near one eighth of the total. Eye moderate, pupil erect. Crown covered with small carinate scales. Supraciliaries small. Scales of the side of the face small. Scales all keeled, in 25 rows, dorsal keels not reaching the end of the scale. Ventrals 132. Anal entire. Subcaudals 33, entire.

A dorsal series of rhomboidal darker-edged brown spots, sometimes complete and distinct, and sometimes triangular and alternating across the back, but united in such a manner as to form a zigzag series. A series of irregular-shaped black spots on the flank and another series alternating with these on the ends of the ventrals. Posteriorly the borders of the dorsal rhombs unite with each other, and the lateral spots inclosing lighter spaces on the flanks. Ventrals yellowish in front, becoming dark leaden posteriorly spotted with light. Labials, infralabials, chin-shields, and scales below the neck white. A dark band behind the eye, above the hinder labials to the neck. End of tail lighter. (Jan.) Guatemala, Mexico.

ADDITIONS AND CORRECTIONS.

It is three years since the foregoing Synopses and Descriptions were written. In that time additional researches and discoveries have necessitated a number of changes. These can not be made in the text. They appear in the Systematic List given below. The latter corrects the preceding wherever they differ, and expresses my views in regard to value of characters, classification, etc., at the latest date. Such descriptions as are lacking in the earlier work, and any changes in the text that may be found necessary, are placed under their proper heads in the List. In many cases the original diagnosis is somewhat closely followed, changes being made only when really necessary.

At first it was the intention to devote these pages exclusively to Kentucky species. The geographical position of this State, however, being such as to make it a sort of neutral ground, of which none of the species were peculiar to itself, it was seen that it would greatly enhance the value of the work for Kentucky, and for the other States, to increase its scope so as to take in all North America from the Isthmus of Tehuantepec. The descriptions include nearly or quite all the species occurring north of the Isthmus, although a few of them are only stragglers from the southward; the illustrations, with but one or two exceptions, are of those of Kentucky. Whenever possible, individuals showing variations from such as may have been figured or described by others have been selected for illustration, with the purpose of adding to or correcting our knowledge of the species.

NOVEMBER, 1882.

SYSTEMATIC LIST AND SYNONYMY.

OPHIDIA.	1
Brongniart, 1800, Bull. Soc. Philom.	
SCOLECOPHIDIA. (Worm Snakes.)	2
<i>Scotceophides</i> Dum. Bibr., 1814, Erp. Gén., VI, 243.	
TYPHLOPIDAE.	2
<i>Typhlopoides</i> Fitz., 1826, Nene Class., pp. 11, 25. <i>Typhlopsidae</i> Gray, 1840, Syn. Brit. Mus., — 1845, Cat. Lizards, 130.	
TYPHLOPINAE	
TYPHLOPS	2
Schneider, 1801, Hist. Amph., II, 339; D. & B., 1844, Erp., VI, 229;	
Gray, 1845, Cat. Lizards, 130; Jan, 1860, Arch. Zoöl., — 1864, Icon. Text, 7.	
TYPHLOPS LONGISSIMUS	2
<i>Typhlops (Ophthalmidion) longissimum</i> Jan, 1861, Arch. Zoöl., I, 182, — 1863, Icon. livr. 5, pl. 5, f. 6. <i>Ophthalmidion longissimum</i> D. & B., 1844, Erp., VI, 263; Dum., 1852, Cat. Meth., II; Baird & Girard, 1853, Serp., 155.	
"North America."	
TYPHLOPS PERDITUS	3
Peters, 1869, Monatsb. Berl. Akad., 435.	
Orizaba, Mexico.	
TYPHLOPS BASMACULATUS	3
Cope, 1866, Pr. Phil. Acad., 320.	
Cordova and Orizaba, Mexico.	
TYPHLOPS EMUNCTUS sp. n.	3
Central America.	
STENOSTOMINAE.	4
ANOMALEPIS	4
Jan, 1861, Arch. Zoöl., 185, — 1863, Sist., 9, — 1864, Icon. Text, 6, — 1857, Ind. Rett. Mus. Milan (Name).	
ANOMALEPIS MEXICANUS	4
Jan, 1861, Arch. Zoöl., 185, — 1861, Icon. livr. 3, pl. 5-6, f. 1, — 1863, Sist., 9, — 1864, Icon. Text, 7.	
Mexico.	

OPHIDIA—Continued.

- STENOSTOMA 4
- Wagler, 1824, Spix, Serp. Bras., 68; D. & B., 1844, Erp., VI, 322; Jan, 1861, Arch. per la Zool., 186. For Synopsis, page 5, read:
- Scales in 15 rows;
- ocular not in contact with the vertebral series;
- infralabials four *dulce*
- ocular in contact with vertebrals *humile*
- Scales in 14 rows;
- two labials, separated by the ocular;
- oculars not in contact with vertebrals;
- infralabials five *rubellum*
- oculars and vertebrals in contact; .
- infralabials five *tenuiculum*
- three labials, two in front of ocular;
- oculars and vertebrals not in contact *myopicum*
- Scales in 13 rows *phenops*
- STENOSTOMA DULCE 5
- Rena dulcis* B. & G., 1853, Serp., 142; Bd., 1859, P. R. R. Rep., X, pl. 33, f. 100. *Stenostoma dulce* Jan, 1861, Arch. Zool., 189, — Icon. livr. 2, pl. 5, 6, f. 5, — 1863, Sist., 15, — 1864, Icon. Text, 36; Cope, 1880, Bull. 20, U. S. Mus., p. 20, — 1875, Checklist, 44.
- Texas.
- STENOSTOMA HUMILE
- Rena humilis* B. & G., 1853, Serp., 143. *Stenostoma humile* Cope, 1861, Pr. Phil. Ac., 305 (Name), — 1875, Checklist, 44.
- Body very slender and cylindrical; tail short, conical, tapering, not acute, one-fifteenth of the total length. Head less depressed (than *S. dulcis*). Eyes and nostril more distinct than in the preceding species (*S. dulce*). The eye-shield in contact with the longitudinal series on top of the head. No supraorbitals. Post parietals much smaller than the parietals. Scales in 15 rows. Scales on the abdomen larger than on the back. Uniform chestnut-brown, lighter beneath.
- Vallicetas, Cal. (B. & G.)
- STENOSTOMA RUBELLUM sp. n.
- Long, slender, cylindrical; head indistinct, slightly depressed, muzzle rounded; tail short, thick, ending in a spine which is directed downward. Eyes distinctly visible beneath the ocular shields. Rostral reaching backward as far as to a vertical from the anterior edge of the ocular. Nasal obliquely divided; lower portion narrow, separated from the ocular by a single labial. Oculars reaching the edge of the lip, broadening upward, separated by three small shields on the top of the head, followed by a large labial. A pair of large broad parietals on each side behind the upper half of the ocular and the small supra-ocular, separated on the vertex by the vertebral series, and the posterior of each pair separated from the labial by a scale-like shield. Nasals separated by the extremity of the rostral and the anterior scale of the dorsal row. Rostral broad, upper extremity rather acute. Infralabials

OPHIDIA—Continued.

five. Scales in 14 rows; ventrals little, if any, larger. Each scale has four angles, and its posterior margin is nearly a semicircle. Back reddish brown; belly reddish white.

Resembles *S. dulce* in coloration, but differs in the number of dorsal rows of scales, complete separation of nasals by the rostral, five infra-labials, instead of four, and in having only the anterior parietal in contact with the posterior labial. Total length 8 inches; tail 0.38.

Uvalde, Texas.

STENOSTOMA TENUCULUM sp. n.
San Luis Potosi, Mexico.

STENOSTOMA MYOPICUM sp. n.
Tampico, Mexico.

STENOSTOMA PHENOPS
Cope, 1875, Jour. Phil. Ac., 128, — Checklist, 44.
Tehuantepec; Guatemala.

ONYCHOPHIDIA. (Clawed Snakes.) 7

ERYCIDAÆ. 7

Erycina Bonap., 1831, Saggio Distrib. Met. An. Vert. *Erycidae* Bonap., 1840, Roy. Acad. Turin, Mem., II; Gthr., 1864, Rept. Brit. Ind., 332.

CHARINA 7

Gray, 1849, Cat. Snakes, 113; B. & G., 1853, Serp., 154. *Wenona* B. & G., 1852, Pr. Phil. Ac., 176, — 1853, Serp., 139. *Pseudocorys* Jan, 1862, Arch. Natgesch., I, 242, — 1865, Icon., Text, 67. *Lichanura* Cope, 1861, Pr. Phil. Ac., 304.

CHARINA BOTTAE 7

Gray, 1849, Cat., 113; B. & G., 1853, Serp., 154. *Tortrix lottee* Blainv., 1835, Nouv. Ann. Mus., 289, pl. 26, f. 1-16. *Pseudocorys bottae* Jan, 1863, Sist., 21, — Icon. livr., 3, pl. 2, f. 1, — 1865, Icon., Text, 67.

California to Mexico.

Var. PLUMBEA

Wenona plumbea B. & G., 1852, Pr. Phil. Ac., 176, — 1853, Serp., 139; Grd., 1858, Wilkes' Exp., Rept., 112, pl. 7, f. 1-7; Jan, 1863, Sist., 21, — 1864, Icon., livr. 3, pl. 2, f. 2, — 1865, Icon., Text, 69. *Wenona isabella* B. & G., 1852, Pr. Phil. Ac., 176, — 1853, Serp., 140; Grd., 1858, Wilkes' Exp., Rept., 113, pl. 7, f. 8-14. *Charina plumbea* Cope, 1861, Pr. Phil., Ac., 305. *Wenona* Bd., 1859, P. R. R. Rep., X, pl. 32, f. 3.

Uniform brownish or leaden. Scales in 45 rows. Ventrals 26+37.
California to Puget Sound.

CHARINA TRIVIRGATA 8

Lichanura trivirgata Cope, 1861, Pr. Phil. Ac., 304; Jan, 1865, Icon., Text, 70. *L. roseofusca* Cope, 1863, Pr. Phil. Ac., 2. *L. myriolepis* Cope, 1868, Pr. Phil. Ac., 2.

Lower California; Mexico.

OPHIDIA—Continued.

- BOAEIDAE 8
- Boidae* Bonap., 1831, Saggio, — 1840, Roy. Acad. Turin.
- BOA
- Linn., 1758, Systema, I, 214, — 1766, Syst., I, 373 (Part); Gray, 1825, Ann. Phil., 209; Wagler, 1830, Syst. Amph., 168; D. & B., 1844, Erp., VI, 500.
- BOA IMPERATOR 8
- Daudin, 1803, Rept., V, 150; D. & B., 1844, Erp., VI, 519; Gray, 1849, Cat. Snakes, 101; Jan, 1857, Ind. Sist. Rett. Mus. Milan, 44, — 1863, Sist., 23, — 1864, Icon., livr. 6, pl. 1, — 1865, Icon., Text, 81; Sumichrast, 1880, Bull. Soc. Zool. de France, 179.
- Central America and Mexico.
- BOA CONSTRICTOR
- Bons constrictor* Linn., 1754, Mus. Ad. Fridr., 38. *Boa constrictor* Linn., 1758, Syst., I, 373; Schneider, 1801, Amph., II, 247; Fitz., 1826, Neue Class., 54; Wagl., 1830, Syst., 163; D. & B., 1844, Erp., VI, 507; Gray, 1849, Cat. Sn., 100; Jan, 1864, Icon., livr. 5, pl. 2, f. 2.
- Brazil to Central America.
- Var. ISTHMICA 9
- Central America:
- BOA MEXICANA (Rapp). 10
- Boa dicinibopue* var. *mexicana* Jan, 1863, Sist., 23, — 1864, Icon., livr. 5, pl. 4, f. 1, — 1865, Icon., Text, 82.
- CHILABOTHIRUS
- Dum. Bibr., 1844, Erp., VI, 562.
- Body stout, slightly compressed; neck smaller; head broad, depressed; tail rather short. Nostrils lateral, anterior. Eyes small, pupil vertical. Anterior teeth larger. Labials imbricate, without pits. Crown-shields large, irregular. Scales smooth, flat. Subcaudals simple. Claws distinct.
- CHILABOTHIRUS INORNATUS
- D. & B., 1844, Erp., VI, 563; Jan, 1863, Sist., 24, — 1864, Icon., livr. 6, pl. 5, f. 1. *Boa inornata* Reinh., 1813, Dansk. Vid. Selsk., pl. 21-23.
- Body stout, slightly compressed; head distinct, crown flat, snout broad; tail short, prehensile, rather thick at the extremity. Rostral large, pentangular, wider at the suture with the labial and nasal. Nasal divided in three—upper portions largest and meeting between internasals and rostral, or in two—anterior part much larger and reaching from the anterior labial to the opposite nasal. Internasals and prefrontals large, subequal or subdivided. Frontal and supraciliaries large, the former often dissected. Parietals irregular, becoming scale-like posteriorly. Loreal large, elongate, sometimes supplemented by small shields. Antecoculars two, upper larger. Postoculars usually four. Temporals scale-like. Labials commonly eleven, the sixth touching the eye. Infralabials about thirteen, anterior four or five broad, anterior pair meeting behind the mental. Mental furrow deep. Submentals small. Scales smooth, flat, in 36 to 40 rows, outer and

OPHIDIA—Continued.

vertebral broader. Ventrals 266 to 271. Subcaudals 36 to 52, entire. Yellowish, reddish, or greyish brown, light to dark, punctulate and clouded with darker, with irregular transverse badly defined bands of darker, which are often confluent into more or less distinct longitudinal bands on the sides of the neck. Belly olivaceous, flecked with lighter. Tail sometimes transversely banded. Some specimens are nearly uniform in coloration, very light to almost uniform black. The specimens described were large ones from Porto Rico. Reinhardt's specimens had 264 to 271 ventrals, and 67 to 69 subcaudals. The tail is shorter in the adult, and commonly the tip is imperfect.

West Indies; (?) Central America; (?) Mexico.

ACACOPHIDIA. (Non-venomous Snakes.) 11

COLUBRIDAE. 11

Bonap., 1831, Saggio, etc.

DIPSADINAE 11

LEPTOGNATHUS 12

Dum., 1852, Mem. Acad., 23, p. 467; D. & B., 1854, Erp., VII, 473; Gthr., 1858, Cat., 177.

LEPTOGNATHUS FASCIATUS 12

Tropidodipsas fasciata Gthr., 1858, Cat., 181. *Leptognathus philippii* Jan., 1863, Syst., 101, — 1870, Icon, livr. 37, pl. 5, f. 1. *L. fasciata* Cope, 1868, Pr. Phil. Ac., 137.

Mexico; Central America.

LEPTOGNATHUS DUMERILII (?). 13

Jan., 1863, Syst., 101, — 1870, Icon, livr. 37, pl. 5, f. 2.

Mexico.

LEPTOGNATHUS NEBULATUS 13

Gthr., 1858, Cat., 177; Cope, 1868, Pr. Phil. Ac., 136.

Coluber nebulatus Linn., 1754, Mus. Ad. Fridr., 32, pl. 24, f. 1, — 1758, Syst., I, 222, — 1766, Syst., I, 383; Weigel, 1782, Abb. Hall. Nat. Ges., I, 32; Mus. Linck, 1783, I, 75; Gmel., 1788, Syst. Linn., I, 1107; Shaw, 1802, Zool., III, 476; Latr., 1802, Rept., IV, 164; Daud., 1803, Rept., VI, 435; Merr., 1820, Syst., 104. *Cerastes nebulatus* Laur., 1768, Synops., 83. *Sibon nebulatus* Fitz., 1826, Nene Class., 60. *Dipsas nebulata* Boie, 1827, Isis, XX, 550; Schl., 1837, Essai, I, 162, II, 275, pl. 11, f. 14, 15; DeFil., 1840, Cat. Serp. *Petalognathus nebulatus* Dum., 1852, Mem. Acad., XXIII, 466; D. & B., 1854, Erp., VII, 464; Martens, 1856, Mus. Berl., 27.

Scha., 1734, Thesaur., I, 22, pl. 14, f. 4, — 1735, pl. 29, f. 3; Scheuch., 1735, Phys. Saer., IV, 1532, pl. 748, f. 8; Linn., 1749, Anoen. Acad., I, 304; Daub., 1771, Diet. Anim., 657, 679; LaC., 1789, Quad. Ovip. & Serp., II, 271, 307; Bonnat., 1789, Ophiol. 36, pl. 20, f. 38; Merr., 1790, Beitr., I, 31, pl. 8; Bechst., 1802, Ueb. LaC. Nat., IV, 39, 96, pl. 4, f. 2.

Brazil to Mexico; West Indies (Gthr.)

OPHIDIA—Continued.

- LEPTO NATHUS IMMIDIATUS 14
 Guenther, 1872, Ann. Mag. Nat. Hist., IX, 31. Mexico.
- LEPTOGNATHUS BREVIS (?)
 Dum., 1852, Mem. Acad., XXIII, 467; D. & B., 1854, Erp., VII, 476;
 Cope, 1868, Pr. Phil. Acad., 136. Mexico.
- DIPSAS 14
 Laur., 1768, Syst. Rept., 89; Gthr., 1858, Cat. Col. Snakes, 169.
 (HIMANTODES.)
- DIPSAS CENCHOA 14
 Max., 1825, Beitr. Brazils, I, 396; Fitz., 1826, Neue Class., 59; Boie,
 1827, Isis, 560; Wagl., 1830, Amph., 181; Gthr., 1858, Col. Snakes,
 174. *Coluber cenchoa* Linn., 1758, Syst. Nat., I, 226, — 1766, Syst.
 Nat., I, 389; Gmelin, 1788, Syst. Nat. Linn., I, 1118; Shaw, 1802,
 Gen. Zool., III, 475; Laur., 1802, Hist. Rept., IV, 129; Daud., 1803,
 Rept., VI, 283; Kuhl, 1820, Beitr., 88. *Bungarus cenchoa* Oppel, 1810,
 Ann. Mus. Hist. Nat., XVI, 392, — 1811, Ord. Fam. Gatt. Rept., 70.
Dipsas weigeli Fitz., 1826, Neue Class., 59; Schlegel, 1837, Ess.
 Phys. Serp., I, 162, II, 278, pl. 11, f. 19-20; DeFil., 1840, Cat. Serp.;
 Fitz., 1843, Syst. Rept., 27. *Himantodes cenchoa* Dum. Bibr., 1854,
 Erp. Gen., VII, 1065; Mart., 1856, Mus. Berl., 31; Jan, 1863, Elenc.
 Sist., 102, — 1871, Icon., livr. 38, pl. 2, f. 1. *Natrix cenchoa* Merr.,
 1820, Amph., 114.
 Seba, 1735, Thesaur., II, Tab. 15, f. 2, 3; Scheuchser, 1735, Phys.
 Sacr., IV, Tab. 678, f. 3; Linn., 1749, Amoen. Acad., I, 306; Daub.,
 1771, Diet. Encycl., 691; Weigel, 1782, Berl. Nat. Ges., III; LaC.,
 1789, Quad. Ovip., II, 316; Bonn., 1789, Encycl. Meth., Oph., 35, pl.
 29, f. 60; Bechst., 1802, Ueb. LaC. Natg., IV, 115.
 Mexico to Brazil.
- DIPSAS LITCOMELAS 15
 Cope, 1861, Pr. Phil. Ac., 296. Vera Cruz, Mexico.
- SIBON 15
 Fitz., 1826, Neue Class., 60; Cope, 1860, Pr. Phil. Ac., 266.
- SIBON BISCIUTATUM 16
Dipsas biscutata D. & B., 1854, Erp. Gen., VII, 1153.
Dipsatomorphus biscutatus Gthr., 1858, Col. Serp., 176. *Eleiroidipsas biscu-*
tata Jan, 1863, Sist., 105, — 1872, Icon., livr. 39, pl. 1, f. 3. *Lycoplon*
lyrophaneus Cope, 1860, Pr. Ac. Phil., 343. *Trimorphodon biscutatus*
 Cope, 1869, Pr. Am. Phil. Soc., 152. *Trimorphodon lyrophaneus* Cope,
 1861, Pr. Phil. Ac., 297. Central America to Mexico.
- Var. LATIFASCIATUM 17
Dipsas biscutata var. *latifascia* Peters, 1869, M. B. Berl. Acad., 877.
 Pueblo, Mexico.
- SIBON UPSILON
Trimorphodon upsilon Cope, 1879, Pr. Am. Phil. Soc., 152.
 Locals 3. Oculars 3-3. Temporals 3+3+3. Blotches forming
 transverse bands. Guadalajara, Western Mexico.

OPHIDIA—Continued.

- SIBON ANNULATUM
- Fitz., 1826, Neue Class., 60; Cope, 1860, Pr. Phil. Ac., 266.
- Calaber annulatus* Linn., 1754, Mus. Ad. Fridr., pl. 8, f. 2, — 1758, Syst., I, 224, — 1766, Syst., I, 386; Mus. Linck, I, No. 76; Gmelin, 1788, Syst. Linn., I, 1111; Shaw, 1802, Gen. Zool., III, 490; Latr., 1802, Rept., IV, 127; Daud., 1803, Rept., VI, 369; Kuhl, 1820, Beitr., 85; Merr., 1820, Amph., 111, — 1821, Beitr., fasc. 3, p. 25, pl. 3-4. *Dipsas annulata* Schleg., 1837, Ess. Phys., I, 165, II, 294; DeFil., 1840, Cat. Serp.; D. & B., 1854, Erp. Gen., VII, 1141; Martens, 1856, Mus. Berl., 32. *Itycodon annulatus* Boie, 1827, Isis, XX, 527. *Leptodécira annulata* Fitz., 1843, Syst., 27; Gthr., 1858, Cat., 166. *Eteirodipsas annulata* Jan, 1863, Syst., 165.
- Seba, 1735, II, pl. 41, f. 3, pl. 57, f. 1, pl. 71, f. 3; Scheuchser, 1735, Phys. Sacr., IV, pl. 652, f. 2; Knorr, 1767, Delic. Surin. Nat., II, pl. 6, f. 2; Daub., 1771, Diet. Encycl., 591; LaC., 1789, Quad. Ovip., II, 312; Bechst., 1802, Ueb. LaC., IV, 107, pl. 13, f. 1; Merr., 1790, Beitr., pt. I, pl. 11; Herm., 1804, Obs. Zool., I, 285.
- Mexico to Brazil.
- Var. SEPTENTRIONALE 17
- Dipsas septentrionalis* Kenn., 1859, Mex. Bound. Surv., II, 16, pl. VIII, f. 1; Bd., 1859, P. R. R. Rep., X, pl. 25, f. 18. *Sibon septentrionalis* Cope, 1860, Pr. Phil. Ac., 226. *Sibon annulatum* subsp. *septentrionale* Cope, 1875, Checklist, 38. *Eteirodipsas annulata* var. *septentrionalis* Jan, 1872, Icon., livr. 39, pl. 1, f. 2.
- Texas; Arizona; Mexico.
- SIBON TORQUATUM 18
- Leptodécira torquata* Gthr., 1860, Ann. Mag. *Hipsiglena torquata* Cope, 1861, Pr. Phil. Ac., 302. Nicaragua.
- Var. PACIFICUM
- Leptodécira pacifica* Cope, 1868, Pr. Phil. Ac., 310.
- Oculars 2—2. Loreal higher than long. Labials 7, third and fourth in orbit. Temporals 1+2. Scales in 19 rows, biporous, all nearly equal. Ventrals 164. Anal bifid. Subcaudals 61 pairs. Back brownish, with four series of blackish brown very small spots, of which the median pair forms usually a short cross-bar by their union. A broad black bar across the nape; a short brown line behind each eye, and a number of dots on the crown. Below uniform cream color.
- Mazatlan, Mexico.
- Var. PERSONATUM
- Leptodécira personata* Cope, 1868, Pr. Phil. Ac., 310.
- Oculars 1—2. Loreal as high as long. Labials 8, fourth and fifth in orbit. Temporals 1+2. Scales in 23 rows, biporous. Ventrals 173; anal divided. Cream color, with broad black cross bands on the dorsal region, 23 between nape and vent. Top of head black. A broad yellowish collar.
- Mazatlan, Mexico.
- SIBON DISCOLOR 18
- Leptodécira discolor* Gthr., 1860, Pr. Zool. Soc. Lond., 317.
- Oaxaca, Mexico.

OPHIDIA—Continued.

- SCYTALINAE 18
Scytalius D. & B., 1854, *Erp. Gen.*, VII, 988. *Scytalidae* Gthr., 1858, *Cat. Col.*, 185.
- OXYRHOPUS 18
 Wagler, 1830, *Amph.*, 185.
- OXYRHOPUS CLOELIA 19
 Guenther, 1858, *Cat. Col. Sn.*, 189. *Coluber cloelia* Daud., 1803, *Rept.*, VI, 330, pl. 78. *Natrix cloelia* Merr., 1820, *Amph.*, 98. *Cloelia daudini* Fitz., 1826, *Neue Class.*, 55, No. 2. *Cloelia occipitalis* Wagler, 1830, *Amph.*, 187. *Ignoton cloelia* Schl., 1837, *Ess. Phys.*, I, 143, II, 114. *Deiopeda cloelia* Fitz., 1843, *Syst.*, 25. *Brachyrylon cloelia* D. & B., 1854, *Erp. Gen.*, VII, 1007. Central America; Mexico (?).
- OXYRHOPUS DOLIATUS 19
 Dum. Bibr., 1854, *Erp. Gen.*, VII, 1020; Gthr., 1858, *Cat.*, 192. Central America to Brazil; Mexico (?).
- DENDROPHINAE 20
Dendrophilae Gthr., 1858, *Cat.*, 142.
- LEPTOPHIS 20
 Bell, 1825, *Zool. Jour.*, 329.
- LEPTOPHIS MEXICANUS 20
 Dum. Bibr., 1854, *Erp. Gen.*, VII, 536; Jan, 1863, *Sist.*, 84, — 1878, *Icon.*, livr. 49, pl. 6, f. 4. *Ahoatulla mexicana* Gthr., 1858, *Cat.*, 154. *Thrasops mexicanus* Cope, 1860, *Pr. Phil. Ac.*, 557. Mexico.
- OXYBELIS 21
 Wagler, 1830, *Amph.*, 183.
- OXYBELIS AENEUS 21
 Wagl., 1830, *Amph.*, 183; Fitz., 1843, *Syst.*, 27. D. & B., 1854, *Erp. Gen.*, VII, 819; Martens, 1856, *Mus. Berl.*, 31. *Dryinus aeneus* Wagl., 1824, *Spix Serp. Bras.*, 12, pl. III; Boie, 1827, *Isis*, 546. *Dryiophis aeneus* Cuv., 1829, *Regn. An.*, II, 83. *Coluber acuminatus* Wied., 1830, *Abbild. Nat. Bras.*, Lief. 14, pl. I, — 1825, *Beitr.*, 322. *Dryiophis aurata* Schleg., 1837, *Essai*, pl. 10, f. 16–18. *Dryinus auratus* Bell, 1825, *Zool. Jour.*, II, pl. 12. *Dryiophis acuminata* Gthr., 1858, *Cat.*, 156. Mexico to Brazil.
- NATRICINAE 21
Natricidae Gthr., 1858, *Cat.*, 50.
- TROPIDONOTUS 22
 Kuhl, 1826, *Isis*, 205; Boie, 1827, *Isis*, 518; Wagl., 1830, *Amph.*, 179; Schleg., 1837, *Ess.*, 296; D. & B., 1854, *Erp. Gen.*, VII, 594; Gthr., 1858, *Cat.*, 59. (CHILOPOMA.)
- TROPIDONOTUS RUFIPUNCTATUS 23
Chilopoma rufipunctatum (Cope) Yarrow, 1875, *Wheeler's Surv.*, V, 543, pl. XX, f. 1; Cope, 1875, *Checklist*, 40. Southern Arizona.

OPHIDIA—Continued.

(EUTAENIA.)

TROPIDONOTUS SAURITA (Ribbon Snake) 23

- Schlegel, 1837, Essai, I, 169, II, 327; D. & B., 1854, Erp., VII, 585; Mart., 1856, Mus. Berl., 25; Gthr., 1858, Cat., 72; Jan, 1863, Sist., 69, — 1865, Arch. Zool., 212, — 1868, Icon., livr. 26, pl. 2, f. 1; Allen, 1868, Pr. B. N. H. Soc., 180. *Coluber saurita* Linn., 1766, Syst., I, 385; Gmel, 1788, Syst. Linn., I, 1109; Harl., 1827, Jour. Phil. Ac., V, 352, — 1835, Med. Res., 115; Shaw, 1802, Zool., III, 532; Fitz., 1826, Neue Class., 59; Daud., 1803, Rept., VII, 81; Bonnat., 1789, Ophiol., 58; Storer, 1839, Mass. Rept., 229; Thomps., 1842, Hist. Vermont, 115. *Leptophis sauritus* Holbr., 1842, Herp., IV, 21, pl. 4; DeK., 1842, N. Y. Fauna, Rept., 47, pl. 11, f. 24. *Eutaenia saurita* B. & G., 1853, Serp., 24; Bd., 1854, Serp. N. Y., 14, — 1859, P. R. R. Rep., X, pl. 26, f. 19; Yarrow, 1875, Wheeler's Rep., V, pp. 545, 546. *Ribbon Snake* Catesby, 1743, Carol., II, 50.

Mississippi Valley to the Atlantic.

Var. SACKENII

- Eutaenia sackenii* Kenn., 1859, Pr. Phil. Ac., 98; Yarr., 1875, Wheeler's Rep., V, pp. 545, 547; Cope, 1877, Pr. Am. Phil. Soc., 64.

Slender. Back olive black, to edge of abdomen. Lateral stripe greenish yellow, on third and fourth rows. No dorsal stripe. Belly greenish.

Florida.

Var. FAIREYI

- Eutaenia faireyi* B. & G., 1853, Serp., 25; Bd., P. R. R. Rep., X, pl. 26, f. 20; Yarr., 1875, Wheeler's Rep., V, pp. 545, 547; Cope, 1880, Bull. 20, U. S. Mus., 23. *Tropidonotus saurita* var. *faireyi* Jan, 1863, Sist., 71; Cooper, 1860, P. R. R. Rep., XII (2), 299, pl. 13, 14.

Stouter than *T. saurita*. Scales in 19 rows. Ventrals 168 to 178. Subcaudals 112—115. Back to edge of abdomen blackish brown. With three longitudinal stripes, outer on third and fourth rows.

Minnesota and Wisconsin Southward.

Var. PROXIMUS

- Coluber proximus* Say, 1823, Long's Exp., I, 187. *Tropidonotus sauritus* var. *proximus* Jan, 1863, Sist., 70. *Eutaenia proxima* B. & G., 1853, Serp., 25, — 1854, Marey's Exp., 191, pl. 2; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 21; Yarr., 1875, Wheeler's Rep., V, pp. 545, 547; Cope, 1880, Bull. 20, U. S. Mus., 23; Sumichrast, 1881, Bull. Soc. Zool. de France, 182; Cones & Yarrow, 1878, Hayd. Rep., IV, 280. *E. macrostemma* Kenn., 1860, Pr. Phil. Ac., 331; Cope, 1866, Pr. Phil. Ac., 307; Yarr., 1875, Wheeler's Rep., V, 548. *E. megalops* Kenn., 1860, Pr. Phil. Ac., 330; Yarr., 1875, Wheeler's Rep., V, pp. 545, 547. *E. flavilabris* Cope, 1866, Pr. Phil. Ac., 306; Yarr., 1875, Wheeler's Rep., V, 547.

Arkansas and Texas to Mexico.

OPHIDIA—Continued.

- TROPIDONOTUS SIRTALIS** (Garter Snake) 24
Coluber sirtalis Linn., 1758, Syst., I, 222, — 1766, Syst., I, 383; Gmel., 1788, Syst. Linn., I, 1107; Shaw, 1802, Zool., III, 535; Merr., 1820, Syst., 132; Harl., 1827, Jour. Phil. Ac., 352; Storer, 1839, Mass. Rept., 221. *Eutaenia sirtalis* B. & G., 1853, Serp., 50; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 23; Yarr., 1875, Wheeler's Rep., V, 550; Cope, 1877, Pr. Am. Phil. Soc., 64. *Tropidonotus sirtalis* Holbr., 1842; Herp., IV, 41, pl. 11; Jan, 1863, Sist., 69, — 1865, Arch. Zool., 210. *T. ordinatus* var. B., Gthr., 1858, Cat., 74. *T. bipunctatus* Schleg., 1837, Essai, I, 168, II, 320; D. & B., 1854, Erp., VII, 582. *T. konnicotti* Jan, 1863, Sist., 70, — 1865, Arch. Zool., 216. *T. sirtalis* var. *dorsalis* Jan, 1863, Sist., 69. *Eutaenia dorsalis* B. & G., 1853, Serp., 31; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 2; Yarr., 1875, Wheeler's Rep., V, 550. *E. sirtalis* subsp. *dorsalis* Yarr., 1875, Wheeler's Rep., V, 554. *E. sirtalis* var. *obscura* Cope, 1880, Bull. 20, U. S. Mus., 23. *Tropidonotus sirtalis* var. *dorsalis* Jan, 1867, Icon., livr. 25, pl. 4, f. 1. *E. cyrtopsis* subsp. *ocillata* Cope, 1880, Bull. 20, U. S. Mus., 22. New England to the Mississippi Valley.
- Var. ORDINATUS** 25
Coluber ordinatus Linn., 1766, Syst., I, 379; Gmel., 1788, Syst. Linn., I, 1097; Harl., 1827, Jour. Phil. Ac., V, 349. *Tropidonotus ordinatus*, Holbr., 1842, Herp., IV, 45, pl. 12; Gthr., 1858, Cat., 73. *Eutaenia ordinata* B. & G., 1853, Serp., 32; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 24; Yarr., 1875, Wheeler's Rep., V, 551. *Spotted Ribbon Snake* Catesby, 1743, Carol., II, pl. 51 and 53. Coast Region, Nova Scotia to Georgia; Alabama.
- Var. RADIX**
Tropidonotus sirtalis var. *radix* Jan, 1863, Sist., 69, — 1865, Arch. Zool., 211. *Eutaenia radix* B. & G., 1853, Serp., 34; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 25, pl. 34, f. 5; Cooper, 1860, P. R. R. Rep., XII, (2), 299; Yarrow, 1875, Wheeler's Rep., V, 548; Coues & Yarrow, 1878, Hayden's Rep., IV, 277. *E. radix triningii* Coues & Yarrow, 1878, Hayd. Rep., IV, 279. *Thamnophis haydeni* Cope, 1862, Hayd., Trans. Am. Assoc., XII, 177. *Eutaenia haydeni* Kenn., 1860, Coop. P. R. R. Rep., XII (2), 298, pl. 14; Jan, 1865, Arch. Zool., 212. Scales in 19 rows, outer broad. Black, with three narrow yellow lines, or dark brown, with the lines and with series of irregular spots on the flanks. Lateral yellow lines on the third rows. Ventrals 153. Subcaudals 51 pairs. Northern part of Mississippi basin, Michigan to Rocky Mountains.
- Var. MARCIANA** 25
Eutaenia marciانا B. & G., 1853, Serp., pp. 36, 156, — 1854, Marcy's Exp., 192, pl. 3; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 26; Yarrow, 1875, Wheeler's Rep., V, pp. 549, 555; Cope, 1880, Bull. 20, U. S. Mus., 22; Sumichrast, 1881, Bull. Soc. Zool. Fr., 182. *Tropidonotus marciانا* Jan, 1863, Sist., 70, — 1865, Arch. Zool., 215. (?) *T. sirtalis* Peters, 1869, Mb. Berl. Akad., 877. Kansas to Texas and Mexico.

OPHIDIA—Continued.

- Var. **PARIETALIS**. 25
Coluber parietalis Say, 1823, Long's Exp., I, 186. *Tropidonotus parietalis* Hall., 1856, Pr. Phil. Ac., 248. *Tropidonotus sietalis* var. *parietalis* Jan, 1863, Sist., 69. *Eutaenia sietalis parietalis* Coles & Yarrow, 1878, Hayd. Rep., IV, 276. *Eutaenia parietalis* B. & G., 1853, Serp., 28. *E. ornata* Bd., 1859, P. R. R. Rep., X, pl. 26, f. 22; Cope, 1866, Pr. Phil. Ac., 305, 306. — 1875, Checklist, 41; Yarr., 1875, Wheel. Rep., V, 550, 553.
 Missouri basin and westward to Utah.
- Var. **VAGRANS**. 25
Eutaenia vagrans B. & G., 1853, Serp., 35; Grd., 1858, Expl. Exp. Herp., 154, pl. XIV, f. 5-10; Cooper, 1860, P. R. R. Rep., XII (2), 297; Cope, 1866, Pr. Phil. Ac., pp. 305, 307, vars.; Yarrow, 1875, Wheeler's Rep., V, 548; Coles & Yarrow, 1878, Hayd. Rep., IV, 274; Coles, 1875, Wheeler's Rep., V, 614; Baird, 1859, P. R. R. Rep., X, Rept., 19, pl. 17. *Eutaenia vagrans vagrans* Yarrow, 1875, Wheel. Rep., V, 551. *E. angustirostris* Kenn., 1860, Pr. Phil. Ac., 332; Yarrow, 1875, Wheel. Rep., V, 548. *E. vagrans* subsp. *angustirostris*, Yarr., 1875, Wheel. Rep., V, 554. *Tropidonotus vagrans* Jan, 1863, Sist., 70. — 1865, Arch. Zool., 215. *Eutaenia couchii* (Kenn.) Bd., 1859, P. R. R. Rep., X, Rept., 10; Yarr., 1875, Wheel. Rep., V, 549.
 Scales in 21 rows. Back brownish, with light vertebral line; belly slate color. Two series of small black spots, 95 to 105, on each side. Ventrals 161 to 179. Subcaudals 70 to 90.
 Rocky Mountains to the Sierras; Sonora.
- Var. **COLLARIS**. 25
Tropidonotus collaris Jan, 1863, Sist., 69. — 1867, Icon., livr. 25, pl. 5, 2.
 Southern Mexico; Panama.
- Var. **LEPTOCEPHALUS**. 25
Tropidonotus sietalis var. *leptocephala* Jan, 1863, Sist., 69. — 1867, Icon., livr. 25, pl. 4, f. 2. *Eutaenia leptocephala* B. & G., 1853, Serp., 29; Bd., 1859, P. R. R. Rep., X, pl. 36, f. 2; Girard, 1858, Expl. Exp., Herp., 151, pl. XII, f. 7-13; Coop., 1860, P. R. R. Rep., XII (2), 297; Jan, 1865, Arch. Zool., 211; Yarr., 1875, Wheel. Rep., V, 550.
 Scales in 19 to 17 rows, outer smooth. Orbitals 2-3. Light olive brown, with two series (about 130) of small black spots on each flank. Lateral stripes obsolete. Scales black at base. Ventrals 140 to 150. Subcaudals 57 to 66.
 Oregon
- Var. **INFERNALIS**. 25
Coluber infernalis Blainv., 1834, Nouv. Ann. Mus., III, 59, pl. XXVI, f. 3. *Eutaenia infernalis* B. & G., 1853, Serp., 26; Grd., 1858, Expl. Exp., Herp., 148, pl. XIV, f. 11-16; Bd., 1859, P. R. R. Rep., X, Rept., 10; Yarr., 1875, Wheel. Rep., V, 550. *Tropidonotus saurita* var. *infernalis* Jan, 1863, Sist., 70. *Eutaenia pickeringii* B. & G., 1853, Serp., 27; Grd., 1858, Expl. Exp., 150, pl. XIII, f. 14-20; Bd.,

OPHIDIA—Continued.

1859, P. R. R. Rep., X, pl. 36, f. 3; Coop., 1860, P. R. R. Rep., XII (2), 296; Yarr., 1875, Wheel. Rep., V, 550; Coues & Yarr., 1875, Hayd. Rep., IV, 281. *Trop. sirtalis* var. Jan, 1863, Sist., 69, — 1865, Arch. Zool., 211. *Eutaenia cooperi* Kenn., 1859, Pr. Phil. Ac.; Coop., 1860, P. R. R. Rep., XII (2), 296; Yarr., 1875, Wheel. Rep., V, 551. *Trop. sirtalis* var. *cooperi* Jan, 1863, Sist., 69, — 1865, Arch. Zool., 212. *Eutaenia elegans* B. & G., 1853, Serp., 34; (Kenn.) Bd., 1859, P. R. R. Rep., X, Rept., 10; Yarr., 1875, Wheel. Rep., V, 549. *Trop. elegans* Jan, 1863, Sist., 70, — 1865, Arch. Zool., 214. *E. cyrtopsis* Kenn., 1860, Pr. Phil. Ac., 333; Cope, 1866, Pr. Phil. Ac., 316; Yarr., 1875, Wheel. Rep., V, 550; Cope, 1880, Bull., 20, U. S. Mus., 23. *Thamnophis cyrtopsis* var. *eylides* Cope, 1860, Pr. Phil. Ac., 299. *E. ordinoides* B. & G., 1853, Serp., 33; Grd., 1858, Expl. Exp., Ierp., 153, pl. 14, f. 1-4; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 3; Yarr., 1875, Wheel. Rep., V, 548. *Tropidonotus ordinoides* B. & G., 1852, Pr. Phil. Ac., 176. *T. sirtalis* var. *ordinoides* Jan, 1863, Sist., 69, — 1865, Arch. Zool., 211. *Trop. concinnus* Hall., 1852, Pr. Phil. Ac., 182. *Eut. concinna* B. & G., 1853, Serp., 146; Bd., 1859, P. R. R. Rep., X, Rept., 10; Coop., 1860, P. R. R. Rep., XII (2), 298; Yarr., 1875, Wheel. Rep., V, 549. *E. hammondi* Kenn., 1860, Pr. Phil. Ac., 332; Yarr., 1875, Wheel. Rep., V, 549. *Trop. hammondi* Jan, 1863, Sist., 70, — 1865, Arch. Zool., 215. *Eutaenia atrata* (Kenn.) Coop., 1860, P. R. R. Rep., XII (2), 296; Yarr., 1875, Wheel. Rep., V, 551. *Trop. sauritus* var. Jan, 1863, Sist., 70, — 1865, Arch. Zool. *Eut. sanichrosti* Cope, 1866, Pr. Phil. Ac., 306; Yarr., 1875, Wheel. Rep., V, 547. *Thamnophis scalaris* Cope, 1860, Pr. Phil. Ac., 369. *Eutaenia scalaris* Cope, 1866, Pr. Phil. Ac., 306; Yarr., 1875, Wheel. Rep., V, 549. *Eutaenia phenax* Cope, 1868, Pr. Ac. Phil., 134; Yarr., 1875, Wheel. Rep., V, 549. (?) *Trop. scalyer* Jan, 1863, Sist., 70. (?) *Trop. glyphyrus* Jan, 1863, Sist., 70. *T. ruficeps* Peters, 1869, Mb. Bd. Akad., 877.

California to Mexico.

Var. QUADRISERIALIS.

Tropidonotus quadriserialis Fischer, 1879, Verh. Nat. Ver. Hamb., 82.

Labials 8. Infralabials 10. Orbitals 1-3. Scales in 19 rows, keeled. Ventrals 149. Anal bifid. Subcaudals 79 pairs. Olivaceous, with four rows of small black spots.

Mazatlan, Mexico.

(NERODIA.)

TROPIDONOTUS SIPEDON (Water Snake) 25

Cularis sipedon Linn., 1758, Syst., I, 219, — 1766, Syst., I, 379; Gmel., 1788, Syst. Nat. Linn., I, 1098; Shaw, 1802, Zool., III, 496; Merr., 1820, Syst., 124; Harl., 1827, Jour. Phil. Ac., 351, — 1835, Med. Res., 114; Storer, 1839, Rept. Mass., 228; Thompson, 1842, Hist. Vermont, 118. *Tropidonotus sipedon* Holbr., 1842, Ierp., IV, 29, pl. VI; Dum. Bibr., 1854, Erp., VII, 568. *Nerodia sipedon* B. & G., 1853, Serp., 38 Baird, 1854, N. Y. Serp., 16, — 1859, P. R. R. Rep., X, pl. 27, f. 27.

OPHIDIA—Continued.

Tropidonotus fasciatus var. Jan, 1863, Sist., 71; *Trop.* (*Nerodia*) *sipedon* Jan, 1865, Arch. Zool., 222. *Coluber fasciatus* Linn., 1766, Syst., I, 378; Gmel., 1788, Syst. Nat. Linn., I, 1094; Holbr., 1838, Herp., I, 93, pl. 20. *Nerodia fasciata* B. & G., 1853, Serp., 19; Bd., 1859, P. R. R. Rep., X, pl. 34, f. 4. *Tropidonotus fasciatus* Schleg., 1837, Essai, I, 169, II, 323, pl. 12, f. 16, 17; Holbr., 1842, Herp., IV, 25, pl. V; D. & B., 1854, Erp., VII, 566; Gthr., 1858, Cat., 76; Jan, 1863, Sist., 71, — 1868, Icon., livr. 26, pl. 3, f. 2; Cope, 1877, Pr. Am. Phil. Soc., 64, — 1880, Bull. 20, U. S. Mus., 22; Mart., 1856, Mus. Berl., 24. *Trop. niger* Holbr., 1842, Herp., IV, p. 37, pl. 9; D. & B., 1854, Erp., VII, 572. *Nerodia niger* B. & G., 1853, Serp., 147; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 31. *Coluber porcatus* Harl., 1827, Jour. Phil. Ac., 356. *Nerodia couchii* Kenn., 1860, Pr. Phil. Ac., 335. *Tropidonotus couchii* Cope, 1860, Pr. Ac. Phil., 342. *Nerodia agassizii* B. & G., 1853, Serp., 41. *Tropidonotus obliquus* Hall., 1856, Pr. Phil. Ac., 248. *Tropidonotus rhombifer* Jan, 1863, Sist., 71, — 1868, Icon., livr. 26, pl. 4, f. 2. *Coluber pavidogaster* Max., 1839, Reise Nord. Amer., 106. *Tropidonotus sipedon woodhousei* Cope, 1880, Bull. 20, U. S. Mus., 22.

Mississippi Valley to Maine.

Var. ERYTHROGASTER 26

Coluber erythrogaster Holbr., 1838, Herp., II, 91, pl. 19. *Tropidonotus erythrogaster* Holbr., 1842, Herp., IV, 33, pl. VII; D. & B., 1854, Erp., VII, 570. *Nerodia erythrogaster* B. & G., 1853, Serp., 40; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 28. Copper-belly Snake, Catesby, 1743, Carol., II, p. 46, pl. 46.

Southeastern States.

Var. RHOMBIFER 26

Nerodia rhombifer B. & G., 1853, Serp., 147, 42; Bd., 1859, P. R. R. Rep., X, pl. 34, f. 2. *Tropidonotus rhombifer* 1852, Pr. Phil. Ac., 177; Cope, 1861, Pr. Phil. Ac., 298, — 1880, Bull. 20, U. S. Mus., 22. *Nerodia woodhousei* B. & G., 1853, Serp., 42; Bd., 1859, P. R. R. Rep., X, pl. 34, f. 3. *Tropidonotus woodhousei* Cope, 1860, Pr. Phil. Ac., 342; Jan, 1863, Sist., 71, — 1868, Icon., livr. 26, pl. 4, f. 1. *Nerodia hallowbrookii* B. & G., 1853, Serp., 43; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 30. *Tropidonotus transeversus* Hall., 1852, Pr. Phil. Ac., 177. *Nerodia transeversa* B. & G., 1853, Serp., 148; Bd., 1859, P. R. R. Rep., X, pl. 26, f. 1.

Mississippi Valley to Wisconsin.

TROPIDONOTUS TAXISPLOTUS 26

Hallowbrook, 1842, Herp., IV, 35, pl. 8; Dum. Bibr., 1854, Erp., VII, 603; Jan, 1863, Sist., 71, — 1868, Icon., livr. 26, pl. 5, f. 1. *Nerodia taxispilota* B. & G., 1853, Serp., 43; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 29. *Tropidonotus pogonius* D. & B., 1854, Erp., VII, 574; Gthr., 1858, Cat., 248. *Trop. fasciatus* var. *pog.* Jan, 1863, Sist., 71, — Icon., 1868, livr. 26, pl. 3, f. 1.

Southeastern States.

OPHIDIA—Continued.

- TROPIDONOTUS CYCLOPEUS 26
 Dum. Bibr., 1854, Erp., VII, 576; Gthr., 1858, Cat., 77, 248; Jan, 1863, Sist., 71, — 1868, Icon., livr. 26, pl. 5, f. 1 and 2; Mart., 1856, Mus. Berl., 24.
 Ohio to Florida.
- TROPIDONOTUS COMPRESSICAUDUS 27
Neotia compressicauda Kenn., 1860, Pr. Ac. Phil., 335. *Tropidonotus compressicauda* Cope, 1860, Pr. Phil. Ac., 368, — 1861, Pr. Phil. Ac., 74. *Tropidonotus ustus* Cope, 1860, Pr. Phil. Ac., 340.
 Florida.
- (REGINA.)
- TROPIDONOTUS LEBERIS 27
Coluber leberis Linn., 1758, Syst., I, 216, — 1766, Syst., I, 375; Gmelin, Syst. Linn., I, 1086; Shaw, 1804, Zool., III, 433. *Tropidonotus leberis* DeKay, 1842, N. Y. Fauna Rept., p. 45, pl. 11; Holbr., 1842, Herp., IV, 49, pl. 13; D. & B., 1854, Erp., VII, 579; Gthr., 1858, Cat., 78; Cope, 1860, Pr. Phil. Ac., 342; Jan, 1863, Sist., 72, — 1868, Icon., livr. 27, pl. 5, f. 1. *Regina leberis* B. & G., 1853, Serp., 45; Baird, 1854, N. Y. Serp., 17, — 1859, P. R. R. Rep., X, pl. 27, f. 32. *Coluber septentrionalis* Say, 1825, Jour. Phil. Ac., 240; Harl., 1827, Jour. Phil. Ac., 335, — 1835, Med. Res., 118.
 Michigan to Texas.
- VAR. RIGIDUS 28
Coluber rigidus Say, 1825, Jour. Phil. Ac., 239. *Tropidonotus rigidus* Holbr., 1842, Herp., IV, 39, pl. X; D. & B., 1854, Erp., VII, 577; Cope, 1860, Pr. Ac. Phil., 342. *Tropidonotus leberis* var. Jan, 1863, Sist., 72. *Regina rigida* B. & G., 1853, Serp., 46; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 33.
 New York southward.
- VAR. CLARKII
Regina clarkii B. & G., 1853, Serp., 48; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 35. *Tropidonotus clarkii* Cope, 1861, Pr. Phil. Ac., 74; Jan, 1863, Sist., 73, — 1868, Icon., livr. 27, pl. 6, f. 1. — 1880, Bull. 20, U. S. Mus., 22. *Tropidonotus medusa* Gthr., 1858, Cat., 78.
 Orbitals 1—3 to 2. Loreal longer than high. Labials 8, fifth and sixth largest. Scales in 19 rows. Ventrals 132. Anal entire. Subcaudals 57 pairs. Yellowish brown, with four longitudinal bands of darker. Belly yellowish, with two brownish black-dotted bands.
 Texas to Mexico.
- VAR. GRAHAMII 28
Regina grahamii B. & G., 1853, Serp., 47; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 34. *Tropidonotus grahamii* Gthr., 1858, Cat., 78. *Tropidonotus leberis* var. Jan, 1863, Sist., 72. *T. baronis mulleri* Trosch., 1865, in Muell Wirbelth., Mex., 79.
 Michigan to Texas.

OPHIDIA—Continued.

- VAR. VALIDUS 28
Regina calida Kenn., 1860, Pr. Phil. Ac., 334. *Tropidonotus validus*
 Cope, 1860, Pr. Phil. Ac., 342, — 1861, Pr. Phil. Ac., 298. *Trop.*
calaco Cope, 1860, Pr. Phil. Ac., 341. *Trop. tephroleuca* Cope, 1860,
 Pr. Ac. Phil., 341. (?) *Tropidonotus mesomelaus* Jan, 1863, Sist., 73,
 — 1868, Icon., livr. 27, pl. 5 and 6, f. 2.
 California to Mexico.
- TROPIDONOTUS KIRTLANDII 28
Regina kirtlandi Kenn., 1856, Pr. Phil. Ac., 95; Bd., 1859, P. R. R.
 Rep., X, pl. 27, f. 36. *Tropidoclonium kirtlandi* Cope, 1860, Pr. Ac.
 Phil., 340, — 1875, Checklist, 42. *Ischnognathus kirtlandi* Jan, 1863,
 Sist., 74, — 1868, Icon., livr. 30, pl. I, f. 1.
 Illinois to Ohio.
- STORERIA 29
 Baird and Girard, 1853, Cat. Serp., 135.
- STORERIA STORERIOIDES 29
Tropidoclonium storerioides Cope, 1865, Pr. Phil. Ac., 190, — 1875, Check-
 list, 42.
 Mexico.
- STORERIA OCCIPITOMACULATA 30
 Baird & Girard, 1853, Serp., 137; Baird, 1854, N. Y. Serp., 26, — Bd.,
 1859, P. R. R. Rep., X, pl. 33, f. 99; Cope, 1877, Pr. Am. Phil. Soc.,
 64, — 1875, Checklist, 42. *Ischnognathus occipitumaculatus* Gthr., 1858,
 Cat., 81; Jan, 1863, Sist., 74, — 1868, Icon., livr. 30, pl. I, f. 2. *Trop-*
idonotus occipitumaculatus Storer, 1839, Rept. Mass., 230. *Coluber caus-*
tus Hall., 1847, Pr. Phil. Ac., 274, — 1849, Pr. Phil. Ac., 245.
 Mississippi Valley and eastward.
- STORERIA DEKAYI 31
 Baird & Girard, 1853, Serp., 135; Baird, 1854, N. Y. Serp., 26, — Bd.,
 1859, P. R. R. Rep., pl. 33, f. 98; Cope, 1880, Bull. 20, U. S. Mus., 22.
Tropidonotus dekayi Holbr., 1842, Herp., IV, 53, pl. 14. *Trop. ordi-*
natus Storer, 1839, Rept. Mass., 223. *Ischnognathus dekayi* Dum.
 Bibr., 1854, Erp., VII, 507; Gthr., 1858, Cat., 81; Jan, 1863, Sist., 74,
 — 1868, Icon., livr. 30, pl. I, f. 3.
 Maine to Mexico.
- STORERIA COPEI 31
Adelphis copei Cope, 1879, Pr. Am. Phil. Soc., 265.
 Guadalaxara, Mexico.
- STORERIA LINEATA 32
Microps lineatus Hall., 1856, Pr. Phil. Ac., 241; Bd., 1859, P. R. R. Rep.,
 X, pl. 34, f. 6. *Tropidoclonium lineatum* Cope, 1860, Pr. Phil. Ac., 76,
 — 1875, Checklist, 42, — 1880, Bull., U. S. Mus., No. 20, p. 22.
 Texas to Kansas.
- HELICOPS 33
 Wagler, 1830, Syst. Amph., 170.
 Page 33, line 3, read: Teeth smooth, posterior maxillary larger and
 often isolated.

OPHIDIA—Continued.

- (TRETANORHINUS.)
- HELICOPS VARIABILIS 33
Tretanorhinus variabilis D. & B., 1854, *Erp. Gen.*, VII, 349; Jan, 1863, *Sist.*, 76. *Helicops agassizii* Jan, 1863, *Sist.*, 76, — 1868, *Icon.*, livr. 28, pl. 2, f. 1. *Tretanorhinus variabilis* var. *aduncus* Jan, 1863, *Sist.*, 76.
 Nicaragua to Mexico.
- HELICOPS NIGROLUTEUS
Tretanorhinus nigroluteus Cope, 1861, *Pr. Ac. Phil.*, 298.
 Head narrow, slightly distinct. Rostral broader than high, separated from internasals by the nasals. Nasal in two parts, nostril between. Prefrontals as long as frontal. Anterior border of frontal equal to lateral. Loricals two, anterior smaller. Oculars 2—2. Labials 8, fourth in orbit. Infralabials 10. Scales keeled, in 21 rows. Ventrals 136. Anal bifid. Black above; yellow beneath; punctulated anteriorly, punctulations forming bands near the flank.
 Central America.
- HELICOPS ALLENI 34
 Garman, 1874, *Pr. Bost. Soc.*, 92; Cope, 1875, *Checklist*, 43.
 Florida.
- HYDROPS 34
 Wagler, 1830, *Amph.*, 170; Gray, 1842, *Zoöl. Misc.*, 67; Fitz., 1843, *Syst.*, 25; Gray, 1849, *Cat. Serp.*, 75.
- HYDROPS ERYTHROGRAMMUS 35
Coluber erythrogrammus Latr., 1802, *Rept.*, IV, 141; Daud., 1803, *Rept.*, VII, 93, pl. 83, f. 2; Harl., 1827, *Jour. Phil. Ac.*, 361; Holbr., 1836, *Herp.*, I, 115, pl. 22. *Abaster erythrogrammus* B. & G., 1853, *Serp.*, 125; Bd., 1859, *P. R. R. Rep.*, X, pl. 33, f. 93; Gray, 1849, *Serp.*, 78; Gthr., 1858, *Cat.*, 275. *Calopisma erythrogrammus* D. & B., 1854, *Erp.*, VII, 336; Mart., 1856, *Mus. Berl.*, 27; Jan, 1863, *Sist.*, 75, — 1868, *Icon.*, livr. 29, pl. 4, f. 2, pl. 5, f. 1. *Homalopsis erythrogrammus* Boie, 1827, *Isis*, 551. *Helicops erythrogrammus* Wagl., 1830, *Amph.*, 170; Holbr., 1842, *Herp.*, III, 107, pl. 25. *Homalopsis plicatilis* var. Schleg., 1837, *Essai*, 173, II, 353. *Natrix erythrogrammus* Merr., 1820, *Amph.*, 117.
 Illinois and Virginia southward.
- HYDROPS ABACURUS 36
Coluber abacurus Holbr., 1836, *Herp.*, I, 119, pl. 23. *Hydrops abacurus* D. & B., 1854, *Erp.*, *Atlas*, pl. 65. *Calopisma abacurum* D. & B., 1854, *Erp.*, VII, 342; Mart., 1856, *Mus. Berl.*, 27. *Helicops abacurus* Holbr., 1842, *Herp.*, III, pl. 26. *Hydrops reinwardtii* Gray, 1842, *Zoöl. Misc.*, 67. *Homalopsis reinwardtii* Schleg., 1837, *Essai*, I, 173, II, 357. *Farancia fasciata* Gray, 1849, *Cat.*, 74; Gthr., 1858, *Cat.*, 275. *Farancia deammondi* Gray, 1842, *Zoöl. Misc.*, 68. *Farancia abacurus* B. & G., 1853, *Serp.*, 123; Bd., 1859, *P. R. R. Rep.*, X, pl. 33, f. 92. *Calopisma reinwardtii* Jan, 1863, *Sist.*, 75, — 1868, *Icon.*, livr. 29, pl. 6, f. 1-2.
 The Carolinas to Texas.

OPHIDIA—Continued.

HYDROPS QUINQUEVITTATUS 36
Homonopsis quinquevittatus D. & B., 1854, Erp., VII, 975. *Calopsisma quinquevittatum* Jan, 1863, Sist., 75, — 1865, Arch. Zool., 244, — 1868, Icon., livr. 30, pl. 2, f. 1. *C. quinquevittatum* var. *mevicana* Jan, 1863, Sist., 75, — 1868, Icon., livr. 30, pl. 2, f. 2.
 Mexico and Central America.

HYDROPS SEPTEMVITTATUS
Calopsisma septemvittatum Fischer, 1879, Verh. Nat. Vereins, Hamb., 84.
 Internasals fused. Nasal entire, grooved. Loreal present. Labials 8. Infralabials 9. Temporals 1+2+3. Scales in 19 rows, smooth. Ventrals 122 to 130. Anal bifid. Subcaudals 63 to 81 pairs (in description 53 to 73). Back chocolate brown, with four longitudinal black bands. Belly yellowish white, with three longitudinal black bands.

Mexico.

COLUBRINÆ 37
Colubridæ Gthr., 1858, Cat., 84.

SALVADORA 38
 Baird and Girard, 1853, Serp., 101. *Phimothya* Cope, 1860, Pr. Phil. Ac., 253 and 566.

SALVADORA GRAHAMII 38
 B. & G., 1853, Serp., pp. 104, 161; Bl., 1859, P. R. R. Rep., X, pl. 32, f. 78, — Mex. Bound. Surv., II, 21, pl. 5, f. 2; Jan, 1861, Icon., livr. 1, pl. 3, f. 1, — 1863, Sist., 58. *Phimothya grahamii* Cope, 1861, Pr. Phil. Ac., 300, — 1866, Pr. Phil. Ac., 310, — 1875, Checklist, 38; Yarr., 1875, Wheel. Rep., V, 538; Coues, 1875, Wheeler's Rep., V, 620.

California to Mexico and Utah to Texas.

VAR. BAIRDII
Salvadora bairdii Jan, 1861, Icon., livr. 1, pl. 3, f. 2, — 1863, Sist., 58. *Phimothya bairdii* Cope, 1861, Pr. Phil. Ac., 300; Sumichrast, 1873, Arch. Sci., 245; Pters., 1869, Mb. Brl. Akad., 876.
 Oculars 2—2 to 3. Loricals 2. Labials 8, sixth largest. Infralabials 10. Temporals 2+2+3. Rostral less produced backward, less prominent, and less free on the margins than that of *S. grahamii*.

Mexico.

VAR. HEXALEPIS 39
Phimothya hexalepis Cope, 1866, Pr. Phil. Ac., 305. *Phimothya grahamiae* subsp. *hexalepis* Cope, 1875, Checklist, 38. *Phimothya grahamiae hexalepis* Coues, 1875, Wheeler's Rep., V, 620.

Arizona.

SALVADORA DECURTATA 39
Phimothya decurtata Cope, 1868, Pr. Phil. Ac., 310, — 1875, Checklist, 38.

Lower California.

OPHIDIA—Continued.

- CYCLOPHIS 39
 Gthr., 1858, Cat. Serp., 119. *Chlorosoma* B. & G., 1853, Serp., 106.
- (CYCLOPHIS.)
- CYCLOPHIS VERNALIS (Green Snake) 39
 Gthr., 1858, Cat., 119; Cope, 1875, Checklist, 38; Yarr., 1875, Wheeler's Rep., V, 539; Cones & Yarrow, 1878, Hayden's Rep., IV, 285.
Coluber vernalis (DeK.) Harlan, 1827, Jour. Phil. Ac., 361, — 1835, Med. Res., 124; Storer, 1839, Rept. Mass., 224; Holbr., 1842, Herp., III, 79, pl. 17; Thomps., 1842, Hist. Vermont Rept., 117; DeKay, 1842, N. Y. Fauna, III, 40, pl. 11, f. 22; Jones, 1865, Rept. Nova Scotia, 6.
Liopeltis vernalis Cope, 1860, Pr. Phil. Ac., 560; Jan, 1863, Sist., 82, — Icon., livr. 31, pl. 5, f. 3; Hayd., 1862, Trans. Am. Phil. Soc., 177; Allen, 1869, Pr. Bost. Soc., 182.
Chlorosoma vernalis B. & G., 1853, Serp., 100; Bd., 1359, P. R. R. Rep., X, pl. 32, f. 81; Verr., 1863, Pr. Bost. Soc., 195. *Herpetodryas vernalis* Hall., 1856, Pr. Phil. Ac., 243.
 East of the Rocky Mountains to Nova Scotia.
- (PHYLLOPHILOPHIS.)
- CYCLOPHIS AESTIVUS (Green Snake) 40
 Gthr., 1858, Cat., 119; Cope, 1875, Checklist, 38, — 1880, Bull. 20, U. S. Mus., 23, — 1877, Pr. Phil. Am. Soc., 64.
Coluber aestivus Linn., 1766, Syst., I, 387; Gmel., 1788, Syst. Linn., I, 1114; Harl., 1827, Jour. Phil. Ac., 357, — 1835, Med. Res., 121. *Leptophis aestivus* Bell, 1826, Zool. Jour., II, 329; Holbr., 1842, Herp., IV, 17, pl. 3; B. & G., 1853, Serp., 106; Bd., 1859, P. R. R. Rep., X, pl. 32, f. 79. *Liopeltis aestivus* Jan, 1863, Sist., 81, — 1869, Icon., livr. 31, pl. 5, f. 1.
Ophiodryas aestivus Cope, 1860, Pr. Ac. Phil., 560. *Herpetodryas aestivus* Schleg., 1837, Essai, I, 151, II, 186, pl. 6, f. 12, 13; D. & B., 1854, Erp., VII, 209; Mart., 1856, Mus. Berl., 26. *Leptophis majalis* B. & G., 1853, Serp., 107, 161, — 1854, Marey's Exp., 203, pl. 9; Bd., P. R. R. Rep., X, pl. 32, f. 80.
The Green Snake Catesby, 1743, Carol., II, pl. 57; Bartram, 1791, Georgia, 16.
 Maryland to Mexico.
- COLUBER 40
Coryphodon D. & B., 1854, Erp., VII, 178; Gthr., 1858, Cat. 107.
- (BASCANIUM.)
- COLUBER CONSTRICTOR (Black Snake) 41
 Linn., 1758, Syst., I, 216. — 1766, Syst., I, 385; Gmel., 1788, Syst. Linn., I, 1109; Shaw, 1802, Zool., III, 464; Latr., 1802, Rept., IV, 178; Merr., 1820, Syst., 108; Daud., 1803, Rept., VI, 402; Harl., 1827, Jour. Phil. Ac., 348, — 1835, Med. Res., 112; Schleg., 1837, Essai, I, 114, II, 133, pl. 5, f. 3 and 4; Stor., 1839, Rept. Mass., 225; Holbr., 1842, Herp., III, 55, pl. 11; Thomps., 1842, Vermont, 117; DeK., 1842, N. Y. Rept., 35, pl. 10, f. 20; Mart., 1856, Mus. Berl., 24; Jones, 1865, Rept. Nova Scotia, 4.

OPHIDIA—Continued.

- Bascanium constrictor* B. & G., 1853, Serp., 93; Baird, 1854, Serp. X, Y., 22; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 67; Allen, 1869, Pr. Bost. Soc., 180; Cope, 1875, Checklist, 40, — 1877, Pr. Am. Phil. Soc., 64.
- Coryphodon constrictor* D. & B., 1854, Erp., VII, 183; Gthr., 1858, Cat., 108; Jan, 1863, Sist., 64, — 1867, Icon., livr. 22, pl. 4, f. 3, — 1876, Icon., livr. 48, pl. 6, f. 1.
- Bascanium Foxii* B. & G., 1853, Serp., 96; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 69. *Hierophis constrictor* Bonap., 1811, It. Fauna, II.
- The Black Snake* Catesby, 1743, Carol., II, 48, pl. 48; Kalm., 1764, Reise Amer., II, 202; Penn., 1792, Arct. Zool., II, Supp., 92.
- Texas to Nova Scotia.
- Var. FLAVIVENTRIS (Blue Racer) 42
- Coluber flaviventris* Say, 1823, Long's Exp., I, 185.
- Bascanium flaviventre* B. & G., 1853, Serp., 96; Bd., 1859, Mex. Bound., II, Rept., 20, — P. R. R. Rept., X, pl. 31, f. 70; Hayd., 1862, Trans. Phil. Soc., 177; Allen, 1874, Pr. Bost. Soc., 69; Yarr., 1875, Wheel. Rep., V, 511, 515, and 541; Cones & Yarrow, 1878, Hayd. Rep., IV, 284. *Coryphodon flaviventris* Hall., 1856, Pr. Phil. Ac., 241.
- C. constrictor* var. *flaviventris* Jan, 1863, Sist., 64, — 1867, Icon., livr. 22, pl. 3, f. 1-2, — 1876, Icon., livr. 48, pl. 6, f. 2.
- Bascanium ectustus* B. & G., 1853, Serp., 97; Girard, 1858, Expl. Exp., Rept., 127; Bd., 1859, P. R. R. Rep., X, pl. 36, f. 6; Cooper, 1860, P. R. R. Rep., XII (2), 301.
- Bascanium constrictor* subsp. *ectustum* Yarrow, 1875, Wheeler's Rep., V, 541; Cope, 1875, Checklist, 40. *Coryphodon constrictor* var. *ectustum* Jan, 1863, Sist., 64, — Icon., livr. 22, pl. 4, f. 1-2. (?) *Bascanium Fremontii* B. & G., 1853, Serp., 95; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 68.
- Mississippi basin westward to the Pacific.
- Var. MENTOVARIVS 42
- Coryphodon mentovarivus* D. & B., 1854, Erp., VII, 187; Jan, 1863, Sist., 64. *Bascanium mentovarivum* Cope, 1879, Pr. Am. Phil. Soc., 271.
- Mexico; Tehuantepec.
- (MASTICOPIHIS.)
- COLUBER FLAGELLIFORMIS ("Coach-whip Snake") 42
- Anguis flagelliformis*, The Coach-whip Snake, Catesby, 1743, Carol., II, 54. *Coluber flagellum* Shaw, 1802, Zool., III, 475. *Coluber flagelliformis* Holbr., 1836, Herp., I, 107, pl. 19. *Psammocephalus flagelliformis* Holbr., 1842, Herp., IV, 11, pl. 2. *Masticophis flagelliformis* B. & G., 1853, Serp., 98; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 71, pl. 32, f. 72; Jan, 1863, Sist., 65, — Icon., livr. 20, pl. 6, f. 1. *Heterodryas flagelliformis* D. & B., 1854, Erp., VII, 210; Mart., 1856, Mus. Berl., 26; Gthr., 1858, Cat., 118. *Drynobius flagelliformis* Cope, 1860, Pr. Phil. Ac., 561. *Bascanium flagelliforme* Cope, 1877, Pr. Am. Phil. Soc., 64. *Bascanium flagelliforme* subsp. *flagelliforme* Cope, 1875, Checklist, 40. *Heterodryas psammophis* Schleg., 1837, Essai, I, 152, II, 195.
- Coach-whip Snake* Bartram, 1791, Travels, 219.
- Southern States westward to Louisiana.

OPHIDIA—Continued.

- VAR. TESTACEUS 43
Coluber testaceus Say, 1823, Long's Exp., II, 48; Harl., 1827, Jour. Phil. Ac., 348, — 1853, Med. Res., 113; Holbr., 1842, Herp., III, 63, pl. 13; B. & G., 1853, Serp., 150. *Coryphodon testaceus* Gthr., 1858, Cat., 108. *Drymobius testaceus* Cope, 1860, Pr. Phil. Ac., 561. *Masticophis testaceus* B. & G., 1859, Mex. Bound., II, Rept., 20; Baird, 1859, P. R. R. Rep., X, 43; Cope, 1866, Pr. Phil. Ac., 305. *Bascanium flagelliforme* subsp. *testaceum* Cope, 1875, Checklist, 40, — 1880, Bull. 20, U. S. Mus., 23; Yarr., 1875, Wheeler's Rep., V, 542; Coues, 1875, Wheel. Rep., V, 617. *Masticophis flagelliformis* var. *testaceus* Jan, 1863, Sist., 65, — 1867, Icon., livr. 20, pl. 6, f. 2. *Psammophis flavigularis* Hall, 1852, Pr. Phil. Ac., 178, — 1853, Sitgreave's Exp., I, 31, 146. *Masticophis flavigularis* B. & G., 1853, Serp., 99, 159, — B. & G., 1854, Marey's Exp., 201; Bd., 1859, P. R. R. Rep., X, pl. 32, f. 73. *Herpodytes flavigularis* Hall, 1859, P. R. R. Rep., X, Rept., 12. (?) *Coluber mormon* B. & G., 1852, Stansbury's Exp., 351. *Masticophis mormon* B. & G., 1853, Serp., 101; Bd., 1859, P. R. R. Rep., X, pl. 32, f. 74.
Dakota to Texas and the Pacific Coast.
- VAR. AURIGULUS. 44
Drymobius aurigulus Cope, 1861, Pr. Phil. Ac., 301, — 1875, Checklist, 40.
Lower California.
- COLUBER MEXICANUS 45
Zamenis mexicanus D. & B., 1854, Erp., VII, 695. (?) *Masticophis mexicanus* Jan, 1863, Sist., 65, — 1867, Icon., livr. 22, pl. 6, f. 1. *Phimothyra mexicana* Cope, 1861, Pr. Phil. Ac., 167.
Mexico.
- COLUBER SPINALIS. 45
Masticophis spinalis Peters, 1866, MB. Berl. Akad., 91.
- COLUBER TAENIATUS. 46
Leptophis taeniatus Hall, 1852, Pr. Phil. Ac., 181, — 1853, Sitgreave's Exp., 133, 146. *Masticophis taeniatus* B. & G., 1853, Serp., 103; Bd., 1859, P. R. R. Rep., X, 20, pl. 23, pl. 32, f. 76; Cooper, 1860, P. R. R. Rep., XII, 302; Jan, 1863, Sist., 64, — 1867, Icon., livr. 22, pl. 5; Cope, 1866, Pr. Phil. Ac., 305. *Drymobius taeniatus* Cope, 1860, Pr. Phil. Ac., 561. *Bascanium taeniatum* subsp. *taeniatum* Cope, 1875, Checklist, 40. *Bascanium taeniatum* Coues, 1875, Wheeler's Rep., V, 616. *Leptophis lateralis* Hall, 1853, Pr. Phil. Ac., 237, — 1859, P. R. R. Rep., X, Rept., 13, pl. 4, f. 3. *Masticophis schottii* B. & G., 1853, Serp., 160; Bd., 1859, Mex. Bound., II, 20, — P. R. R. Rep., X, pl. 32, f. 77. *Bascanium taeniatum* subsp. *laterale* Cope, 1875, Checklist, 40; Yarr., 1875, Wheel. Rep., V, 543. *Bascanium laterale* Yarr., 1875, Wheeler's Rep., V, 512. *Bascanium taeniatum* var. *ornatum* Coues, 1875, Wheeler's Rep., V, 617; Cope, 1875, Checklist, 40. *Masticophis ornatus* B. & G., 1853, Serp., 102, 159; Bd., 1859, P. R. R. Rep., X, pl. 32, f. 75. *Bascanium flagelliforme* subsp. *piccum* Cope, 1875, Checklist, 40; Coues, 1875, Wheeler's Rep., V, 617. *Bascanium taeniatum* subsp. *piccum* Cope, 1880, Bull. 20, U. S. Mus., 24.
From the Plains to the Pacific.

OPHIDIA—Continued.

- Var. **BILINEATUS**
- Masticophis bilineatus* Jan, 1863, Sist., 40, — 1867, Icon., livr. 22, pl. 6, f. 1.
- Scales smooth, in 17 rows. Oculars 2-2, lower antecular very small. Temporals 2+2+2. Labials 8, fourth and fifth in orbit, or 9, fifth and sixth touching the eye, seventh largest. Infralabials 10 to 11. Back black, with two narrow lines of light color along each flank. Orbitals, loreal, and nasals each with white spot. Labials and ventral surface light.
- Mexico.
- SPILOTES** 47
- Wagler, 1830, Amph., 179; Gthr., 1858, Cat., 96. *Georgia* B. & G., 1853, Serp., 92.
- (GEORGIA.)
- SPILOTES COUPERI** 48
- Cope, 1860, Pr. Phil. Ac., 342, 564, — 1875, Checklist, 39. *Coluber couperi* Holbr., 1842, Herp., III, 75, pl. 16. *Georgia couperi* B. & G., 1853, Serp., 92; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 66.
- Gulf States.
- SPILOTES OBSOLETUS** 48
- Coluber obsoletus* Holbr., 1842, Herp., III, 61, pl. 12. *Georgia obsoleta* B. & G., Serp., 158; Bd., 1859, P. R. R. Rep., X, pl. 31, f. 66. *Spilotes crebennus* Cope, 1860, Pr. Phil. Ac., 342, 564, — 1864, Pr. Phil. Ac., 167, — 1875, Checklist, 39; Sumichr., 1873, Arch. Sci., 246. *Spilotes corais crebennus* Cope, 1877, Pr. Am. Phil. Soc., 64.
- Florida to Texas.
- (SPILOTES.)
- SPILOTES CORAIS** 48
- Coluber corais* (Cuv.) Boie, 1827, Isis, 537; Schleg., 1837, Essai, II, 139, pl. 5, f. 9-11, 1844 (?), Abbild., pl. 28, f. 9-11 (skull).
- Spilotes corais* D. & B., 1854, Erp., VII, 222; Gthr., 1858, Cat., 98; Jan, 1863, Sist., 62, — 1876, Icon., livr. 48, pl. 4, f. 6, pl. 5, f. 1.
- Brazil to Mexico.
- SPILOTES VARIABILIS** 49
- Dum. Bibr., 1854, Erp., VII, 220; Gthr., 1858, Cat., 99; Sumichr., 1873, Arch. Sci., 245. *Coluber variabilis* Max., 1825, Beitr., 271, — 1830, Abbild., Lief. 14, pl. 2, f. 3-6; Boie, 1827, Isis, 537. *Coluber pallatus* Linn., 1754, Mus. Ad. Fridr., pl. 20, f. 3. *Crasastes mericanus* & *C. coronatus* Laur., 1768, Synops., 83. *Coluber mericanus* Gmel., 1788, Syst. Nat. Linn., I, 1088. *Coluber plutonius* Daud., 1803, Rept., 324; Merr., 1790, Beitr. (2), pl. 12. *Coluber caninana* Merr., 1820, Syst., 121. *Spilotes pallatus* Wagl., 1830, Amph., 179; Schleg., 1837, Essai, 149, pl. 6, f. 1, 2; Fitz., 1843, Syst., 26; Cope, 1861, Pr. Phil. Ac., 300.
- Scha, 1735, Thesaur., II, pl. 20, f. 1, pl. 105, f. 4; Scheuch., 1736, Bibl. Sacr., pl. 662, f. 11, pl. 747, f. 3.
- Mexico to Brazil.

OPHIDIA—Continued.

- Var. *AURIBUNDUS* 50
Spilotes pollatus subsp. *auribundus* Cope, 1861, Pr. Phil. Ac., 300. *Spilotes auribundus* (N. Salmii Gthr.) Cope, 1864, Pr. Phil. Ac., 167; Sumichr., 1873, Arch. Sci., 245.
Mexico.
- SPILOTES MELANURUS* 50
Dum. Bibr., 1854, Erp., VII, 224; Jan., 1863, Sist., 63, — 1876, Icon., livr. 48, pl. 5, f. 2. *S. coxalis* var. *melanurus* Gthr., 1858, Cat., 99.
Mexico.
- SPILOTES PORCHILONOTUS* 50
Guenther, 1858, Cat., 100; Sumichr., 1873, Arch. Sci., 245.
Mexico.
- PITYOPHIS* 51
Holbrook, 1842, Herp., IV, 7; Harl., 1852, Pr. Phil. Ac., 181; B. & G., 1853, Serp., 64; D. & B., 1854, Erp., VI, 232; Gthr., 1858, Cat., 85.
- PITYOPHIS MELANOLEUCUS* (Bull Snake) 51
Holbr., 1842, Herp., IV, 7, pl. 1; B. & G., 1853, Serp., 65; D. & B., 1854, Erp., VII, 233; Mart., 1856, Mus. Berl., 26; Gthr., 1858, Cat., 86; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 44; Cope, 1875, Checklist, 39, — 1877, Pr. Am. Phil. Soc., 64. *Coluber melanoleucus* Daud., 1803, Rept., VI, 409; Harl., 1827, Jour. Phil. Ac., 359, — 1835, Med. Res., 122.
Pine Snake or *Bull Snake* Bartram, 1791, Travels, 276.
East of the Mississippi, south of Ohio and Pennsylvania.
- PITYOPHIS CATENIFER* 52
Coluber catenifer Blainville, 1835, Nouv. Ann. Mus., IV, 290, pl. 26, f. 2. *Pityophis catenifer* B. & G., 1853, Serp., 69; Grd., 1858, Expl. Exp., Rept., 135; Bd., 1859, P. R. R. Rep., X, pl. 36, f. 4; Gthr., 1858, Cat., 87; Cope, 1875, Checklist, 39; Bd., 1859, P. R. R. Rep., X, Williamson's Route, 11. *P. novaeclaus* B. & G., 1853, Serp., 72; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 48; *P. wilkisi* B. & G., 1853, Serp., 71; Grd., 1858, Wilkes' Exp., Rept., 137, pl. 9, f. 1—7; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 47. *P. melanoleucus* var. *catenifer* Jan, 1863, Sist., 59, — 1867, Icon., livr. 22, pl. 1, f. 1. *P. hermanni* Hall., 1853, Pr. Phil. Ac., 236. *Coluber vertebralis* Blainv., 1835, Ann. Mus., IV, 293, pl. 27, f. 2; B. & G., 1853, Serp., 152. *Pityophis vertebralis* D. & B., 1854, Erp., VII, 238; Gthr., 1858, Cat., 86; Cope, 1861, Pr. Phil. Ac., 300, — 1875, Checklist, 39; Hall., 1859, P. R. R. Rep., X, 14. *P. hornatais* Cope, 1860, Pr. Phil. Ac., 342. *P. melanoleucus* var. *vertebralis* Jan, 1863, Sist., 59, — 1867, Icon., livr. 22, pl. 1, f. 3.
Oregon to Mexico.
- Var. *SAYI* 52
Coluber sayi Schleg., 1837, Essai, II, 157; B. & G., 1853, Serp., 151. *C. melanoleucus* var. Harl., 1827, Jour. Phil. Ac., 360, — 1835, Med. Res., 123. *Pityophis melanoleucus* Max., 1863, Reise N. Amer., 95. *P. sayi* B. & G., 1853, Serp., 152; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 45;

OPHIDIA—Continued.

- Coop., 1860, P. R. R. Rep., XII (2), 300; Hayden, 1862, Trans. Phil. Soc., 177. *P. sayi* subsp. *sayi* Cope, 1875, Checklist, 39. *P. McClintockii* B. & G., 1853, Serp., 68. 1854, Macey's Exp., 196, pl. 5; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 47.
Rocky Mountains to Illinois.
- Var. MEXICANUS. 52
Pityophis mexicanus D. & B., 1854, Erp., VII, 236; Jan, 1863, Sist., 59, — 1867, Icon., livr. 22, pl. 2, f. 1. *Amsime mexicanus* D. & B., 1854, Erp., Atlas, pl. 62. (?) *Pityophis sayi* var. *mexicanus* Cope, 1875, Checklist, 39. (?) *P. sayi* subsp. *mexicanus* Yarr., 1875, Wheeler's Rep., V, 539. (?) *P. sayi mexicanus* Cope, 1880, Bull. 20, U. S. Mus., 23. Mexico.
- Var. DEPPEI
Elaphis deppci D. & B., 1854, Erp., VII, 268. *Pityophis deppci* Jan, 1863, Sist., 59, — 1867, Icon., livr. 22, pl. 2, f. 2. *P. deppci* var. *pholidotictus* Jan, 1863, Sist., 59. *Arizona elegans* (Kenn.) Bd., 1859, P. R. R. Rep., X, 42, — 1859, Mex. Bound., II, pl. XIII. *Pityophis elegans* Cope, 1875, Checklist, 39; Yarr., 1875, Wheel. Rep., V, 541, & Coues, p. 618. (?) *Arizona jani* Cope, 1860, Pr. Phil. Ac., 369. (?) *Arizona lineaticollis* Cope, 1861, Pr. Ac. Phil., 300.
Scales in 29 to 31 rows, keels of vertebral low to obsolete. Oculars 1 to 2—2. Loreal narrow. Temporals 3+3 to 4. Prefrontals commonly two, sometimes four. Ventrals 235; anal entire; subcaudals 67 pairs. Dorsal and lateral series of blotches more or less irregular in shape. Head nearly uniform light.
Arizona to Mexico.
- Var. BELLONA. 53
Churchillia bellona B. & G., 1852, Stansbury's Exp., 350. *Pityophis bellona* B. & G., 1853, Serp., 66, 157; Gthr., 1858, Cat., 87; (Kenn.) Bd., 1859, P. R. R. Rep., X, 42, pl. 28, f. 46, — Mex. Bound., II, 18; Cope, 1866, Pr. Ac. Phil., 305; Allen, 1874, Pr. Bost. Soc., 69. *P. sayi* var. *bellona* Cope, 1875, Checklist, 39. *P. sayi* subsp. *bellona* Yarr., 1875, Wheeler's Rep., V, 540, & Coues, p. 617; Coues & Yarr., 1878, Hayd. Rep., IV, 282. *P. affinis* Hall, 1852, Pr. Phil. Ac., 181, — 1853, Sitgr. Exp., 130, 146.
The Utah basin.
- ELAPHIS 53
Aldrovandus, 1640, Hist. Serp. Draec., 267 (re-print, 1765); Bonap., 1831, Saggio. Dum., 1852, Mem. Sci. Nat., XXIII, 453; D. & B., 1854, Erp., VII, 243; Gthr., 1858, Cat., 92. *Elaphe* Fitz., 1833, Wagler's Icon., pl. 27, — 1843, Syst., 26.
(SCOTOPHIS.)
- ELAPHIS OBSOLETUS (Black Snake) 54
Coluber obsoletus Say, 1823, Long's Exp., I, 140; Harl., 1827, Jour. Phil. Ac., 347, — 1835, Med. Res., 112. *Scotophis obsoletus* Kenn., 1860, Pr. Phil. Ac., 330. *Coluber obsoletus* subsp. *obsoletus* Cope, 1875, Checklist, 39. *Elaphis spiloides* D. & B., 1854, Erp., VII, 269; Gthr., 1858, Cat., 90. Mississippi Valley.

OPHIDIA—Continued.

Var. ALLEGHANIENSIS (Black Pilot Snake) 54

Coluber alleghaniensis Holbr., 1842, Herp., III, pl. 19. *Scotophis alleghaniensis* B. & G., 1853, Serp., 73; (Kenn.) Bd., 1859, P. R. R. Rep., X, pl. 29, f. 49. *Elophis alleghaniensis* Hall., 1856, Pr. Phil. Ac., 243; Mart., 1856, Mus. Berl., 27; Jan., 1863, Sist., 62, — 1867, Icon., livr. 24, pl. 2, f. 1-2; Allen, 1869, Pr. Bost. Soc., 181. *E. Holbrookii* D. & B., 1854, Erp., VII, 272. *E. rubriceps* D. & B., 1854, Erp., VII, 270. *E. alleghaniensis* vars. *rubriceps* & *holbrookii* Jan., 1863, Sist., 62. *Scotophis confinis* B. & G., 1853, Serp., 76; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 52.

New England to Alabama.

Var. LINDHEIMERI 54

Scotophis lindheimeri B. & G., 1853, Serp., 74; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 50. *Coluber lindheimeri* Cope, 1875, Checklist, 39, — 1880, Bull. 20, U. S. Mus., 23. *Scotophis rhinomegas* Cope, 1860, Pr. Phil. Ac., 255. *S. latus* B. & G., 1853, Serp., 77, — 1854, Macey's Exp., 198, pl. 6; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 53. *S. calligaster* Kenn., 1859, Pr. Phil. Ac., 98. *Elophis guttatus* var. *calligaster* Jan., 1863, Sist., 62, — 1867, Icon., livr. 21, pl. 6, f. 2. *E. guttatus* var. *rhinomegas* Jan., 1863, Sist., 62. *Scotophis emoryi* B. & G., 1853, Serp., 157; (Kenn.) Bd., 1859, P. R. R. Rep., X, 42, pl. 30, f. 56. *Coluber emoryi* Cope, 1880, Bull. 20, U. S. Mus., 23.

Illinois to Texas.

Var. FAIRDI

Coluber fairdi (Yarrow) Cope, 1880, Bull. U. S. Nat. Mus., p. 41.

Vertical plate longer than broad, with a slight notch in anterior border; posterior portion very large, broader than long. Nine upper labials, seventh largest. Lower labials twelve, seventh largest. Dorsal rows of scales 27, long and lozenge-shaped; three upper dorsal rows slightly carinated. General color above (alcoholic) warm grayish-ash; beneath yellowish; behind occipitals two converging oblong brown blotches, and posterior to these a series of narrow transverse brown blotches, eighty in number, becoming obsolete near caudal extremity; these blotches are six scales in width (Yarrow).

Fort Davis, Texas.

ELAPHS GUTTATUS (CORN SNAKE) 55

Coluber guttatus Linn., 1766, Syst., I, 385; Daub., 1771, Encycl. Meth., 602; Gmel., 1788, Syst. Linn., I, 1110; Schleg., 1837, Essai, I, 149, II, 168; Holbr., 1842, Herp., III, 65, pl. 14; Harl., 1827, Jour. Phil. Ac., 363, — 1835, Med. Res., 126; Holbr., 1838, Herp., II, 109; Gthr., 1858, Cat., 89; Cope, 1875, Checklist, 39. *C. carolinianus* Shaw, 1802, Zööl., III, 460, pl. 19. *C. maculatus* Latr., 1802, Rept., IV, 73; Harl., 1827, Jour. Phil. Ac., 362, — 1835, Med. Res., 125. *C. molossus* Daud., 1803, Rept., VI, 269; Harl., 1827, Jour. Phil. Ac., 363, — 1835, Med. Res., 126. *C. floridanus* Harl., 1827, Jour. Phil. Ac., 360, — 1835, Med. Res., 124. *C. compressus* (Donnd.) Bechst., 1802, Ueb.

OPHIDIA—Continued.

- LaC., Amph., IV, 236, pl. 36, f. 2. *C. pantherinus*, p. 102, *Natrix guttata*, p. 99, and *N. maculatus*, p. 124, Merr., 1820, Syst. Amph. *Scotophis guttatus* D. & B., Erp., VII, 273; Mart., 1856, Mus. Berl., 27; Jan, 1863, Sist., 62, — 1867, Icon., livr. 21, pl. 6, f. 1. *Elaphis guttatus* D. & B., 1854, Erp., VII, 273; Mart., 1856, Mus. Berl., 27; Jan, 1863, Sist., 62, — 1867, Icon., livr. 21, pl. 6, f. 1.
- LaC., 1789, Quad. Ovip., II, 329; Bonn., 1790, Encycl. Meth., Ophiol., 19; Merr., 1790, Beitr., H. t. II.
- Corn Snake* Catesby, 1743, Carol., II, pl. 55.
Southeastern States to Virginia.
- Var. VULPINUS** 56
Scotophis vulpinus B. & G., 1853, Serp., 75; Bd., 1859, P. R. R. Rep., X, pl. 29, f. 51; (Kenn.) Coop., 1860, P. R. R. Rep., XII (2), 99. *Coluber vulpinus* Cope, 1875, Checklist, 39.
Massachusetts to Nebraska.
- ELAPHIS QUADRIVITTATUS** (Chicken Snake) 56
Coluber quadrivittatus Holbr., 1842, Herp., III, 89, pl. 20; Gthr., 1858, Cat., 88; Cope, 1875, Checklist, 39, — 1877, Pr. Am. Phil. Soc., 64. *Scotophis quadrivittatus* B. & G., 1853, Serp., 80; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 55. *Elaphis quadrivittatus* D. & B., 1854, Erp., VII, 265; Jan, 1863, Sist., 62.
Florida to Virginia.
- DROMICUS** 56
Bibron, 1843, Rept. Cuba, Sagra, 221; D. & B., 1854, Erp., VII, 646; Gthr., 1858, Cat., 126.
- DROMICUS LAUREATUS** 57
Guenther, 1868, Ann. Mag., 419.
"City of Mexico."
- DROMICUS FLAVILATUS** 58
Cope, 1871, Pr. Phil. Ac., — 1875, Checklist, 38, — 1877, Pr. Am. Phil. Soc., 65.
Florida to North Carolina.
- DROMICUS MARGARITIFERUS** 58
Guenther, 1858, Cat., 126. *Herpetodryas margaritifera* Schleg., 1837, Essai, I, 151, II, 184, — Abbild., t. 44. *Leptophis margaritifera* D. & B., 1853, Erp., VII, 539; Mart., 1856, Mus. Berl., 26. *Thamnosophis margaritifera* Jan, 1863, Sist., 82, — 1868, Icon., livr. 31, pl. 6, f. 3. *Zamenis tricolor* Hall, 1855, Jour. Phil. Ac., 31, f. 3. *Coluber hickmanella* Shaw, 1802, Zool., III, 511.
Mexico and Central America.
- DROMICUS PUTNAMI** 59
Jan, 1863, Sist., 67, — 1867, Icon., livr. 6, f. 3. *Tomodon nasutus* Cope, 1864, Pr. Phil. Ac., 166.
Southern Mexico.

OPHIDIA—Continued.

- CORONELLINAE 59
Coronellator Gthr., 1858, Cat., 22.
- TACHYMENIS 60
 Wiegmann, 1854, Nov. Act. Acad. Caes. Leop., 252; Gthr., 1858, Cat., 33.
- TACHYMENIS LINEATA 60
Tomoloba lineatum D. & B., 1854, Erp., VII, 936, — Atlas, pl. 73 (named *Eudromus lineatus*); Jan, 1863, Sist., 57, — 1866, Icon., livr. 19, pl. 6, f. 3. *Pseudomphis lineatus* Gthr., 1858, Cat., 135. Mexico.
- TACHYMENIS IMPERIALIS 61
Tachymphis imperialis B. & G., 1855, Gilliss' Exp., II, 215 (name); Bd., 1859, Mex. Bound. Rept., 23, pl. 19, f. 1, — P. R. R. Rep., X, pl. 33, f. 87. *Coniophanes imperialis* Cope, Pr. Phil. Ac., 74, — 1875, Checklist, 38. Texas and Mexico.
- TACHYMENIS LATERITA 61
Coniophanes laterita Cope, 1861, Pr. Phil. Ac., 524.
 "Guadalaxara, Mexico."
- TACHYMENIS PROTEROPS 62
Coniophanes proterops Cope, 1860, Pr. Phil. Ac., 249.
 Mexico, New Grenada.
- TACHYMENIS FISSIDENS 62
Coronella fissidens Gthr., 1858, Cat., 36.
Glyphyrophis lateralis Jan, 1863, Sist., 54, — 1866, Icon., livr. 18, pl. 5, f. 3. Mexico.
- TACHYMENIS BIPUNCTATA 63
Coronella bipunctata Gthr., 1858, Cat., 36.
Glyphyrophis pictus Jan, 1863, Sist., 54, — 1866, Icon., livr. 18, pl. 5, f. 4. Mexico.
- TACHYMENIS MELANOCEPHALA 63
 Peters, 1869, Berl. Akad. Monatsb., 876. Mexico.
- ERYTHROLAMPRES 63
 Boie, 1826, Isis, 981; Wagl., 1830, Syst., 187; D. & B., 1854, Erp., VII, 843; Gthr., 1858, Cat., 47.
- ERYTHROLAMPRES GUENTHERI 63
E. caustissimus var. D. Gthr., 1858, Cat., 48.
 "Mexico(?)"
- OPHIBOLUS 64
 Baird & Girard, 1853, Serp., 82.
- OPHIBOLUS DOLIATUS 64
Coluber doliatus Linn., 1766, Syst., I, 379; Gmel., 1788, Syst. Linn., I, 1096; Harl., 1827, Jour. Phil. Ac., 362, — 1835, Med. Res., 125. *Coronella doliata* Holb., 1842, Herp., II, 105, pl. 24; D. & B., 1854, Erp., VII, 621; Gthr., 1858, Cat., 41. *C. coccinea* Schleg., 1837, Essai, I, 135, II, 57, pl. 2, f. 1. *Ophibolus doliatus* sub-sp. *doliatus* Cope, 1875, Checklist, 37 (part). *Sphenophis coccinea* Fitz., 1843, Syst., 25.
 Southeastern States.

OPHIDIÆ—Continued.

- Var. ELAPSOIDEUS 65
Oscrota elapsoidea Baird & Girard, 1853, Serp., 133; Mart., 1856, Mus. Berl., 24; Bd., 1859, P. R. R. Rep., X, pl. 33, f. 97; Cope, 1875, Checklist, 36, — 1877, Pr. Am. Phil. Soc., 65. *Calamaria elapsoidea* Holbr., 1842, Herp., III, 119, pl. 28.
Coronella coccinea Jan, 1863, Sist., 46, — 1866, Icon., livr. 17, pl. 1, f. 1.
Cenophora coccinea Jan, 1865, Icon., livr. 11, pl. 5, f. 1.
Southern States east of the Mississippi.
- OPHIBOLUS TRIANGULUS 65
La Triangle LaC., 1789, Serp., II, 331. *Coluber triangulus* Boie, 1827, Isis, 537. *Ablates triangulum* D. & B., 1854, Erp., VII, 315. *Lampropeltis triangula* Cope, 1860, Pr. Phil. Ac., 256; Allen, 1869, Pr. Bost. Soc., 180. *Coluber eximius* Stor., 1839, Mass. Rep., 227; Holbr., 1842, Herp., III, 69, pl. 15; DeK., 1842, N. Y. Rept., 28, pl. 12, f. 25; Gthr., 1858, Cat., 91. *Ophibolus eximius* & *clericus* B. & G., 1853, Serp., 87 and 88; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 61, 62, — 1854, Serp. N. Y., 21. *O. doliatus* var. *triangulus* Cope, 1875, Checklist, 37. *Coronella eximia* Jan, 1863, Sist., 46, — 1866, Icon., livr. 17, pl. 1, f. 3.
Pseudelaps y Berth., 1843, Abh. Gött., I, 67, pl. 1, f. 11 and 12. *Ablates triangulum* var. *clericus* Hall, 1856, Pr. Phil. Ac., 246.
Coluber eximius of Harlan, with 250 + 33 to 60 ventrals and subcaudals; is probably a species of *Scotophis*.
House Snake; Chicken Snake; Milk Snake; or Thunder Snake.
East of the Mississippi to Canada.
- Var. CALLIGASTER 66
Coluber calligaster Harl., 1835, Med. Res., 122. *Ablates triangulum* var. *calligaster* Hall, 1856, Pr. Ac. Phil., 244. *Lampropeltis calligaster* Cope, 1860, Pr. Phil. Ac., 255. *Ophibolus calligaster* Cope, 1875, Checklist, 37. *O. evansi* Kenn., 1859, Pr. Phil. Ac., 99. (?) *Coronella evansi* Jan, 1863, Sist., 47, — 1866, Icon., livr. 17, pl. 2, f. 3.
Northern portion of the Mississippi Valley.
- Var. MEXICANUS 66
San Luis Potosi.
- Var. DOLIATUS 66
Ophibolus doliatus B. & G., 1853, Serp., 89; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 63. *Coronella doliata* var. *gentilis* Jan, 1863, Sist., 46, — 1866, Icon., livr. 17, pl. 1, f. 2. *Lampropeltis multistriata* Kenn., 1860, Pr. Phil. Ac., 328. *Coronella doliata* Jan, 1863, Sist., 46, — 1865, Icon., livr. 14, pl. 4, f. 1.
Nebraska and southward.
- Var. GENTILIS 66
Ophibolus gentilis B. & G., 1853, Serp., 90, — 1854, Marey's Exp., 229, pl. 8; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 64. *Lampropeltis doliata* Cope, 1860, Pr. Phil. Ac., 256. *Ophibolus doliatus* subsp. *doliatus* Cope, 1875, Checklist, 37.
Arkansas to Utah.

OPHIDIA—Continued.

- VAR. ZONATUS 67
Coluber (zucholus) zonatus Blainv., 1835, Nouv. Ann. Mus., IV, 293; B. & G., 1853, Serp., 153. *Ballophis zonatus* Lockington, 1875, Pr. Cal. Acad. California.
- VAR. ANNULATUS
Lampropeltis annulata Kenn., 1860, Pr. Phil. Ac., 329; Cope, 1860, Pr. Phil. Ac., 257. *Ophibolus dolatus* subsp. *annulatus* Cope, 1875, Checklist, 36; Yarrow, 1875, Wheeler's Rep., V, 537. ? *Lampropeltis polyzona* Cope, 1860, Pr. Phil. Ac., 258. ? *L. anaurus* Cope, l. c., 258.
- Bright red, 18 to 22 pairs of black rings from head to vent, each pair inclosing a spotless yellow ring, which encircles the body, widening but little on the flanks. Belly between the yellow rings black. Top of head black, this color extending back upon the occipitals in an acute angle. A broad occipital yellow ring. (Kenn.)
 Mexico.
- VAR. RHOMBOMACULATUS
Coronella rhombomaculata Holbr., 1842, Herp., III, 103, pl. 23; Jan, 1863, Sist., 47, — 1866, Icon., livr. 17, pl. II, f. 1—2. *Ophibolus rhombomaculatus* B. & G., 1853, Serp., 86; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 60; Cope, 1875, Checklist, 37. *Lampropeltis rhombomaculatus* Cope, 1860, Pr. Phil. Ac., 255.
- Top of head and back chestnut brown, with a vertebral series of rhomboid reddish-brown spots. Lower surface salmon-colored, more or less bright. Ventrals 211; subcaudals 45. (Holbr.)
 Southeastern United States.
- OPHIBOLUS GETULUS (Chain Snake) 68
 Baird & Girard, 1853, Serp., 255; Bd., 1854, Serp. N. Y., 20, — 1859, P. R. R. Rep., X, pl. 31, f. 65. *Coluber getulus* Linn., 1766, Syst., I, 382; Gmel., 1788, Syst. Linn., I, 1106; Harl., 1827, Jour. Phil. Ac., 358, — 1835, Med. Res., 122; Peale, 1829, Macl. Lye., I, pl. V; Gthr., 1858, Cat., 249. *Coronella getula* Holbr., 1842, Herp., III, 95, pl. 21; D. & B., 1854, Erp., VII, 616; Jan, 1863, Sist., 47, — 1865, Icon., livr. 14, pl. 5, f. 1. *Lampropeltis getula* Cope, 1860, Pr. Phil. Ac., 255. *Ophibolus getulus* subsp. *getulus* Cope, 1875, Checklist, 37, — 1880, Bull. 20, U. S. Mus., 23. *Pseudolaps getulus* Fitz., 1826, Neue Class. Rept., 26.
- Chain Snake* Catesby, 1743, Carol., II, 52. Chain Snake; King Snake; Thunder Snake. Southern States to Texas.
- VAR. SAYI (King Snake) 68
Coronella sayi Holbr., 1842, Herp., 99, pl. 22; D. & B., 1854, Erp., VII, 619; Gthr., 1858, Cat., 41. *Coluber sayi* DeK., 1842, Rept., 41. *Herpobatrachus getulus* Schleg., 1837, Essai, I, 153, II, 198. *Lampropeltis getulus* Fitz., 1843, Syst., 25. *Lampropeltis sayi* Cope, 1860, Pr. Phil. Ac., 251. *Ophibolus sayi* B. & G., 1853, Serp., pp. 84, 159, — 1854, Marey's Exp., pl. 7; Bd., 1859, P. R. R. Rep., X, pl. 30, f. 59. *O. getulus* var. *sayi* Cope, 1875, Checklist, 37. *O. getulus sayi* Cope, 1880, Bull. 20, U. S. Mus., 23. *Coronella getulus* var. *sayi* Jan, 1863, Sist., 47, — 1865, Icon., livr. 14, pl. 5, f. 2. Mississippi Valley to Illinois and Texas.

OPHIDIA—Continued.

- Var. BOYLIU 69
Ophibolus boylii B. & G., 1853, Serp., 82; Bd., 1859, P. R. R. Rep., X, 11, pl. 30, f. 57, — 1859, Mex. Bound., II, Rept., 20; Cope, 1866, Pr. Phil. Ac., 305. *O. gyalus* subsp. *boylii* Cope, 1875, Checklist, 37; Yarrow, 1875, Wheeler's Rep., V, 538; Cones, 1875, Wheeler's Rep., V, 618. *Coronella ballata* Hall., 1853, Pr. Phil. Ac., 236, — 1859, P. R. R. Rep., X, 14. ? *C. gyalus* Mart., 1856, Mus. Berl., 25. *C. gyalus* var. *boylii* Jan, 1863, Sist., 47. *Lampropeltis boylii* Cope, 1860, Pr. Phil. Ac., 255. *L. boylii* var. *conjuncta* Cope, 1860, Pr. Phil. Ac., 255, — 1861, l. c., p. 301. *Coronella gyalus* var. *pseudogyalus* Jan, 1863, Sist., 47, — 1865, Icon., livr. 12, pl. 6, f. 2.
 California to Mexico.
- Var. SPLENDIDUS
Ophibolus splendidus B. & G., 1853, Serp., 83; Bd., 1859, Mex. Bound., II, Rept., pl. 14, p. 20, — P. R. R. Rep., X, pl. 30, f. 58; Cope, 1866, Pr. Phil. Ac., 310. *Lampropeltis splendida* Cope, 1860, Pr. Phil. Ac., 255. *Ophibolus gyalus* subsp. *splendidus* Cope, 1875, Checklist, 37; Cones, 1875, Wheeler's Rep., V, 619. *Coronella gyalus* var. *splendida* Jan, 1863, Sist., 47, — 1865, Icon., livr. 12, pl. 6, f. 1.
 Back black, crossed by 50 to 63 broad bands of light color, forming a spot in the middle of each scale. Nearly all scales on the flank, with more or less white in the center. The dorsal bands bifurcate on the flank, inclosing rhomboid spots of darker, which often extend upon the ventrals. Belly white, blotched with black.
 Arizona and Southern California to Mexico.
- Var. CALIFORNIÆ
Coluber (Ophis) californiæ Blainv., 1835, Ann. Mus., IV, pl. 27, f. 1; B. & G., 1853, Serp., 153. *Coronella californiæ* D. & B., 1854, Erp., VII, 623. *C. gyalus* var. *californiæ* Jan, 1863, Sist., 47, — 1865, Icon., livr. 14, pl. 5, f. 3. *Ophibolus californiæ* Cope, 1875, Checklist, 37.
 More yellow than black, the latter forming more or less effaced, anastomosing and broken longitudinal lines on the back. The vertebral lines darker. Oculars 1-2.
 California.
- Var. PYRRHOMELAS 67
Ophibolus pyrrhomelas Cope, 1866, Pr. Phil. Ac., 305. *O. pyrrhomelas* Cope, 1875, Checklist, 37; Yarrow, 1875, Wheeler's Rep., V, 537. *O. pyrrhomelas* Cones, 1875, Wheeler's Rep., V, 619.
 Arizona; Sonora.
- LIOPHIS 69
 Wagler, 1839, Amph., 187; D. & B., 1851, Erp., VII, 697; Gilr., 1858, Cat., 42.
- LIOPHIS EUAPOIDES 69
Pliocercus elapoides Cope, 1860, Pr. Phil. Ac., 253; Peters, 1869, Mb., Berl. Akad., 876. *Elapochrus deppii* Peters, 1860, Mb. Berl. Akad., 263. *Liophis tricinetus* Jan, 1863, Sist., 53, — 1866, Icon., livr. 18, pl. 4, f. 4-6.
 Mexico.

OPHIDIA—Continued.

- DIADOPHIS 70
 Baird and Girard, 1853, Serp., 112.
- DIADOPHIS ANNULLATUS 70
Enicognathus annulatus D. & B., 1854, Erp., VII, 335, pl. 80, f. 1-3; Jan, 1863, Sist., 51, — 1866, Icon., livr. 16, pl. 4, f. 3.
 Mexico.
- DIADOPHIS DECORATUS 71
 Cope, 1860, Pr. Phil. Ac., 250. *Coronella decorata* Gthr., 1858, Cat., 35.
Enicognathus vittatus Jan, 1863, Sist., 50, — 1866, Icon., livr. 16, pl. 2, f. 2-3. *Rhadinea decorata* Cope, 1875, Jour. Phil. Ac., 138.
 Southern Mexico.
- DIADOPHIS FULVIVITTIS 71
Rhadinea fulviritta Cope, 1875, Jour. Phil. Ac., 139.
 Southern Mexico.
- DIADOPHIS PUNCTATUS 72
 Baird & Girard, 1853, Serp., 112; Bd., 1859, P. R. R. Rep., X, pl. 33, f. 82; Cope, 1860, Pr. Phil. Ac., 250; Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 1; Bd., 1854, Serp. N. Y., 24; Allen, 1869, Pr. Bost. Soc., 182. *Coluber punctatus* Linn., 1766, Syst., I, 376; Gmel., 1788, Syst. Linn., I, 1089; Harl., 1827, Jour. Phil. Ac., 354, — 1835, Med. Res., 117; Storer, 1839, Rept. Mass., 225; Holbr., 1842, Herp., III, 81, pl. 18; DeK., 1842, Rept. N. Y., 39, pl. 14, f. 29; Jones, 1865, Rept. Nova Scotia, 5. *Cubimaria punctata* Schleg., 1837, Essai, I, 132, II, 39. *Spilotes punctatus* Swains., 1839, Hist. Amph., II, 364. *Ablabus punctatus* D. & B., 1854, Erp., VII, 310; Gthr., 1858, Cat., 28. *Diadophis punctatus* subsp. *punctatus* Cope, 1875, Checklist, 37. *D. punctatus* var. *pallidus* Cope, 1860, Pr. Phil. Ac., 250.
 Southern and Eastern States to Nova Scotia.
- Var. ARNYI 72
Diadophis arnyi Kenn., 1859, Pr. Phil. Ac., 99; Cope, 1875, Checklist, 38. *Liophis* (*Diadophis*) *arnyi* Gthr., 1868, Ann. Mag., 413. *Diadophis punctatus* var. *arnyi* Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 5. *D. dysops* Cope, 1860, Pr. Phil. Ac., 251, — 1875, Checklist, 38. *D. punctatus* var. *lactus* Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 6.
 Illinois to Arkansas.
- Var. DOUGLIS 72
Diadophis douglis B. & G., 1853, Serp., 114; ? Bd., 1859, P. R. R. Rep., X, pl. 32, f. 1, pl. 33, f. 84; Coates, 1875, Wheeler's Rep., V, 623. *D. punctatus* var. *douglis* Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 2. *Ablabus occipitalis* Gthr., 1858, Cat., 29. *Diadophis occipitalis* Cope, 1860, Pr. Phil. Ac., 250. *D. punctatus* var. *stictogynus* Cope, 1860, Pr. Phil. Ac., 250, — 1875 (subspecies), Checklist, 37, — 1880, Bull. 29, U. S. Mus., 23. *D. texensis* Kenn., 1860, Pr. Phil. Ac., 328.
 Texas to Mexico.

OPHIDIA—Continued.

- VAR. *AMABILIS*
- Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 4. *Diadophis amabilis* B. & G., 1853, Serp., 113; Bl., 1859, P. R. R. Rep., X, pl. 33, f. 83. *Diadophis pulchellus* B. & G., 1853, Serp., 115; Bl., 1859, P. R. R. Rep., X, pl. 33, f. 85; Yarrow, 1875, Wheeler's Rep., V, 538. *D. punctatus* var. *pulchellus* Cope, 1860, Pr. Phil. Ac., 250; Jan, 1863, Sist., 49, — 1866, Icon., livr. 15, pl. 6, f. 3.
- Greenish to blackish-brown, punctulate with black. Belly orange, more or less sprinkled or spotted with black. Occipital ring moderate. Scales in 15 rows. Ventrals 184 to 204. Subcaudals 59 to 60.
- Arizona to California.
- DIADOPHIS REGALIS** 73
- Baird & Girard, 1853, Serp., 115, 161; Bl., 1859, P. R. R. Rep., X, pl. 33, f. 86; Coues, 1875, Wheeler's Rep., V, 623; Cope, 1875, Checklist, 38. Mexico.
- RHINOCHEILUS** 73
- Baird & Girard, 1853, Serp., 120.
- RHINOCHEILUS LECONTEI** 73
- B. & G., 1853, Serp., 120, 161; Bl., 1859, P. R. R. Rep., X, pl. 33, f. 90; Cope, 1866, Pr. Phil. Ac., 304; Jan, 1863, Prodr. Icon. Ofid., II, — 1863, Sist., 43, — 1876, Icon., livr. 48, pl. 3, f. 1; Coues, 1875, Wheeler's Rep., V, 623; Cope, 1875, Checklist, 36, — 1880, Bull. 20, U. S. Mus., 23.
- Texas to California.
- VAR. *TESSELLATUS* VAR. *N.* 74
- Mexico.
- HETERODON** 74
- (Pal. de Beauv., 1799) Latr., 1802, Rept., IV, 32; Fitz., 1826, Class., 56; Gray, 1831, Synops. (An. King. Griff., IX), 85; Troost, 1836, Ann. N. Y. Lyc., III, 183; Schleg., 1837, Essai, II, 96; Holbr., 1842, Herp., III, 37; B. & G., 1853, Serp., 51; D. & B., 1854, Erp., VII, 764; Gthr., 1858, Cat., 82.
- HETERODON PLATYRHINUS** (Hog-nose Snake) 75
- Latreille, 1802, Rept., IV, 32, pl. 28, f. 1-3; Holbr., 1838, Herp., II, 97, pl. 21, — 1842, Herp., IV, pl. 17; B. & G., 1853, Serp., 51; Bl., 1859, P. R. R. Rep., X, pl. 28, f. 38; Stor., 1839, Rept., Mass., 231; D. & B., 1854, Erp., VII, 766; Gthr., 1858, Cat., 82; Jan, 1863, Sist., 44, — 1876, Icon., livr. 48, pl. 3, f. 2; Cope, 1875, Checklist, 43, — 1877, Pr. Am. Phil. Soc., 61; DeK., 1842, Rept. N. Y., 55; Bl., 1854, Serp. N. Y., 18; Allen, 1869, Pr. Bost. Soc., 182. *H. cognatus* B. & G., 1853, Serp., 54; Bl., 1859, Mex. Bound. Surv., II, 47, — P. R. R. Rep., X, pl. 28, f. 39. *Coluber heterodon* Daud., 1803, Rept., VII, 153, pl. 60, f. 28; Say, 1818, Am. Jour. Sci., I, 261; Harl., 1827, Jour. Phil. Ac., 357, — 1835, Med. Res., 120. *H. tigrinus* Troost, 1836, Ann. N. Y. Lyc., III, 189; DeK., 1842, Rept. N. Y., 52. *Hog-nose Snake* Catesby, 1743, Carol., II, 56.
- Eastern, Middle, and Southern States.

OPHIDIA—Continued.

- Var. *niger*. 76
 Jan, 1863, Sist., 44. *Heterodon niger* Troost, 1836, Ann. N. Y. Lyc., 186;
 Holbr., 1838, Herp., II, 105, pl. 23, — 1842, Herp., IV, 63, pl. 16;
 DeK., 1842, Rept. N. Y., 52; B. & G., 1853, Serp., 55; D. & B., 1854,
 Erp., VII, 769; Gthr., 1858, Cat., 83; Bd., 1859, P. R. R. Rep., X, pl.
 28, f. 40. *H. atmodus* B. & G., 1853, Serp., 57; Bd., 1859, P. R. R.
 Rep., X, pl. 28, f. 41. *H. platyrhinus* subsp. *atmodus* Cope, 1875,
 Checklist, 43. *Scytale niger* Daud., 1803, Rept., V, 342; Harl., 1827,
 Jour. Phil. Ac., 367, — 1835, Med. Res., 130. *Coluber cacodaemon*
 Shaw, 1802, Zool., III, 377, pl. 102. *C. thraso* Harl., 1835, Med. Res.,
 120. *Heterodon annulatus* Troost, 1836, Ann. Lye. N. Y., III, 188;
 DeK., 1842, Rept. N. Y., 52. *The Black Viper* Catesby, 1743, Carol.,
 II, 44. Southern States east of the Mississippi.
- HETERODON SIMUS. 76
 Holbr., 1842, Herp., III, 57, pl. 15; DeK., 1842, Rept. N. Y., 52; B. &
 G., 1853, Serp., 59; Mart., 1856, Mus. Berl., 29; Bd., 1859, P. R. R.
 Rep., X, pl. 28, f. 42, 42 b. *H. platyrhinus* Schleg., 1827, Essai, I, 140,
 II, pl. 3, f. 20-22. *Coluber sinus* Linn., 1766, Syst., I, 375; Gmel.,
 1788, Syst., Linn., I, 1086. *H. sinus* subsp. *sinus* Cope, 1875, Check-
 list, 43. Mississippi Valley to the Atlantic.
- Var. *nasicus*. 77
Heterodon nasicus B. & G., 1852, Stansbury's Exp., 352, — 1854, Marey's
 Exp., 193, pl. 4, — 1853, Serp., 61, 157; Bd., 1859, P. R. R. Rep., X,
 pl. 28, f. 43; Hall, 1853, Sitgreave's Exp., 147, — 1856, Pr. Phil. Ac.,
 249; Bd., 1859, P. R. R. Rep., X, Whipple's 41, Beckwith's 19, —
 1859, Mex. Bound., II, Rept., 18, pl. 11, f. 1; Hayd., 1862, Trans. Am.
 Phil. Soc., 177; Cope, 1866, Pr. Phil. Ac., 307; Allen, 1874, Pr. Bost.
 Soc., 69; Jan, 1863, Sist., 46, — 1865, Icon., livr. 10, pl. 5. *H. catesbyi*
 Gthr., 1858, Cat., 83. *H. sinus* subsp. *nasicus* Cope, 1875, Checklist,
 43; Yarrow, 1875, Wheeler's Rep., V, 555; Cones, 1875, Wheeler's
 Rep., V, 611. California to Texas and Nebraska.
- Var. *KENNERLYI*. 77
Heterodon kennerlyi Kenn., 1860, Pr. Phil. Ac., 336; Jan, 1865, Icon.,
 livr. 10, pl. 5, f. 2. Sonora, Mexico.
- CEMOPHORA. 77
 Cope, 1860, Pr. Phil. Ac., 244.
- CEMOPHORA COCCINEA. 78
 Cope, 1860, Pr. Phil. Ac., 244; Jan, 1863, Sist., 45, — 1865, Icon., livr.
 11, pl. 5, f. 2; Cope, 1875, Checklist, 36, — 1877, Pr. Am. Phil. Soc.,
 65. *Coluber coccineus* Blumenbach, 1788, Licht. & Voigt, Mag., pl. V;
 Gmel., 1788, Syst. Linn., I, 1097; Latr., 1802, Rept., IV, 138; Daud.,
 Rept., VII, pl. 83, f. 1; Harl., 1827, Jour. Phil. Ac., 356, — 1835,
 Med. Res., 119. *Heterodon coccineus* Schleg., 1827, Essai, I, 141, II,
 102, pl. 3, f. 15, 16. *Sinotes coccineus* D. & B., 1854, Erp., VII, 637, pl.
 82, f. 2; Mart., 1856, Mus. Berl., 25; Gthr., 1858, Cat., 26. *Rhinostoma*
coccinea Holbr., 1842, Herp., III, 125, pl. 30; B. & G., 1853, Serp.,
 118; Bd., 1859, P. R. R. Rep., X, pl. 33, f. 89. *Elaps coccineus* Merr.,
 1820, Syst. Amph., 145. Southern States east of Texas.

OPHIDIA—Continued.

- Var. *COPEI* 78
Cunophora copei Jan, 1863, Sist., 45, Icon., livr. 11, pl. 5, f. 3.
Tennessee.
- XENODON 79
Boie, 1827, Isis, 541; Schleg., 1837, Essai, II, p. 80; D. & B., 1854, Erp., VII, 733; Gthr., 1858, Cat., 53; Jan, 1863, Arch. Zool., 106.
- XENODON BERTHOLDI 79
Jan, 1863, Arch. Zool., II, 108, — 1863, Sist., 56, — 1866, Icon., livr. 19, pl. 4, f. 2. A variety of *X. colubrinus*, from Para, of Guenther.
Mexico.
- HYPSIGLENA 79
Cope, 1860, Pr. Phil. Ac., 246.
- HYPSIGLENA OCHRORHYNCHUS 80
H. ochrorhynchus Cope, 1860, Pr. Phil. Ac., 246, — 1875, Checklist, 38.
Lower California.
- Var. CHLOROPHAEA
Hypsiglena chlorophaea Cope, 1860, Pr. Phil. Ac., 247. *H. ochrorhynchus*
var. *chlorophaea* Cope, 1866, Pr. Phil. Ac., 304. *H. ochrorhynchus*
subsp. *chlorophaea* Cope, 1875, Checklist, 38. *H. ochrorhynchus chloro-*
phaea Coues, 1875, Wheeler's Rep., V, 622.
More slender. Colors darker; spots black. Ventrals 167. Subcau-
dals 55 pairs.
Arizona to Sonora.
- CALAMARINAE 81
Calamariidae Gthr., 1858, Cat., 2; Jan, 1862, Arch. Zool., I.
- FICMIA 81
Gray, 1849, Cat. Serp., 80.
- FICMIA FRONTALIS 82
Tolaca frontalis Cope, 1864, Pr. Phil. Ac., 167.
Colima, Mexico.
- FICMIA CANA 83
Gyalopion canum Cope, 1860, Pr. Phil. Ac., 243, 310, — 1875, Checklist,
36; Coues, 1875, Wheeler's Rep., V, 624, pl. 18, f. 2.
Arizona.
- FICMIA NASUS 83
Conopsis nasus Gthr., 1858, Cat., 6. California.
- FICMIA LINEATA
Tolaca lineata (Kenn.) Bd., 1859, Mex. Bound., II, Rept., 23, pl. 21, f. 2,
— 1859, P. R. R. Rep., X, pl. 35, f. 8; Cope, 1860, Pr. Phil. Ac., 241.
Oxyrhina de Filippii Jan, 1862, Arch. Zool., II, pp. 54, 61, — 1863,
Sist., 41. *Stenorhina de Filippii* Jan, 1857, Ind. Rett. Mus. Milan, 48.
Stenorhina varians Jan, 1857, Ind. Rett. Mus. Milan, 48. *Oxyrhina*
varians Jan, 1862, Arch. Zool., II, pp. 54, 60, — 1863, Sist., 41.
Body stout; head short, indistinct, pointed; snout acute, promi-
nent; crown arched; tail short, about one sixth of total. Rostral

OPHIDIA—Continued.

- large, turned back on the top of the snout. Frontal large, anterior angle separating the prefrontals. Nasal entire. No loreal. Oculars 1-2. Labials 7, first small. Infralabials 6. Scales smooth, in 17 rows, outer row broader than long. Anal bifid. Subcaudals in two rows. Uniform light-brownish ash above, with three imperfect longitudinal stripes of blackish, each on a single row of scales.
- Mexico.
- FICMIA MACULATA 84
Oxyrhina maculata Jan, 1862, Arch. Zoöl., II, 54, 61, — 1863, Sist., 41, — 1876, Icon., livr. 48, pl. 2, f. 2-4.
- Mexico.
- FICMIA VARIEGATA 84
 Cope, 1866, Pr. Phil. Ac., 126. *Amblymetopon variegatum* Gthr., 1858, Cat. 7.
- Mexico.
- FICMIA OLIVACEA 84
 Gray, 1849, Cat. Serp., 80; Cope, 1863, Pr. Phil. Ac., 126, — 1866, Pr. Phil. Ac., 132; Peters, 1869, Mb. Berl. Akad., 875; Steind., 1870, Sitzungsab. Ak. Wien, 344, pl. 6.
- Mexico.
- CHEILORHINA 85
 Jan, 1862, Arch. Zoöl., II, 57.
- CHEILORHINA VILLARSHI 85
 Jan, 1862, Arch. Zoöl., II, 57, — 1863, Sist., 41, — 1876, Icon., livr. 48, pl. 1, f. 5.
- Western Mexico.
- STENORHINA 85
 Dum. Bibr., 1854, Erp., VII, 865.
- STENORHINA FREMUSVILLEI 85
 D. & B., 1854, Erp., VII, 868. *S. degenhardtii* var. *fremusvillei* Jan, 1862, Arch. Zoöl., II, 65, — 1863, Sist., 41. ? *Bergenia mexicana* Steind., 1867, Novara Exp., Rept., 92.
- Central America and Mexico.
- STENORHINA QUINTQUELINEATA 86
Microphis quinquelineatus Hall., 1854, Pr. Phil. Ac., 97. *S. degenhardtii* var. *quinquelineata* Jan, 1862, Arch. Zoöl., 55, 65. *Stenorhina quinquelineata* Cope, 1869, Pr. Phil. Ac., 243.
- Mexico.
- TANILLA 86
 Baird & Girard, 1853, Serp., 131.
- TANILLA GRACILIS 87
 B. & G., 1853, Serp., 132, 161; Cope, 1866, Pr. Phil. Ac., 126, — 1875, Checklist, 35, — 1880, Bull. 20, U. S. Mus., 20; Bd., 1859, Mex. Bound., II, Rept., 23. *Homonolocanion gracile* Jan, 1862, Arch. Zoöl., II, 50, — 1863, Sist., 39, — 1866, Icon., livr. 15, pl. 2, f. 1.
- Texas.

OPHIDIA—Continued.

- Var. HALLOWELLII
Tantilla Hall, 1856, Pr. Phil. Ac., 246. *T. Hallowellii* Cope, 1860, Pr. Phil. Ac., 77, — 1866, Pr. Phil. Ac., 126, — 1875, Checklist, 55.
 Oculars 1. No loreal. Frontal broad, short, hexagonal. Temporals 2. Labials 6, (third and fourth in orbit, fourth to sixth large. Infralabials 6, fourth largest. Scales smooth, in 15 rows, outer larger. Ventrals 130. Anal entire. Subcaudals 42 pairs. Light brown or olive above; light beneath. A vertebral narrow dark-colored line from the back of the head.
 Kansas.
- Var. CALAMARINA 87
Tantilla calamarium Cope, 1866, Pr. Phil. Ac., 320. *T. bimaculata* Cope, 1875, Rept. Costa Rica, 143.
 Guadalupe, Mexico.
- TANTILLA NIGRICEPS 88
 Kennicott, 1860, Pr. Phil. Ac., 328; Cope, 1866, Pr. Phil. Ac., 126, — 1875, Checklist, 55, — 1889, Bull. 20, U. S. Mus., 20; Coles, 1875, Wheeler's Rep., V, 626. *Scolocophis fusciceps* Cope, 1860, Pr. Phil. Ac., 371, — 1861, Pr. Phil. Ac., 74.
 Texas and New Mexico.
- TANTILLA CORONATA 88
 Baird & Girard, 1853, Serp., 131; Bd., 1859, P. R. R. Rep., X, pl. 33, f. 96; Cope, 1866, Pr. Phil. Ac., 126, — 1875, Checklist, 55, — 1877, Pr. Am. Phil. Soc., 65. *Hemalocration melanocepalum* Jan, 1862, Arch. Zool., II, 50, 51 (mixed).
 Gulf States.
- Var. WAGNERI 88
Hemalocration wagneri Jan, 1862, Arch. Zool., II, 50, 51, — 1863, Sist., 40, — 1866, Icon., livr. 15, pl. 2, f. 3.
 Florida.
- TANTILLA PLANICEPS 89
 Cope, 1875, Checklist, 55 (name). *Coluber planiceps* Blainv., 1835, Nouv. Ann. Mus. Paris, IV, 294, pl. 27, f. 3; B. & G., 1853, Serp., 154. *Hemalocration planiceps* Dum., 1852, Mem. Ac. Sci., XXIII, 490; D. & B., 1854, Exp., VII, 857; Jan, 1862, Arch. Zool., II, 51, 55, — 1863, Sist., 40, — 1866, Icon., livr. 15, pl. 2, f. 2.
 California.
- ELAPOMORPHUS 89
 (Wiegmann) Fitz., 1843, Syst. Rept., 25.
- ELAPOMORPHUS MEXICANUS 89
 Guenther, 1862, Ann. Mag. Nat. Hist., pl. 9, f. 1 (extr. p. 6).
 Mexico.
- CONTIA 90
 Baird and Girard, 1853, Serp., 110.

OPHIDIA—Continued.

- (SONORA.)
- CONTIA SEMIANNULATA 90
Sonora seminannulata B. & G., 1853, Serp., 117; Btl., 1859, Mex. Bound., II, Rept., 21, pl. 19, f. 3, — P. R. R. Rep., X, pl. 33, f. 88; Cope, 1875, Checklist, 36.
 Sonora.
- CONTIA OCCIPITALIS 91
Rhinostoma occipitale Hall, 1854, Pr. Phil. Ac., 95, — 1856, Pr. Phil. Ac., 311, — 1859, P. R. R. Rep., X, Rept., 15. *Lamprosona occipitale* Btl., 1859, Mex. Bound., II, Rept., 21, pl. 21, f. 1, — P. R. R. Rep., X, pl. 35, f. 67. *Chionactis occipitalis* Cope, 1860, Pr. Phil. Ac., 241, — 1875, Checklist, 35.
 Arizona.
- VAR. ANNULATA 91
Lamprosona annulatum Btl., 1859, Mex. Bound., II, 22. *Chionactis occipitalis* subsp. *annulata* Cope, 1875, Checklist, 36.
 Arizona.
- CONTIA ISOZONA 92
 Cope, 1866, Pr. Phil. Ac., 304, — 1875, Checklist, 36; Yarrow, 1875, Wheeler's Exp., V, 537, pl. 18, f. 1.
 Arizona to Utah.
- (PROCNURA.)
- CONTIA AEMULA (? position) 92
Procnura aemula Cope, 1879, Pr. Am. Phil. Soc., 262.
 Mexican Plateau.
- (CONTIA.)
- CONTIA PYGAFA 93
 Cope, 1871, Pr. Phil. Ac., 223, — 1875, Checklist, 36, — 1877, Pr. Am. Phil. Soc., 65.
 Florida.
- CONTIA MITIS 93
 Baird & Girard, 1853, Serp., 110; Btl., 1859, P. R. R. Rep., X, pl. 36, f. 7; Cope, 1860, Pr. Phil. Ac., 251, — 1861, Pr. Phil. Ac., 74. *Ablabes purpurocauda* Gthr., 1858, Cat., 245. *Homonosoma mile* Jan, 1862, Arch. Zool., 11, 33, 35, — 1863, Sist., 36, — 1865, Icon., livr. 13, pl. 4, f. 1.
 California.
- CONTIA EPISCOPA 94
 Cope, 1860, Pr. Phil. Ac., 251, — 1875, Checklist, 36, — 1880, Bull. 20, U. S. Mus., 20. *Lamprosona episcopum* Kenn., 1859, Mex. Bound., II, Rept., 22, pl. 8, f. 2. ? *Contia episcopa* subsp. *torquata* Cope, 1880, Bull. 20, U. S. Mus., 20. *C. episcopa* subsp. *episcopa* Cope, l. c., 20. *Homonosoma episcopum* Jan, 1862, Arch. Zool., 11, 33, 35, — 1863, Sist., 37, — 1865, Icon., livr. 13, pl. 4, f. 2.
 Texas.

OPHIDIA—Continued.

- LODIA**
- Baird & Girard, 1853, Cat. N. A. Serp., 116.
- Head distinct. Frontal divided transversely. Nasal in two parts. Loreal entering orbit, fused with the lower antecular. Oculars 1—2. Mentals one pair. Eyes rather large, pupil round. Scales smooth. Anal bifid. Subcaudals in two rows.
- LODIA TENUIS**
- B. & G., 1853, Serp., 116; Bd., 1859, P. R. R. Rep., X, pl. 36, f. 8; Cope, 1875, Checklist, 36. *Calamaria tenuis* B. & G., 1852, Pr. Phil. Ac., 176.
- Body slender, subcylindrical; tail short, conical, tapering. Frontal hexagonal, lateral borders parallel. A small subelliptical shield between prefrontals and frontal. Internasals subtriangular, about half as large as prefrontals. Rostral broad. Nasal in two parts, nostril between. Loreal large, elongate, entering the orbit, fused with lower antecular. Oculars 1—2. Anterior temporal largest, elongate. Labials 6. Infralabials 6, fourth largest. One pair of mentals. Scales rather large, smooth, in 15 rows, outer broader. Ventrals 150. Anal entire. Subcaudals 33 pairs. Brown above, bluish laterally, with a longitudinal stripe of lighter on each flank. Belly lighter, bases of scutes bluish.
- Oregon and Washington Territories.
- NINIA** 94
- B. & G., 1853, Serp., 49. *Streptophorus* D. & B., 1854, Erp., VII, 514.
- NINIA LIEBMANNI** 95
- Chersodromus liebmanni* Reinh., 1860, Vid. Medd. Kjobenh., 35, pl. IV, f. 10, 11; Cope, 1861, Pr. Phil. Ac., 302; Jan, 1862, Arch. Zool., II, 25, — 1865, Icon., livr. 12, pl. 3, f. 2.
- Vera Cruz, Mexico.
- NINIA SERAE** 95
- Cope, 1860, Pr. Phil. Ac., 340. *Streptophorus serae* D. & B., 1854, Erp., VII, 513; Gthr., 1858, Cat., 16; Jan, 1862, Arch. Zool., II, 26, 27, — 1863, Sist., 34, — 1865, Icon., livr. 12, pl. 3, f. 4; Mart., 1856, Mus. Berl., 28.
- Mexico.
- NINIA COLLARIS** 95
- Streptophorus serae* var. *collaris* Jan, 1865, Icon., livr. 12, pl. 3, f. 6.
- Mexico.
- NINIA SIEROLDI** 96
- Elopoites sieroldi* Jan, 1862, Arch. Zool., II, 21, — 1863, Sist., 33, — 1865, Icon., livr. 12, pl. 1, f. 4.
- Mexico.
- NINIA DIADEMATA** 96
- B. & G., 1853, Serp., 49; Bd., 1859, P. R. R. Rep., X, pl. 27, f. 37. *Tropidonotus dimidiatus* Cope, 1861, Pr. Phil. Ac., 297. *Streptophorus bifasciatus* D. & B., 1854, Erp., VII, 520; Cope, 1860, Pr. Phil. Ac., 77; Jan, 1862, Arch. Zool., II, 26, — 1863, Sist., 34, — 1865, Icon., livr. 12, pl. 3, f. 3.
- Mexico.

OPHIDIA—Continued.

	Southeastern States.	
VIRGINIA		96
B. & G., 1853, Serp., 127		
VIRGINIA STRIATULA		97
<i>Coluber striatulus</i> Linn., 1766, Syst., I, 375; Gmel., 1788, Syst. Linn., I, 1087; Daub., 1784-92, Encycl. Meth., II, 684; Latr., 1802, Rept., IV, 84; Daud., 1803, Rept., VII, 200; Merr., 1820, Syst., 118; Harl., 1827, Jour. Phil. Ac., V, 354. — 1835, Med. Res., 117. <i>Calamaria striatula</i> Schleg., 1837, I, 133, II, 43; Holbr., 1842, Herp., III, 123, pl. 29. <i>Haldia striatula</i> B. & G., 1853, Serp., 122; Mart., 1856, Mus. Berl., 23; Bl., 1859, P. R. R. Rep., X, pl. 32, f. 91; Cope, 1875, Checklist, 35, — 1880, Bull. 20, U. S. Mus., 29. <i>Conocophidius striatulus</i> D. & B., 1854, Exp., VII, 140; Gthr., 1858, Cat., 17; Jan., 1862, Arch. Zool., II, 24, — 1863, Sist., 34, — Icon., livr. 12, pl. 3, f. 1.		
	Virginia to Texas.	
VIRGINIA INORNATA sp. n.		97
	Texas.	
VIRGINIA ELEGANS		98
Kenn., 1859, Pr. Phil. Ac., 99; Jan., 1862, Arch. Zool., II, 24, — 1863, Sist., 34, — 1865, Icon., livr. 12, pl. 2, f. 6; Cope, 1880, Bull. 20, U. S. Mus., p. 20.		
	Southern Illinois.	
VIRGINIA VALERIAE		98
B. & G., 1853, Serp., 127; Bl., 1859, P. R. R. Rep., X, pl. 33, f. 94; Jan., 1862, Arch. Zool., II, 24, — 1863, Sist., 34, — 1865, Icon., livr. 12, pl. 2, f. 5; Cope, 1880, Bull. 20, U. S. Mus., p. 20.		
	Maryland to Georgia and Illinois.	
VAR. HARPERTI		99
<i>Corphophis harperti</i> D. & B., 1854, Exp., VII, 135; Cope, 1880, Bull. 20, U. S. Mus., 20. <i>Virginia harperti</i> Cope, 1875, Checklist, 35. <i>V. valerie</i> Jan., 1863, Sist., 34. <i>Cluta harperti</i> Mart., 1856, Mus. Berl., 23.		
	Georgia to Texas.	
CARPHOPHIS		99
Gervais, 1843, D'Orb. Diet. d'Hist. Nat., III, 191; D. & B., 1854, Exp., VII, 133. <i>Cluta</i> B. & G., 1853, Serp., 129		
CARPHOPHIS STRAMINEA		99
<i>Chilomeniscus stramineus</i> Cope, 1860, Pr. Phil. Ac., 33, — 1861, Pr. Phil. Ac., 302, — 1875, Checklist, 35.		
	Lower California.	
CARPHOPHIS CINCTA		100
<i>Chilomeniscus cinctus</i> Cope, 1861, Pr. Phil. Ac., 303, — 1875, Checklist, 35. <i>C. cphippicus</i> Cope, 1867, Pr. Phil. Ac., 85, — 1875, Checklist, 35.		
	Guaymas, Sonora.	
CARPHOPHIS HELENÆ		100
<i>Cluta helenæ</i> Kenn., 1859, Pr. Phil. Ac., 100. <i>Corphophis helenæ</i> Cope, 1875, Checklist, 34. <i>Corphophis amonia</i> var. <i>helenæ</i> Jan., 1862, Arch. Zool., 23, — 1863, Sist., 33.		
	Illinois to Mississippi.	

OPHIDIA—Continued.

- CARPHOPIUS AMOENA ("Ground Snake"). 100
 Gervais, 1843, D'Orb. Dict. d'Hist. Nat., III, 191; D. & B., 1854, Erp., VII, 131; Gthr., 1858, Cat., 18; Cope, 1860, Pr. Phil. Ac., 79; Jan., 1862, Arch. Zool., 23, — 1863, Sist., 33, — 1865, Icon, livr. 12, pl. 2, f. 4; Allen, 1869, Pr. Bost. Soc., 182. *Cuber amoenus* Say, 1825, Jour. Phil. Ac., IV, 237; Harl., 1827, Jour. Phil. Ac., 355, — 1835, Med. Res., 118; Storer, 1839, Rept. Mass., 226; Linsley, 1844, Am. Jour., vol. 45, p. 43. *Brachyorrhos amoenus* Holbr., 1842, Herp., III, 115, pl. 27. *Corphophiops vermiformis* Gerv., 1843, Dict. Hist. Nat., III, 191. *C. amoenus* Cope, 1875, Checklist, 34. *Calamaria amoena* Schleg., 1837, Essai, I, 139, II, 31, pl. 1, f. 19-20. *Cluta amoena* B. & G., 1853, Serp., 129; Mart., 1856, Mus. Berl., 23; Baird, 1854, Serp. N. Y., 25, — 1859, P. R. R. Rep., X, pl. 33, f. 95.
 Massachusetts to Illinois and southward.
- Var. VERMIS 101
Cluta vermīs Kenn., 1859, Pr. Phil. Ac., 99. *Corphophiops vermīs* Cope, 1875, Checklist, 34.
 Missouri and southward.
- GEOPHIS 101
 Wagler, 1830, Syst. Amph., 342.
- GEOPHIS SEMIDOLIATUS 102
 Peters, 1859, Mb. Berl. Akad., 275. *Rhabdosoma semidoliatum* D. & B., 1854, Erp., VII, 93. *Rhabdosoma semidoliatum* Gthr., 1858, Cat., 10; Mart., 1856, Mus. Berl., 23; Cope, 1860, Pr. Phil. Ac., 78. *Catostoma semidoliatum* Cope, 1860, Pr. Phil. Ac., 339. *Elapoides semidoliatum* Jan., 1862, Arch. Zool., 21, 22, — 1863, Sist., 33, — 1865, Icon., livr. 12, pl. 2, f. 1.
 Mexico.
- GEOPHIS BICOLOR 102
 Guenther, 1868, Ann. Mag. Nat. Hist., 413.
 Valley of Mexico.
- GEOPHIS LATIFRONTALIS sp. n 103
 San Luis Potosi, Mexico.
- GEOPHIS LIPPIENS 103
Sympholis lippicus Cope, 1861, Pr. Phil. Ac., 524
 Guadalajara, Mexico.

- TOXICOPHIDIA** (Venomous Snakes) 104
- PROTEROGLYPHA** 104
- CONOCERCA** 104
- ELAPIDAE** 104
- Gthr., 1858, Cat., 209.
- ELAPS** 104
- Schneider, 1801, Hist. Amph., II, 289.
- ELAPS FULVIUS** ("Harlequin Snake") 105
- Cuvier, 1817, Regn. Anim., II, 84; Fitz., 1826, Neue Class., 61; Holbr., 1833, Herp., II, 87, pl. 18. — 1842, Herp., III, 49, pl. X; B. & G., 1853, Serp., 21; D. & B., 1854, Erp., VII, 1215; Gthr., 1858, Cat., 235; Bd., P. R. R. Rep., X, pl. 25, f. 15; Cope, 1859, Pr. Phil. Ac., 344, — 1875, Checklist, 34, — 1880, Bull. 20, U. S. Mus., 23; Jan, 1859, Rev. & Mag. Zool. (Extr., p. 6), — 1863, Sist., 113, — 1872, Icon., livr. 42, pl. 2, f. 1; Dum. Boc., 1874, Miss. Sci. Mex., pl. 23; Audubon, Ornithol., I, pl. 41. *Coluber fulvius* Linn., 1766, Syst., I, 381; Gmel., 1788, Syst. Linn., I, 1104; Latr., 1802, Rept., IV, 140; Shaw, 1802, Zool., III, 469; Daud., 1803, Rept., VI, 300; Say, 1825, Am. Jour. Sci., I, 262. *Vipera fulva* Harl., 1827, Jour. Phil. Ac., 364, — 1835, Med. Res., 127. *Le Noir et fauve* LaC., 1789, Quad. Ovip. et Serp., II, 299. ? *Elaps fulvius* var. *leypostema* Jan. 1863, Sist., 113.
- Southern States east of the Mississippi.
- Var. NIGROINCTUS** 106
- Elaps nigroinctus* Grd., 1854, Pr. Phil. Ac., 226, — 1855, Gilliss' Exp., II, 210; Cope, 1859, Pr. Phil. Ac., 345; Sumichrast, 1881, Bull. Soc. Zool. de France, 184. *E. divaricatus* Hall., 1855, Jour. Phil. Ac., 36. *E. fitzingeri* Jan, 1859, Rev. & Mag. Zool., pp. 5 and 10. *E. fulvius* var. *fitzingeri* Jan, 1863, Sist., 113, — 1872, Icon., livr. 42, pl. 2, f. 3.
- Central America to Mexico.
- Var. AFFINIS** 106
- Jan, 1863, Sist., 113, — 1872, Icon., livr. 42, pl. 1, f. 2. *Elaps affinis* Jan, 1859, Rev. & Mag. Zool., pp. 6 and 14, pl. B, f. 2.
- Mexico.
- Var. BIPUNCTIGER**
- Elaps bipunctiger* D. & B., 1854, Erp., VII, 1227. *Elaps distans* Kenn., 1860, Pr. Phil. Ac., 338; Cope, 1875, Checklist, 34. *E. ornaticissimus* Jan, 1859, Rev. & Mag. Zool. (Extr. pp. 5 and 10, pl. A, f. 3), — 1863, Sist., 113, — 1872, Icon., livr. 42, pl. 1, f. 1.
- Scales unicolor. Throat white, perhaps yellowish in life. Vertex black, followed by a collar of white or red. The specimen described had forty-three circles of white or red. The rings were narrow, occupying but two of the ventrals. Muzzle without a black spot. The name bipunctiger was given on account of a couple of light-colored spots in the black of the crown. Tip of tail black.
- Mexico; Florida.

TOXICOPHIDIA—Continued.

- Var. **TENER**
Elaps tener B. & G., 1853, Serp., pp. 22, 156; Bl., 1859, P. R. R. Rep., X, pl. 25, f. 16. *E. tristis* B. & G., 1853, Serp., 23; Bl., 1859, P. R. R. Rep., X, pl. 25, f. 17. *E. fulvius* subsp. *tener* Cope, 1875, Checklist, 34.
 Anterior portion of head black. A yellow occipital ring. Black rings fifteen in number. Red rings broadest. Yellow rings about half as wide as the black, and twice as many as either black or red. Tip of tail black. Ventrals 224 to 227. Subcaudals 26 to 38.
 Texas.
- Var. **APIATUS** 106
 Jan, 1863, Sist., 113. *Elaps apiatus* Jan, 1859, Rev. & Mag. Zool. (Extr., pp. 6, 11, pl. A, f. 4).
 Vera Cruz.
- Var. **EPISTEMA** 106
Elaps epistema D. & B., 1854, Erp., VII, 1222; Gthr., 1858, Cat., 236; Jan, 1859, Rev. & Mag. Zool. (Extr., p. 6), — 1863, Sist., 113.
 Mexico.
- Var. **DIATEMA** 107
 Jan, 1863, Sist., 113. *Elaps diatema* D. & B., 1854, Erp., VII, 1222; Sumichr., 1881, Bull. Soc. Zool. de France, 185.
 Mexico.
- Var. **CEREBRIPUNCTATUS** 107
Elaps cerebripunctatus Pet., 1869, Mb. Berl. Akad., 877.
 Pueblo, Mexico.
- ELAPS EURYXANTHUS** 107
 Kenn., 1860, Pr. Phil. Ac., 337; Cope, 1861, Pr. Phil. Ac., 296, — 1866, Pr. Phil. Ac., 307, — 1875, Checklist, 34; Coues, 1875, Wheeler's Rep., V, 611.
 Arizona to Mexico.
- ELAPS LATICOLLARIS** 107
Elaps maregravii var. *laticollaris* Pet., 1869, Mb. Berl. Akad., 877.
 "Pueblo, Mexico."
- ELAPS DECORATUS** 108
 Jan, 1859, Rev. & Mag. Zool. (Extr., pp. 7, 14, pl. B, f. 5), — 1863, Sist., 114, — 1872, Icon., livr. 42, pl. 6, f. 4.
 Mexico.
- ELAPS ELEGANS** 108
 Jan, 1859, Rev. & Mag. Zool. (Extr., pp. 6, 13, pl. B, f. 1), — 1863, Sist., 114, — 1872, Icon., livr. 42, pl. 5, f. 2; Cope, 1861, Pr. Phil. Ac., 295.
 Mexico.
- PLATYCERCA** 109
- HYDROPHIDAE** 109
- PELAMIS** 109
 Daudin, 1803, Rept., VII, 357.

TOXICOPHIDIA—Continued.

PELAMIS PLATURA

Anguis platyce Linn., 1766, Syst., I, 391. *Hydrophis platura* Latr., 1802, Rept., IV, 197. *Hybros bicolor* Schneid., 1801, Amph., I, 242; Shaw, 1802, Zool., III, 566. *Peltamis bicolor* Daud., 1803, Rept., VII, 366, pl. 89; Merr., 1820, Syst., 138; Gray, 1842, Zool. Misc., 60, — 1849, Cat. Snakes, 41; D. & B., 1854, Erp., VII, 1335; Cope, 1859, Pr. Phil. Ac., 347, — 1875, Jour. Phil. Ac., 95, 148; Jan, 1859, Rev. & Mag. Zool., Extr., p. 24; Fischer, 1856, Abhandl. Naturw. Hamb., 61; Gthr., 1864, Rept. Brit. India, 382; Trosch., 1865, Mueller's Wirbelth. Mex., 78. *Hydrophis bicolor* Mart., 1853, Mus. Berl., 32; Jan, 1863, Sist., 109, — 1872, Icon., livr. 40, pl. 2, f. 3-4, pl. 3, f. 2. *H. peltamis* Schleg., 1837, Essai, I, 187, II, 508, pl. 18, f. 13-15. *H. variegata* Schleg., 1838, Fauna Jap. Rept., tab. 8. *Peltamis ornata* Gray, 1849, Cat. Snakes, 43.

Seba, 1735, Thesaur., II, pl. 77, f. 2; Vosmaer, 1774, Nat. Hist. Plattstaart Slang uit Mexico; Russell, 1796, Ind. Serp., I, 47, pl. 41, — 1801, vol. II, pl. 12.

West coast of Mexico and Central America to the East Indies and China.

SOLENOGLYPHA 110

BOTHROPERA 110

CROTALIDAE 110

Crotali Oppel, 1811, Sur la Class. Rept., Mem. I, 44, — Ann. Mus., 387. *Crotaloidea* Fitz., 1826, Neue Class., 62. *Crotalidae* Gray, 1842, Zool. Misc., 47, — 1849, Cat. Serp., 3.

CROTALUS 110

Crotalus Linn., 1754, Mus. Ad. Fridr., 39. *Crotalus* Linn., 1758, Syst., I, 214, — 1766, Syst., I, 372 (incl. *C. miliaris*); Latr., 1801, Rept., III, 166; Daud., 1803, Rept., V, 297; Cuv., 1817, Regn. An., II, 77, — 1826, R. An., I, 412; Merr., 1820, Amph., 156; Fitz., 1826, Neue Class., 34, 63; Wagl., 1830, Amph., 176; Gray, 1849, Cat. Serp., 19; B. & G., 1853, Serp., 1; D. & B., 1854, Erp., VII, 1453; Cope, 1859, Pr. Phil. Ac., 336. *Crotalophocetus* Linn., 1749, Amoen. Acad., I, pp. 297, 500, — 1756, Syst. Nat., Ed. 9, p. 35; Gronow, 1756, Hist. Amph., Gen., 45, — 1763, Zoophylac., p. 26, sp. 137-138. *Crotalisona* Linn., 1735, Syst. Nat., Ed. 1; Laurent, 1768, Syn. Rept., 92; Cope, 1861, Mitch. Res., 119. *Crotalinus* Raf., 1818, Am. Month. Mag., III, 446, IV, 41. *Cropsophus* Wagler, 1830, Amph., 176; Gray, 1849, Cat. Serp., 18; Fitz., 1843, Syst., 29. *Urocrotalon* Fitz., 1843, Syst., 29.

CROTALUS DERRISSI 111

Crotalus derrissi Linn., 1758, Syst., I, 214, — 1766, Syst., I, 372; Gmel., 1788, Syst. Linn., I, 1081; Bonn., 1790, Ophiol., 2, pl. 3, f. 4; Cuv., 1817, R. An., II, 78, — 1826, R. An., I, 413; Merr., 1820, Syst., 156; Boie, 1827, Isis, 562; Griffith, 1831, Cuv. An. King., IX, 267; LeC., 1853, Pr. Phil. Ac., 416; Cope, 1859, Pr. Phil. Ac., 337; Yarrow, 1875, Wheeler's Rep., V, 532; Sumichr., 1881, Bull. Soc. Zool. de France, 185. *Crotalisona derrissa* Cope, 1861, Pr. Phil. Ac., 292, — 1861,

SOLENOGLYPHA—Continued.

Mitch. Res., 120, — 1866, Pr. Phil. Ac., 308. *Crotalus horridus* Latr., 1801, Rept., III, 186; Daud., 1803, Rept., V, 311; Max., 1824, Beitr. Bras., I, 435, — 1827, Abbild. Bras., lief. 11, pl. 4; Fitz., 1826, Neue Class., 63; Wagl., 1830, Amph., 176; Schleg., 1837, Essai, II, 561, pl. 20, f. 12-14; Gray, 1849, Cat., 20 (mixed); D. & B., 1854, Erp., VII, 1472, pl. 84 bis, f. 2; Mart., 1856, Mus. Berl., 34; Jan., 1859, Rev. & Mag. Zoöl., Extr., p. 28, — 1863, Sist., 124, — 1874, Icon., livr. 46, pl. 3, f. 1-2. *Caudisoma terrificum* Laur., 1768, Syn. Rept., 93; Cope, 1861, Mitch. Res., 120, — 1866, Pr. Phil. Ac., 308. *Crotalus terrificus* Yarrow, 1875, Wheeler's Rep., V, 532. *Crotalus cascavella* Wagl., 1824, Spix. Bras. Serp., 69, pl. 24, — 1830, Syst., 176. *Crotalus simus* Latr., 1801, Rept., III, 202; Daud., 1803, Rept., V, 321. *Crotalus longlingii* Humb., 1833, Recueil d'Obs. Zoöl., p. 6; Cope, 1861, Mitch. Res., 120. *Le Boitiquet*, p. 390, and *Le durissus*, p. 423, LaC., 1739, Serp., II, text only.

Brazil to Mexico.

Var. MOLOSSUS 113

Crotalus molossus B. & G., 1853, Serp., 10; D. & B., 1854, Erp., VII, 1482; Bd., 1859, Mex. Bound., II, Rept., 14, pl. 3, — 1859, P. R. R. Rep., X, Rept., pl. 24, f. 5; Cope, 1859, Pr. Phil. Ac., 338, — 1875, Checklist, 33; Yarrow, 1875, Wheeler's Rep., pp. 531, 533; Coles, 1875, Wheeler's Rep., V, 605. *Caudisoma molossus* Cope, 1861, Mitch. Res., 124, — 1866, Pr. Phil. Ac., 308. *Crotalus ornatus* Hall., 1854, Pr. Phil. Ac., 192, — 1859, P. R. R. Rep. (Parke's), X, pl. 2, pl. 24, f. 5, — Cope, 1859, Pr. Phil. Ac., 338.

Northern Mexico; Arizona; New Mexico.

Var. BASILISCUS

Caudisoma basilisca Cope, 1864, Pr. Phil. Ac., 166, — 1866, Pr. Phil. Ac., 308. *Crotalus basiliscus* Yarrow, 1875, Wheeler's Rep., V, 532.

Muzzle plates two pairs, a third pair subdivided. Rostral higher than wide, in contact with nasals and internasals. Labials 14, separated from suborbitals by two and three rows. Scales in 29 rows, outer largest, outer three smooth. Ventrals 199. Subcaudals 24. Pale yellowish-brown, head and tail darker. Markings on entire scales. Dorsal rhombs about thirty; five obscure rings on the tail. Two elongated occipital spots. No lines on the neck. Labials yellowish.

Western Mexico.

CROTALUS ADAMANTEUS (Diamond Rattlesnake) 112

Crotalus adamantus Beauv., 1799, Trans. Am. Phil. Soc., IV, 368; Holbr., 1842, Herp., III, 17, pl. 2; DeK., 1842, N. Y. Fauna, Rept., 57; B. & G., 1853, Serp., 3; Bd., 1859, P. R. R. Rep., X, pl. 24, f. 2; LeC., 1853, Med. Jour., IX, 664; Jan., 1859, Rev. Mag. Zoöl., Extr., p. 28, — 1863, Sist., 123, — 1874, Icon., livr. 46, pl. 2, f. 2; Yarrow, 1875, Wheeler's Rep., V, 534; Cope, 1877, Pr. Am. Phil. Soc., 64. *Crotalus terrificus* Cope, 1859, Pr. Phil. Ac., 337; LeC., 1853, Pr. Phil. Ac., 419. *Caudisoma adamantea* Cope, 1861, Mitch. Res., 121, — 1866,

SOLENOGLYPHA—Continued.

Pr. Phil. Ac., 309. *Crotalus adamanteus* subsp. *adamanteus* Cope, 1875, Checklist, 33. *C. rhombifer* Latr., 1801, Rept., III, 197; Daud., 1803, Rept., V, 323, pl. 60, f. 22, 23, pl. 69, f. 2; Merr., 1820, Syst., 157; D. & B., 1854, Erp., VII, 1470; Mart., 1856, Mus. Berl., 34; Duges, 1879, La Natureza, IV, 22. *C. durissus* Shaw, 1802, Zool., III, pt. 2, 333, pl. 89. *C. horridus* Latr., 1801, Rept., III, 199.

Texas to the Carolinas.

VAR. SCUTULATUS 113

Caulisoma scutulatus Kenn., 1861, Pr. Phil. Ac., 207; Cope, 1866, Pr. Phil. Ac., 307, 309. *Crotalus scutulatus* Yarr., 1875, Wheeler's Rep., V, 533. *Caulisoma adamantea* subsp. *scutulata* Cope, 1875, Checklist, 33. *Caulisoma adamantea scutulata* Cones, 1875, Wheeler's Rep., V, 607.

Arizona; Mexico.

VAR. ATROX 113

Crotalus atrox B. & G., 1853, Serp., pp. 5, 156; D. & B., 1854, Erp., VII, 1482; Bd., 1859, Mex. Bound., II, 14, pl. 1, — 1859, P. R. R. Rep., X, Whipples, 39, pl. 24, f. 3; Cope, 1859, Pr. Phil. Ac., 337; Yarr., 1875, Wheeler's Rep., V, 534; Duges, 1879, La Natureza, IV, 26. *Caulisoma atrox* Cope, 1861, Mitch. Res., 121, — 1866, Pr. Phil. Ac., 309. *Caulisoma adamantea atrox* Cones, 1875, Wheeler's Rep., V, 607. *Crotalus adamanteus atrox* Cope, 1880, Bull. 20, U. S. Mus., 24. *Crotalus adamanteus* subsp. *atrox* Cope, 1875, Checklist, 33; Yarr., 1875, Wheeler's Rep., V, 529. *Crotalus adamanteus* Jan, 1874, Icon., livr. 46, pl. 2, f. 1. *C. adamanteus* var. *atrox* Jan, 1859, Rev. Mag. Zool., Extr., p. 28, — 1863, Sist., 123. *Caulisoma atrox* var. *sonoricensis* Kenn., 1861, Pr. Phil. Ac., 205. *Caulisoma atrox sonoricensis* Cope, 1861, Pr. Phil. Ac., 292. *Crotalus adamanteus* var. *sonoricensis* Jan, 1863, Sist., 124.

Texas to Mexico.

CROTALUS CONFLUENTUS 114

Crotalus confluentus Say, 1823, Long's Exp., II, 48; B. & G., 1853, Serp., 8, — 1854, Macey's Exp., 188, pl. 1; D. & B., 1854, Erp., VII, 1475; Bd., 1859, Mex. Bound., II, Rept., 14, — 1859, P. R. R. Rept., X, Whipples, 40, pl. 24, f. 4; Cooper, 1860, P. R. R. Rep., XII, pt. 2, 295; Hall, 1856, Pr. Phil. Ac., 250; Cope, 1859, Pr. Phil. Ac., 337; Yarrow, 1875, Wheeler's Rep., V, 530, 533; Cope, 1875, Checklist, 33, 1880, Bull. 20, U. S. Mus., 24; Cones & Yarrow, 1878, Hayd. Rep., IV, 263. *Caulisoma confluenta* Cope, 1861, Mitch. Res., 122, — 1866, Pr. Phil. Ac., 307, 309; Allen, 1874, Pr. Bost. Soc., 309; Cones, 1875, Wheeler's Rep., V, 604. *Crotalus lecontei* Hall, 1851, Pr. Phil. Ac., 189, — 1859, P. R. R. Rep., X, Williamson's, 18, — 1853, Sitgreave's Exp., 139. *Caulisoma lecontei* Cope, 1861, Mitch. Res., 121; Hayd., 1862, Trans. Am. Phil. Soc., 177. *Caulisoma confluenta* var. *lecontei* Cope, 1866, Pr. Phil. Ac., 307. *Crotalus adamanteus* var. *confluentus* Jan, 1859, Rev. Mag. Zool., Extr., p. 28, — 1863, Sist., 124.

Dakota to Texas.

SOLENOGLYPHA—Continued.

- Var. PYRRHUS
Caulisoma pyrrhus Cope, 1866, Pr. Phil. Ac., 308, 310; Coues, 1875, Wheeler's Rep., V, 608. *Crotalus pyrrhus* Cope, 1875, Checklist, 33; Yarrow, 1875, Wheeler's Rep., V, 535. A doubtful variety.
 Arizona.

- CROTALUS OREGONUS
Crotalus oregonus Holbr., 1842, Herp., III, 21, pl. 3; DeK., 1842, N. Y. Fauna, Rept., 57; B. & G., 1853, Serp., 145; Bd., P. R. R. Rep., X, pl. X, 24, f. 6; D. & B., 1854, Erp., VII, 1482; Cope, 1859, Pr. Phil. Ac., 337. *Crotalus lucifer* B. & G., 1852, Pr. Phil. Ac., 177, — 1853, Serp., 6 (part); D. & B., 1854, Erp., VII, 1482; Grd., 1858, Herp., Wilkes' Exp., 187, pl. 15, f. 1-6; Bd., 1859, P. R. R. Rep., X, Williamson, 10, pl. 11, pl. 36, f. 1; Cooper, 1860, P. R. R. Rep., XII, pt. 2, 295. *Caulisoma lucifer* Cope, 1861, Mitch. Res., 121, — 1866, Pr. Phil. Ac., 307, 309.

Internasals and prefrontals small irregular. Labials 15 to 16, two to three rows of scales between them and the orbit. Scales in 25 (25 to 27) rows. Ventrals 168 to 177. Subcaudals 22 to 25. Light brown, tinged with yellow or ash, darker posteriorly. Irregular rhomboid dark-bordered areas of the ground color or darker form a vertebral series. Posteriorly the spots become transverse bands. Outside of the dark borders there are more or less indistinct lines of light color which cross on the flank inclosing a similar small spot on the lower part of the side opposite each of the dorsal series. Of spots and bands there are about forty to the series. A light line crosses the frontals, and passes through the supraciliary to the posterior labial. Another line of light color passes in front of the eye, and includes the lower portions of the labials. The dark colors are darker and the light whiter than those of *confluentus*.

Oregon to California.

- Var. LUCIFER 114
Crotalus lucifer B. & G., 1852, Pr. Phil. Ac., 177, and 1853, Serp., 6, part; Cope, 1859, Pr. Phil. Ac., 337, — 1875, Checklist, 33; Yarr., 1875, Wheeler's Rep., V, 533. *Caulisoma lucifer* Coues, 1875, Wheeler's Rep., V, 606. *Crotalus adamantus* var. *lucifer* Jan, 1863, Sist., 124. *Crotalus horridus* Gthr., 1858, Cat., 267, ? part.

California to Mexico.

- Var. CERBERUS
Caulisoma lucifer var. *cerberus* Coues, 1875, Wheeler's Rep., V, 607.
 "Nearly black, especially on the head."

Arizona.

- Var. MITCHELLI
Caulisoma mitchellii Cope, 1861, Pr. Phil. Ac., 293, 1866, Pr. Phil. Ac., 310. *Crotalus mitchellii* Yarr., 1875, Wheeler's Rep., V, 535; Cope, 1875, Checklist, 33.

Crown scales small, rough. One loreal. Anterior portion of nasal small, higher than long, separated from rostral and labials by small

SOLENOGLYPHA—Continued.

scales. Labials 16, the last large, three rows between them and the orbit. Scales in 25 rows, keels of outer row weaker. Ventrals 198. Subcaudals 26. Greyish-yellow, punctulate with brown. Dorsal spots about 42, lateral angles produced down the flank. Five black cross bands on the tail. A yellow band from the nasals to the upper borders of the hinder labials. Above this a brown band from the eye.

Lower California.

Var. ENYO

Crotalus enyo Cope, 1861, Pr. Phil. Ac., 293, — 1866, Pr. Phil. Ac., 309. *Crotalus enyo* Yarr., 1875, Wheeler's Rep., V, 534; Cope, 1875, Checklist, 33.

Internasals and frontals small, scale-like. Rostral low, equilateral. Nasal in contact with rostral. Numerous small scales in front of the pit. Labials 13 or 14, separated from orbit by three scales. Infralabials 14 or 15. Scales in 23 rows, keels in outer two or three obsolete. Ventrals 166. Subcaudals 26. Light greyish-brown, shaded with yellow. A pair of small brown spots on vertex. A light band bordered with dark across the supraciliaries. A band from the eye over the last labial to the neck. A series of about thirty-three vertebral spots, longer anteriorly, broader posteriorly, and fusing with the lateral spots, forming vertical bands on the flanks. Belly yellow, mottled with darker. Resembles *C. molossus* in coloration.

Lower California.

CROTALUS EXUL sp. n. 114

Cedros Island.

CROTALUS HORRIDUS "Banded Rattlesnake" 115

Crotalus horridus Linn., 1758, Syst., I, 214, — 1766, Syst., I, 372; Bonn., 1790, Ophiol., I, pl. 2, f. 3; Shaw, 1802, Zool., III, 317, pl. 83; Cuv., 1817, R. An., II, 78, — 1836, R. An., I, 413; Guerin, 1829-38, Icon. R. An., Rept., 15, pl. 23, f. 2; Griffith, 1831, An. King., IX, 267; Gray, 1831, Syn. Rept., 78, mixed; LeC., 1853, Pr. Phil. Ac., 417; Cope, 1859, Pr. Phil. Ac., 338, — 1875, Checklist, 33; Yarr., 1875, Wheeler's Rep., V, 534. *Crotaliona horrida* Cope, 1861, Mitch. Res., 122, — 1866, Pr. Phil. Ac., 309. *Crotaliona durissa* Laur., 1768, Syn. Rept., 93. *Crotalus durissus* Gmel., 1788, Syst. Linn., I, 1081, after Laur., mixed: Latr., 1801, Rept., III, 190; Daud., 1803, Rept., V, 304, pl. 68, f. 12; Harl., 1827, Jour. Phil. Ac., 368, — 1835, Med. Res., 132; Schlegel, 1837, Essai, I, 192, II, 365, pl. 20, f. 15, 16; Storer, 1839, Rept. Mass., 233; Holbr., 1842, Herp., III, 9, pl. 1; DeK., 1842, N. Y. Fauna, III, Rept., 55, pl. 9, f. 19; B. & G., 1853, Serp., 1; LeC., 1853, Med. Jour., 663; Bd., 1854, N. Y. Serp., 9, — 1859, P. R. R. Rep., X, Whipple, 39, pl. 24, f. 1; D. & B., 1854, Erp., VII, 1465; Mart., 1856, Mus. Berl., 34; Jan., 1859, Rev. Mag. Zool., Extr., p. 28, 1863, Sist., 123, — 1874, Icon., livr. 46, pl. 1, f. 1, 2; Allen, 1869, Pr. Bost. Soc., 179. *Crotalium cyanurus* Raf., 1818, Am. Month. Mag., III, 446, IV, 41. *Urocrotalon durissus* Fitz., 1843, Syst., 29. *Urop-*

SOLENOGLYPHA—Continued.

sophus durissus Gray, 1849, Cat., 19. *Crotalus atricaudatus* Latr., 1801, Rept., III 209; Daud., 1803, Rept., V, 316; Merr., 1820, Syst., 157, Boie, 1827, Isis, 562; Wagl., 1830, Amph., 177; Gray, 1842, Zool. Misc., 51. ? *C. durissus* var. *concolor* Jan, 1859, Rev. Mag. Zool., Extr., p. 28. *C. durissus* var. *melanurus* Jan, 1859, Rev. Mag., Extr., p. 28, — 1863, Sist., 123. *Le durissus* LaC., 1789, Serp., pl. 18, f. 3, not text. *Le Boiguira* LaC., 1789, pl. 18, f. 1, not text. *Vipera caudisona* Catesby, 1743, Carol., II, 41.

Texas to New England.

CROTALUS CERASTES 116

Crotalus cerastes Hall., 1854, Pr. Phil. Ac., 95, — 1859, P. R. R. Rep., X, Williamson, 17, pl. IV, f. 1, pl. 35, f. 4, 5; (Kenn. Bd., 1859, Mex. Bound., II, 14, pl. 3; Cope, 1829, Pr. Phil. Ac., 337, — 1875, Checklist, 33; Jan, 1863, Sist., 124, — 1874, Icon., livr. 46, pl. 3, f. 5; Yarr., 1875, Wheeler's Rep., V, 534. *Caudisona cerastes* Cope, 1861, Mitch. Res., 124, — 1866, Pr. Phil. Ac., 309. *Caudisona (Aechmophrys) cerastes* Coles, 1875, Wheeler's Rep., V, 609. *Aechmophrys cerastes* Coles, 1875, Wheeler's Rep., V, 609.

California; Arizona; Mexico.

? CROTALUS LEPIDUS 117

Caudisona lepida Kenn., 1861, Pr. Phil. Ac., 206; Cope, 1861, Mitch. Res., 124. *Aphouspis lepida* Cope, 1866, Pr. Phil. Ac., 310, — 1875, Checklist, 33; Yarr., 1875, Wheeler's Rep., V, 535.

Western Texas.

CROTALUS TIGRIS 117

Crotalus tigris (Kenn.) Bd., 1859, Mex. Bound., II, Rept., 14, pl. 4, — 1859, P. R. R. Rep., X, pl. 35, f. 1; Cope, 1859, Pr. Phil. Ac., 338, — 1875, Checklist, 33; Yarr., 1875, Wheeler's Rep., V, 534. *Caudisona tigris* Cope, 1861, Mitch. Res., 122, — 1866, Pr. Phil. Ac., 309; Coles, 1875, Wheeler's Rep., V, 608.

Mexico.

CROTALUS TRISERIATUS 112, 117

Crotalus triseriatus Wieg., 1828, Mus. Berl.; Mart., 1856, Mus. Berl., 34; Cope, 1864, Pr. Phil. Ac., 166; Yarr., 1875, Wheeler's Rep., V, 533. *Uropsophus triseriatus* Wagl., 1830, Amph., 176; Gray, 1842, Zool. Misc., 51. *Caudisona triseriata* Cope, 1866, Pr. Phil. Ac., 308. *Crotalus lugubris* Jan, 1859, Rev. Mag. Zool., Extr., pp. 28, 31, pl. E, f. 4, — 1863, Sist., 124; Cope, 1859, Pr. Phil. Ac., 338, — 1864, Pr. Phil. Ac., 166; Duges, 1879, La Natureza, IV, 25. *Caudisona lugubris* Cope, 1861, Mitch. Res., 122. *Crotalus lugubris* var. *multinaculata* Jan, 1863, Sist., 124, — 1874, Icon., livr. 46, pl. 3, f. 3. *Caudisona polysticta* Cope, 1865, Pr. Phil. Ac., 191, — 1866, Pr. Phil. Ac., 308. *Crotalus polystictus* Yarr., 1875, Wheeler's Rep., V, 533. *Crotalus (Crotalophorus) miliaris* var. *triseriatus* Jan, 1859, Rev. Mag. Zool., Extr., p. 29.

Mexico.

SOLENOGLYPHA—Continued.

Var. *JIMENEZII*

Crotalus jimenezii Duges, 1879, La Naturaleza, IV, 23.

Internasals four. Prefrontals four. Labials 14 to 15, separated from the orbit by two or three series. Scales in 25 rows, outer three smooth. Brownish-yellow, tinted with flesh color below, with five series of brown white-margined spots, of which the vertebral are larger. On each side of the occiput and neck a brown band. A brown band from the eye above the angle of the mouth to the neck; above this another from the supraocular. A vertical band from the eye to the labials; a similar one from the pit to the mouth. Mexico.

SISTRURIUS 110, 118

Caudisoma Fitz., 1826, Nene Class., 63 (not of Laurent); Wagl., 1830, Amph., 176; Bonap., 1832, Saggio, 24; Gray, 1842, Zool. Misc., 51; Fitz., 1843, Syst. Rept., 29; Cope, 1875, Checklist, 33. *Crotalophorus* Gray, 1825, Ann. Phil., 205 (not of Linné); Holbr., 1842, Herp., III, 25; Gray, 1849, Cat. Serp., 17; B. & G., 1853, Serp., 11; Cope, 1859, Pr. Phil. Ac., 336. *Crotalus* Cope, 1861, Mitch. Res., App., 119 (not of Linné).

SISTRURIUS CATENATUS 118

Crotalinus catenatus Raf., 1818, Am. Month. Mag., IV, 41. *Crotalus tergeminus* Say, 1823, Long's Exp., I, 499; Boie, 1824, Isis, 270 (Cope), — 1827, Isis, 563; Harl., 1827, Jour. Phil. Ac., 372, — 1835, Med. Res., 135; D. & B., 1854, Erp., VII, 1479, pl. 84 bis, f. 5; Cope, 1861, Mitch. Res., 125; Hayd., 1862, Trans. Am. Phil. Soc., 177. *Crotalophorus tergeminus* Gray, 1831, Syn. Rept., 78, — 1849, Cat. Serp., 18; Holbr., 1842, Herp., III, 29, pl. V; DeK., 1842, N. Y. Rept., 57; B. & G., 1853, Serp., 14; Bd., 1854, Serp. N. Y., 11, — 1859, P. R. R. Rep., X, 14, pl. 25, f. 9; Cope, 1859, Pr. Phil. Ac., 336. *Caudisoma tergemina* Wagl., 1830, Syst. Amph., 176; Cope, 1875, Checklist, 34; Coues and Yarrow, 1878, Hayd. Rep., IV, 269. *Crotalophorus kirtlandi* Holbr., 1842, Herp., III, 31, pl. 6; DeK., 1842, N. Y. Rept., 57; Gray, 1849, Cat. Serp., 18; B. & G., 1853, Serp., 16; D. & B., 1854, Erp., VII, 1482; Bd., 1854, Serp. N. Y., 12 (*C. missassauga* Kirtland), — 1859, P. R. R. Rep., X, 14, pl. 25, f. 11; Cope, 1859, Pr. Phil. Ac., 336. *Crotalophorus* sp. Agassiz, 1850, Lake Superior, 381, pl. 6, f. 6-8. Ohio and Michigan to the Plains and southward to Mississippi.

Var. *CONSORIS*

Crotalophorus consors B. & G., 1853, Serp., 12; D. & B., 1854, Erp., VII, 1482; Bd., 1859, P. R. R. Rep., pl. 24, f. 8.

Frontal cordiform. Prefrontals moderate. Parietals rather broad. Scales in 25 rows, outer two smooth. Ventrals 147; subcaudals 25. No vertebral red line. Olivaceous brown, with seven series of about fifty spots each, all small. Blotches darker, with a narrow border of darker, surrounded by a light margin. Two lines from the vertex along the neck to the first blotch. A band from the eye to the neck. A whitish line from the nostril to the angle of the mouth. A vertical band on each side of the pit. Belly yellowish-white, blotched with darker. Texas.

SOLENOGLYPHA—Continued.

SISTER RUS MILLARIUS 119

Crotalus millarius Linn., 1766, Syst., I, 372; Gmel., 1788, Syst. Linn., I, 1080; Bonn., 1790, Ophiol., 1; Latr., 1801, Rept., II, 205; Shaw, 1802, Zool., III, 336; Daud., 1803, Rept., V, 328; Cuv., 1817, R. An., II, 79; Merr., 1820, Syst., 156; Boie, 1827, Isis, 562; Harl., 1827, Jour. Phil. Ac., 370. — 1835, Med. Res., 134; Schleg., 1837, Essai, I, 192, II, 569, pl. 20, f. 17, 18; Holbr., 1838, Herp., II, pl. 15; D. & B., 1854, Erp., VII, 1477; Cope, 1861, Mitch. Res., 124; Jan., 1859, Rev. Mag. Zool., Extr., p. 28, — 1863, Sist., 124. *Crotalophorus millarius* Gray, 1825, Ann. Phil., 205, — 1831, Syn. Rept., 78; Holbr., 1842, Herp., III, 25, pl. 4; DeK., 1842, N. Y. Rept., 57; Gray, 1849, Cat. Snakes, 17; B. & G., 1853, Serp., 11; Gthr., 1858, Cat., 267; Bd., 1859, P. R. R. Rep., X, Whipple, 40, pl. 24, f. 7; Cope, 1859, Pr. Phil. Ac., 336. *Caulisoma millaria* Fitz., 1826, Neue Class., 63, — 1843, Syst. Rept., 29; Wagl., 1830, Amph., 176; Gray, 1842, Zool. Misc., 51; Cope, 1875, Checklist, 34, — Pr. Am. Phil. Soc., 64, — 1880, Bull. 20, U. S. Mus., 24. *Crotalus millarius* var. *tergenius* Jan., 1859, Rev. Mag. Zool., Extr., 28, — 1863, Sist., 124, — 1874 Icon., livr. 46, pl. 3, f. 4. ? *Crotalus tergenius* Mart., 1856, Mus. Berl., 34. *Vipera caudisoma minor* Catesby, 1743, Carol., II, 42. *Le Millet* LaC., 1789, Quad. Ovip. Serp., II, 421, pl. 18, f. 2.

Southern States.

Var. EDWARDSII

Crotalophorus edwardsii B. & G., 1853, Serp., 15; D. & B., 1854, Erp., VII, 1485; Bd., 1858, Mex. Bound., II, Rept., 15, — 1859, P. R. R. Rep., X, pl. 25, f. 10; Duges, 1879, La Natureza, IV, 27. *Crotalus edwardsii* Cope, 1861, Mitch. Res., 125; Coues, 1875, Wheeler's Rep., V, 610. *Caulisoma edwardsii* Cope, 1875, Checklist, 34; Yarrow, 1875, Wheeler's Rep., V, 531. ? *Crotalus millarius* var. *edwardsii* Jan., 1863, Sist., 124. *Crotalus millarius* Jan., 1874, Icon., livr. 46, pl. 3, f. 6.

Scales in 23 rows, outer two smooth. Ventrals 143 to 153. Subcaudals 23. Yellowish-brown, with seven (five to seven) series of spots. A band of brown from prefrontal over eye to neck; beneath this a yellowish stripe from nostril to neck. Yellowish bars downward from each side of the pit. About forty-two black-bordered light-margined blotches in the vertebral series. Belly light yellowish, mottled with brown.

Texas; Arizona; Sonora; Mexico.

Var. RAVUS 120

Crotalus rarus Cope, 1865, Pr. Phil. Ac., 191. *Caulisoma rava* Cope, 1875, Checklist, 33.

Mexican Plateau.

ANCISTRODON 120

Aghistredon Pal. de Beauv., 1799, Trans. Am. Phil. Soc., IV, 381; B. & G., 1853, Serp., 17. *Cenchris* Daud., 1803, Rept., VIII, 388. *Tisiphone* Fitz., 1826, Neue Class., 63. *Taricophis* Troost., 1833, Ann. Lyc. N. Y., III, 176.

SOLENOGLYPHA—Continued.

Ancistrodon contourrix (Copperhead) 120

B. & G., 1853, Serp., 17; Bl., 1854, Serp. N. Y., 13, — 1859, P. R. R. Rep., X, pl. 25, f. 12; Cope, 1859, Pr. Phil. Ac., 336, — 1875, Checklist, 34, — 1880, Bull. 20, U. S. Mus., 24; Allen, 1869, Pr. Bost. Soc., 11. *Coluber contourrix* Linn., 1758, Syst., I, 216. *Boa contourrix* Linn., 1766, Syst., I, 373; Gmel., 1788, Syst. Linn., I, 1083. *Trigonocephalus contourrix* Holbr., 1842, Herp., III, 39, pl. 8; D. & B., 1854, Erp., VII, 1494; Mart., 1856, Mus. Berl., 35; Hall., 1856, Pr. Phil. Ac., 249; Jan., 1859, Rev. Mag. Zoöl., Extr., p. 29, — 1863, Sist., 125, — 1874, Icon., livr. 46, pl. 5, f. 1. *Trigonocephalus hibernicus* Dum., 1852, Mem. Acad. Sci., XXIII, 534. *T. cenchris* Schleg., 1837, Essai, I, 191, II, 553, pl. 20, f. 10, 11. *Cenchris contourrix* Gray, 1825, Ann. Phil., — 1849, Cat. 16. *C. mockson* Daud., 1803, Rept., V, 358, pl. 60, f. 25, pl. 70, f. 3, 4; Harl., 1827, Jour. Phil. Ac., 266, — 1835, Med. Res., 128; Wagl., 1830, Syst. Amph., 175. *Aglistrodon mokason* Beauv., 1799, Trans. Am. Phil. Soc., IV, 380. *Scytalus cupreus* Raf., 1834, Am. Jour. Sci., I, 85. Mississippi Valley to the Atlantic.

Var. *ATROFUSCUS* ("Highland Moccasin")

Acontias atrofuscus Troost, 1836, Ann. N. Y. Lyc., 181. *Toxicophis atrofuscus* Troost, 1836, Ann. N. Y. Lyc., 190; B. & G., 1853, Serp., 150. *Cenchris atrofuscus* Gray, 1849, Cat., 16. *Trigonocephalus atrofuscus* Holbr., 1842, Herp., III, 43, pl. 9; D. & B., 1854, Erp., VII, 1495. *Ancistrodon atrofuscus* Cope, 1875, Checklist 34.

"The colour of the upper parts of the body must be considered as black, variegated with brown spots, which are not of a uniform shade, but pass from dark into a lighter colour; they are irregularly scattered over the body; nevertheless, they exhibit a tendency to run from one side toward the other, widening towards the back. The light spots are composed of smoky gray, passing into blackish brown, exhibiting all the intermediate shades between these two colours; they are rather small stripes of the breadth of from two to four scales, and disappear near the tail altogether, which is entirely black, having only four small white points, which are probably accidental." Top of head black, whitish behind the eye. A band from the eye above posterior labial. Belly whitish, blotched with black.

Mountain region from Virginia southward.

Ancistrodon piscivorus (Moccasin) 121

Cadalus piscivorus Latc., 1789, Quad. Ovip. Serp., II, pp. 130, 424. *Trigonocephalus piscivorus* Holbr., 1838, Herp., II, 63, pl. 13, — 1842, Herp., III, 33, pl. 7; D. & B., 1854, Erp., VII, 1491; Mart., 1856, Mus. Berl., 35; Jan., 1859, Rev. Mag. Zoöl., Extr., p. 29, — 1863, Sist., 125, — 1874, Icon., livr. 46, pl. 4, f. 2-4. *Cenchris piscivorus* Gray, 1849, Cat. Snakes, 16; Gthr., 1858, Cat., 267. *Toxicophis piscivorus* B. & G., 1853, Serp., 19; Bl., 1859, P. R. R. Rep., X, Whipple's, 40, pl. 25, f. 13. *Ancistrodon piscivorus* Cope, 1859, Pr. Phil. Ac., 336, — 1877, Pr. Am. Phil. Soc., 64, — 1880, Bull. 20, U. S. Mus., 24. *Ancistrodon piscivorus* subsp. *piscivorus* Cope, 1875, Checklist, 34. *Acontias leucostomus* Troost, 1833, Ann. N. Y. Lyc., III, 176. South Carolina to Texas.

SOLENOGLYPHA—Continued.

Var. PUGNAX

Toxicophis pugnax B. & G., 1853, Serp., pp. 20, 156; Bd., 1859, P. R. R. Rep., X, pl. 25, f. 14. *Toxicophis piscivorus* subsp. *pugnax* Cope, 1875, Checklist, 31. *Trigonocephalus piscivorus* var. *pugnax* Jan, 1863, Sist., 125.

No loreal. Second labial with a narrow edge or acute angle at the mouth. Third and fourth labials in orbit. Light olive-brown above; yellowish, with brownish blotches on each side, beneath. Tail black. About thirty blotches of darker on the back. Margins irregular, often indistinct. Check band obsolete. Ventrals 145. Subcaudals 22, plus 23 pairs. Scales in 25 rows.

Texas.

ANCISTRODON BILINEATUS

Guenther, 1863, Ann. Mag. Nat. Hist., 364; Cope, 1865, Pr. Phil. Ac., 191; Sumichrast, 1881, Bull. Soc. Zool. de France, 185.

"Shining deep black, with scattered white spots, arranged in narrow, distant, transverse bands; the white spots are more numerous and irregular on the belly; a yellow line runs from the rostral along the canthus rostralis and the supraciliary edge to behind the angle of the mouth. A yellow band along the upper labials, the lower margin of which is black. Rostral, with a vertical yellow band, continued on the chin. Upper labials eight. Scales keeled, in 23 series. Ventrals 137; subcaudals 65, the fourteen last double."

Western Mexico; Isthmus of Tehuantepec.

LACHESIS 122

Daudin, 1803, Rept., V, 351; Wagl., 1830, Syst., 175; D. & B., 1854, Erp., VII, 1483; Gray, 1849, Cat., 13. *Craspedocephalus* Gray, 1825, Ann. Phil., 205.

LACHESIS MUTUS 122

Crotalus mutus Linn., 1766, Syst., I, 373; Gmel., 1788, Syst. Linn., I, 1082; Schlegel, 1837, Essai, II, 570, pl. 20, f. 19, 20; Jan, 1859, Rev. Mag. Zool., Extr., p. 29. *Lachesis mutus* Daud., 1803, Rept., V, 351; Wagl., 1830, Syst., 175; Gray, 1842, Zool. Misc., 50, — 1849, Cat., 13; D. & B., 1854, Erp., VII, 1485; Mart., 1856, Mus. Berl., 35; Cope, 1859, Pr. Phil. Ac., 339; Jan, 1863, Sist., 124. *Lachesis rhomboides* Max., 1824, Abbild. Bras. Bief. 5, pl. 5; Fitz., 1826, Neue Class., 63, — 1843, Syst., 29; Gray, 1831, Syn. Rept., 79. *Craspedocephalus crocotalinus* Gray, 1825, Ann. Phil., 205. *Cophis crocotalius* Merr., 1820, Syst., 154. *Bothrops surucucu* Wagl., 1824, Spix. Serp. Bras., p. 59, pl. 23? *Boa muta* LaC., 1789, Quad. Ovip. Serp., II, 389. *Boa crocotalina* Shaw, 1802, Zool., III, 352. *Segtate catenatus* Latr., 1801, Rept., III, 162. *Trigonocephalus rhombifer* Cuv., 1829, R. An. II, 90.

Brazil to Mexico.

Var. STENOPHRYS 123

Lachesis stenophrys Cope, 1875, Jour. Phil. Ac., 152.

Central America.

SOLENOGLYPHA—Continued.

- TRIGONOCEPHALUS 123
- Trigonocephalus* Oppel, 1811, Class. Rept., 55. *Cophias*, Merrem, 1820, Syst., 154; Gray, 1831, Syn. Rept., 79. *Bothrops* Wagler, 1824, Spix. Serp. Bras., 50, — 1830, Syst. Amph., 174; D. & B., 1854, Erp., VII, 1502; Fitz., 1843, Syst. Rept., 29. *Craspedocephalus* Fitz., 1826, Class. Rept., 62; Gray, 1849, Cat., 4.
- TRIGONOCEPHALUS ASPER 124
- Possibly a variety of *T. laucobatus*, which it resembles, but which is a West Indian species.
- Isthmus.
- TRIGONOCEPHALUS ATROR 124
- Culther atror* Linn., 1754, Mus. Ad. Fridr., pl. 22, f. 2; Gmel., 1788, Syst. Linn., I, 1107. *Cophias atror* Merr., 1820, Syst., 154. *Viperæ atror* Laur., 1768, Syst., 103. *Bothrops atroræ* Wagl., 1830, Syst. Amph., 174; D. & B., Erp., VII, 1507; Mart., 1856, Mus. Berl., 35. *Bothrops tessellatus* pl. 21, f. 2, *B. tucnatus* pl. 21, f. 3, and *B. leucurus* pl. 22, f. 2, Wagl. in Spix. Serp. Bras. *B. atror* var. *tessellatus* Jan, 1875, Icon., livr. 47, pl. 2, f. 4. *Craspedocephalus atror* Fitz., 1826, Neue Class., 66; Gray, 1849, Cat., 6. *Trigonocephalus atror* Kuhl., 1820, Beitr., pp. 91, 100; Schleg., 1837, Essai, II, pl. 19, f. 5, 6.
- Brazil to Mexico, through varieties.
- TRIGONOCEPHALUS AFFINIS 125
- Bothrops affinis* Bocourt, 1863, Ann. Sci. Nat., pt. II, 201. *Bothriopsis affinis* Cope, 1875, Jour. Phil. Ac., 150.
- Guatemala.
- TELECRASPIS 126
- Cope, 1869, Pr. Phil. Ac., 345.
- TELECRASPIS UNDULATUS 126
- Trigonocephalus undulatus* Jan, 1859, Rev. Mag. Zool., Extr., p. 30. *T. (Atropos) undulatus* Jan, 1859, l. c. p. 32. *Atropos undulatus* Jan, 1859, l. c., pl. E, f. 6, — 1863, Sist., 127.
- Mexico.
- TELECRASPIS NIGROADSPERSUS 126
- Bothrops (Telecraspis) nigroadspersus* Steind., 1870, Sitzb. Ak. Wien, 348, pl. 8.
- Central America to Mexico.
- TELECRASPIS NUMMIFER 127
- Atropos nummifer* Ruepp., 1845, Mus. Senck., 313. *Telecraspis nummifer* Gthr., 1863, Ann. Mag. Nat. Hist., 25. *Atropos mexicanus* D. & B., 1854, Erp., VII, 1521, pl. 83 bis, f. 1, 2. *Bothriochis mexicanus* Cope, 1861, Pr. Phil. Ac., 294. ? *Trigonocephalus nummifer* Jan, 1859, Rev. Mag. Zool., Extr., 30. ? *Bothrops nummifer* Jan, 1863, Sist., 126, — 1875, Icon., livr. 47, pl. 5, f. 2.
- Mexico.

INDEX.

A.			
ahacurus	36, 144	atrofuscus	178
Ablabes, Contia	164	Atropos	126, 180
Ablabes, Ophibolus	155	atrox, Crotalus	114, 172
Abothropha	xxii, 104	atrox, Trigoniceps	124, 180
Acacophitia	xx, 1, 11, 133	aurata	136
Acontiade	xv	auribundus	50, 150
Acontias	xv	aurigulus	44, 148
Acontias, Ancistrodon	178		
acuminatus	136	B.	
adamanteus	112, 171	baipilii, Elaphis	152
Additions	128	bairelli, Salvadoria	145
Adelophis	143	balteata	157
adnexus	144	baronis-muelleri	142
Aechmophrys	175	Basanium	40, 146
aemula	92, 164	Basiliscus	xiv
aeneus	21, 136	basiliscus, Crotalus	171
aestivus	40, 146	basimaculatus	3, 129
affinis, Elaps	106, 168	Bastard Turtle	vii
affinis, Pityophis	151	Batrachia	xxvii
affinis, Trigoniceps	125, 180	Batrachians	iii
Agamidae	xii	Bellophs	156
agassizii, Helicops	144	bellona	53, 151
agassizii, Nerodia	141	Bergia	162
Agkistrodon	177	bertholdi	79, 161
alleganiensis	54, 152	bicolor, Geophis	102, 167
allegheni	31, 144	bicolor, Pelamys	109, 170
Alytes	xxx	bifasciatus	165
ambilis	159	bifasciatus, Ancistrodon	179
amarus	156	bifasciatus, Coluber	149
Amblymetopon	162	bimaculata	87, 163
Amblyrhynchus	xiii	bipunctata, Coronella	63, 154
amoena	100, 167	bipunctata, Tropidonotus	138
Amphiuma	xxxix	bipunctiger	168
Amphisbaena	xv	biscutatum	16, 131
Amyda	vi	Black Snake	xix, 41, 146
Anasime	151	Black Snake, Pilot	151
Ancistrodon	xxvi, 110, 120, 177	Blue Tail	xv
Anguis	xiv	Boa	8, 132
angustirostris	159	Boacidae	7, 8, 132
annectens	159	Boiquira	171
Anniella	xv	Bohrichis	180
Anniellidae	xv	Bothriopsis	180
annulata, Contia	91, 164	Bothropha	xxiii, 110, 170
annulatum, Sibon	17, 135	Bothrops	179, 180
annulatus, Diadophis	70, 158	bottae	7, 131
annulatus, Heterodon	160	boylii	69, 157
annulatus, Ophibolus	156	Brachyorrhos	167
Anoliidae	xiv	breyvis	134
Anomalepis	2, 4, 129	Bufo	xxx
anthracinus	xv	Bull Frog	xxx
Anura	xxviii-xxx	Bull Snake	150
apiatus	106, 169		
Apoaspis	175	C.	
Apoda	xxviii	cacodaemon	160
Arizona	151	Caecilia	xxviii
army	72, 158	Calamaria	165
asper	124, 180	calamarina	87, 163
Aspidonectes	vi	Calamarinae	xxii, 11, 81, 161
atmodes	160	Calamys	v
atrata	140	calliformae	157
atricaudatus	175	calligaster, Elaphis	152
		calligaster, Ophibolus	66, 155
		Calopisma	143
		cana	83, 161
		canina	149
		Carphophis	166
		Carphophis	99, 166
		carolinianus	152
		caseavella	171
		catenatus, Scytale	179
		catenatus	118, 176
		catenifer	52, 150
		cafe-byi	160
		Catostoma	167
		Caudisoma, Sistrurus	176
		Caudisoma, Crotalus	170
		caelano	143
		Celuta	101, 166
		Cemophora	77, 160
		ceucha	14, 134
		Cenebris	177, 178
		ceneoalt	134
		cerastes	116, 175
		Ceratophora	xiii
		Ceratophrys	xxx
		cerberus	173
		cerebripunctatus	107, 169
		Chain Snake	68, 156
		Chamaeleonidae	xi
		Changes of color	x
		Charina	7, 131
		Cheliorhina	85, 162
		Chelonia	vii
		Chelydra	vi
		Chersodromus	95, 165
		Chicken Snake	153
		Chilabothrus	xvii, 132
		Chilomeniscus	166
		Chilopoma	136
		Chionactis	164
		Chirotes	xvi
		Chlamydosaurus	xiii, xiv
		chlorophaca	161
		Chlorosoma	146
		Churehillia	151
		cincta	100, 166
		Cinosternum	vi
		Cistudo	vi
		clarkii	142
		clericius	155
		cloelia	19, 136
		Cnemidophorus	xiv
		Coach-whip Snake	147
		coccinea, Cemophora	78, 160
		coccinea, Coronella	154, 155
		cognatus	159
		Coleonyx	xii
		collaris, Nimia	95, 165
		collaris, Tropidonotus	25, 139
		Colpochelys	vii
		Coluber	40, 146
		Colubridae	xxii, 11, 133
		Colubrinae	xxii, 11, 37, 145
		colubrinus, Xenodon	79, 161

- compressicaudus 27, 142
 compressus 152
 comp-solemis 142
 concinus 140
 concolor 175
 confinis 152
 confluentus 114, 172
 Coniophanes 154
 conjuncta 157
 Conocephalus 166
 Conocera xxiii, 104, 168
 Conopsis xiii
 consors 83, 161
 constrictor, Boa 9, 132
 constrictor, Coluber xix, 41, 146
 Conia 90, 163
 contortrix 120, 178
 cooperi 140
 copei, Conophora 78, 161
 copei, Storeia 31, 143
 Cophias 180
 Copperhead 120, 178
 corais 48, 149
 corallinus 107
 coronata, Tantilla 88, 163
 coronatus, Spilotes 119
 Coronellinae xxii, 11, 59, 154
 Corn Snake 55, 152
 Corrections 128
 Corythodon 146
 couchii, Eutaenia 139
 couchii, Nerodia 141
 cooperi 48, 149
 Craspedocephalus 179, 180
 Crotalidae xxiii, 110, 170
 Crotalini 170
 crotalinus, Craspedocephalus 179
 Crotalinus, Crotalus 170
 Crotaloidea 170
 Crotalophorus, Crotalus 170
 Crotalophorus, Sistrurus 118, 176
 Crotalus xxvi, 110, 170
 cyanurus 174
 Cyclodus xv
 cyclophus 39, 146
 cyclopium 26, 142
 cyrtopsis 158, 140
 Cystigathiidae xxix
- D.**
- Dactylethra xxx
 daudini 136
 decoratus, Diadophis 71, 158
 decoratus, Elaps 108, 169
 decurtata 39, 145
 de Filippii 161
 degenhardtii 162
 dekayi 31, 143
 Dendrophinae xxi, 11, 20, 135
 deppi, Elapochrus 157
 deppi, Pityophis 154
 diademata 96, 165
 Diadophis 70, 158
 diastema 107, 169
 dimidiatus, Leptognathus 14, 134
 dimidiatus, Ninia 165
 Diploclactylus xii
 Dipsadinae xxi, 11, 133
 Dujesi 12, 14, 14
 discolor 18, 135
 distans 105, 168
 divaricatus 168
 divinioluquax 132
 docilis 72, 158
 doliatus, Ophibolus 64, 154
 doliatus, triangulus var. 66, 155
 doliatus, Oxyrhopus 19, 136
 dorsalis 138
 Draco xiii
 Dromicus 56, 153
 drummondii 144
 Dryinus 136
 Drymobius 147
 duice 5, 139
 dulcis 130
 dumerillii 13, 133
 duris-sus 111, 170
 duris-sus 174
 dysopes 158
- E.**
- edwardsii 177
 Elaphis 53, 151
 Elapidae 104, 168
 Elapochrus 157
 Elapoides 165
 elapoides, Liophis 69, 157
 Elapomorplus 89, 163
 Elaps 104, 168
 elapsoides 65, 155
 elegans, Elaps 108, 169
 elegans, Pityophis 151
 elegans, Tropicodonotus 110
 elegans, Virginia 98, 166
 emoryi 152
 emunctus 3, 129
 Enys xv
 Enicognathus 158
 enyo 174
 epihippius 166
 episcopa 94, 164
 epistema 106, 169
 erbenus 149
 Eretmochelys v, vii
 Erycidae 7, 131
 Erycina 131
 erythrogaster 26, 141
 erythrogrammus 55, 144
 Eudromus 63, 151
 Eutaenia 157
 Evansii 155
 Evesia xv
 Eunectes xv
 Eumetes xx
 euryxanthus 107, 169
 eximius 155
 exsul 114, 174
- F.**
- Farancia 144
 fauxi 137
 Faus xxiii
 fasciata, Farancia 144
 fasciata, Tropicodipsas 133
 fasciatus, Eunectes xv
 fasciatus, Leptognathus 12, 133
 fasciatus, Tropicodonotus 141
 Fascination xxiv
 Ficinia 81, 161
 fissidens 62, 154
 fizingeri 168
 flagelliformis 42, 147
 flagellum 147
 flavigularis 148
 flavilabris 137
 flavilatus 58, 153
 flaviventris 42, 147
 floridanus, Lepidosternum xvi
 floridanus, Coluber 152
 foxii 147
 freminvillei 85, 162
 fremontii 147
 Freshwater Turtles vi
 frontalis 82, 161
 fulvius 105, 168
 fulviventris 71, 158
 funiceps 163
- G.**
- Garter Snake 24, 138
 Gecco xii
 Geconidae xii
 gentilis 66, 155
 Geophis 101, 167
 Georgia 47, 149
 Gerrhonotus xv
 getulus 68, 156
 glyphyros 140
 Glass Snake xv
 gracilis 87, 162
 grahamii, Regina 28, 142
 grahamii, Salvadora 38, 145
 Green Snake 40, 146
 Ground Snakes xxi
 Ground Snake 167
 gnotheri 63, 154
 gnathaus, Elaphis 55, 152
 Gyalopion 83, 161
 Gypochelys vi
- H.**
- haematois 150
 Haldea 166
 halwellii 163
 hammondi 140
 Harquin Snake 105, 168
 harperi 99, 166
 Hatteria x
 haydeni 138
 heermanni 150
 helena 100, 166
 Helicops 33, 143
 Heloderma xi
 Helodermidae xi
 Herpetodryas 146
 Herpeton xxi
 Heterodon xxi, 74, 159
 hexalepis 39, 145
 Hog-nose Snake 75, 159
 holbrookii, Elaphis 152
 holbrookii, Nerodia 141
 Homalacramion 162, 163
 Homalopsis 144
 Homalosoma 164
 Horned Toads xiv
 horridus 115, 174
 Hibernation xxviii
 hickamella 153
 Hierophis 147
 Highland Moccasin 178
 Himantodes 14, 134
 histicus 178
 humile 130
 Hydrophidae 109, 169
 Hydrophis xxx
 Hydrops 34, 144
 Hydrus 170
 Hyllidae xxx
 Hyllodes xxx

- Hypsiglena 79, 161
 hypostema 168
- I.**
- Icthyodi xxviii
 Idiophlops 3
 Iguana xiii
 Iguanidae xiii
 imperator 8, 132
 imperialis 61, 154
 infernalis 25, 139
 inornata, Virginia 97, 166
 inornatus, Chilabothrus, xvii, xx, 132
 intermedius, Crotalus 117
 isabella 131
 Icthmognathus 143
 isozona 92, 164
 isthmica 9, 132
- J.**
- jani 151
 jinuensis 176
- K.**
- kemnerlyi 77, 169
 kennebottii 138
 King Snake 68, 156
 kirtlandi, Regina 28, 147
 kirtlandi, Sistrurus 176
- L.**
- Lacertidae xiv
 Lachesis 122, 179
 laetus, Diadophis 158
 laetus, Scotophis 152
 Lamppropeltis 155
 Lamprosoma 164
 lanceolatus 180
 Land Turtles v
 laterale, Oligosoma xv
 lateralis, Coluber 148
 latericia 61, 154
 latifasciatus 107, 169
 latifasciatus 134
 latifrontalis 103, 167
 laurcutus 57, 153
 Leatherbacks vi
 leberis 27, 142
 lecontei, Crotalus 172
 lecontei, Rhinocheilus 72
 Le durissus 171
 Le Millet 176
 Lepidosternon xvi
 lepiscus 117, 175
 leptocephalus 139
 Leptognathus 12, 133
 leptogrammus xv
 Leptophis, Cyclophis 116
 Leptophis 20, 136
 leucomelas 15, 144
 leucostomus 178
 Liekeana 131
 hobmani 95, 165
 lindheimeri 54, 152
 lineata, Ficinia 161
 lineata, Storeria 32, 143
 lineata, Tachymenis 60, 154
 lineatocollis 151
 lineatus 136
 liophis 69, 157
 lippinsi 103, 167
 Lodia 165
- loetlingii 171
 Loggerheads vi
 longissimum 2, 129
 lucifer 114, 173
 lugubris 175
 Lyriocephalus xiii
 lyrophanes 134
- M.**
- maculata, Ficinia 84, 162
 maculatus, Coluber 152
 macrostemma 137
 majalis 146
 malacolemmys vi
 margravii 169
 mariana 25, 138
 margariferus 58, 153
 Marine Toad xxvi
 Massasauga 118, 176
 Masticeophis 41, 147
 McClellani 151
 medusa 142
 Megalobatrachus xxix
 Megalophrys xxx
 megalops 137
 melanocephala, Tachymenis, 63, 154
 melanocephalum, Homalocera-
 nion 163
 melanoleucus 51, 150
 melanurus, Crotalus 175
 melanurus, Spilotes 50, 150
 Menopoma xxviii
 mentovarius 42, 147
 mesomelanus 143
 mesotes 59
 Metapoceros xii
 mexicana, Boa 10, 132
 mexicana, Bergenia 162
 mexicana, Calopisma 145
 mexicanus, Anomalepis 4, 129
 mexicanus, Atropis 180
 mexicanus, Coluber 45, 148
 mexicanus, Elapomorphus, 89, 163
 mexicanus, Leptophis 20, 136
 mexicanus, Ophibolus 66, 155
 mexicanus, Pityophis 52, 151
 mexicanus, Spilotes 149
 Microphis 162
 microps 32, 143
 millirius 119, 177
 mitchellii 173
 mitis 93, 164
 Moccasin 121, 178
 Mookason 178
 Mohoch xii
 molossus, Coluber 152
 molossus, Crotalus 113, 171
 moroni 138
 myopium 5, 6, 131
 myriolepis 131
 Mud Turtles vi
 mullenbergii v
 multimaclata 175
 multistriata 155
 mutus 122, 179
- N.**
- nasicus 77, 160
 natus 83, 161
 Natriumae xxi, 11, 21, 136
 nebulatus 15, 135
 Necturus xxviii
 Nerodia xxi, 140
- Nessia xv
 niger, Heterodon 76, 160
 niger, Nerodia 141
 nigriceps 88, 163
 nigrocolpersens 126, 180
 nigrocinetus 166, 168
 nigroluteus 114
 Nimia 94, 165
 Nototrema xxx
 numifer 127, 180
- O.**
- obliquus 141
 obscura 138
 obsoletus, Elaphis 54, 151
 obsoletus, Spilotes 48, 149
 occipitalis, Conia 91, 164
 occipitalis, Diadophis 158
 occipitomaclata 30, 143
 ocellata 178
 ochrorhyncha 80, 161
 Oligosoma xv
 oliyacea 81, 162
 Onychophidia xx, 1, 7, 131
 Ophedrys 146
 Ophibolus 64, 154
 Ophidia iv, xvi, 1
 Ophiodes xiv
 Ophis 157
 Ophisaurus xv
 Ophthalmidion 129
 Ophthoglossa xxx, xxai
 ordinatus 25, 138
 ordinoides 146
 oregonus 173
 ornata, Pelamys 176
 ornatus, Pelamys 168
 ornatus, Coluber 148
 ornatus, Crotalus 171
 Osceola 155
 Oxybelis 21, 136
 Oxyrhina 162
 Oxyrhopus 18, 136
 Ozotheca vi
- P.**
- pacificum 135
 pallidus 158
 pantherinus 153
 parietalis 25, 139
 palustris vi
 platura 179
 Pelamys xviii, 169, 169
 perditus 3, 129
 personatum 135
 Petalognathus 13
 pictus 148
 pickeringi 24, 139
 pinus vi
 Pine Snake 150
 Pipa xxix, xxx
 piscivorus 121, 178
 Pityophis 51, 150
 Podocnemys vi
 poecilonotus 59, 150
 poizonias 111
 Poison Glands xxiv
 Polychrus xiv
 polycteta 175
 polyzona 156
 porcutus 141
 phenax 140
 phenops 5, 6, 151
 philippii 133


- pholidiatus 151
 Phrynosoma XIV
 Phylloclactus XI
 Phyllophilophis 39, 146
 Phymothyrus 145
 plainiceps 89, 163
 Platycerca XXIII, 109, 169
 Platypeltis VI
 platyrhinus 75, 179
 Pleurodeles XXIX
 plicatilis 144
 Pliocercus 157
 plimbea 131
 platinus 149
 plethura 92, 164
 proteoglypha XXII, 104, 168
 proterops 62, 154
 Proteus XXVIII, XXIX
 proximus 24, 137
 psammophis 147
 Pseudelaps 155
 Pseudocoryx 131
 pseudocrotalus 157
 Pseudopis XV
 Psyllozoon XII
 puzosax 179
 pulchellus 159
 pullatus 149
 punctatus 72, 158
 purpureocauda 164
 putnami 59, 153
 pygmaea 93, 161
 pyromelanus 67
 pyrhoneclax 67, 157
 pyrillus 114, 173
- Q.**
- quadriserialis 140
 quadrivittatus 56, 153
 quinqueimaculata 86, 162
 quinquevittatus 36, 145
- R.**
- Rhabdosoma 167
 Rachiolon XXI
 radix 138
 Ramiidae XXIX
 Rattles XXV
 ravus 120, 177
 regalis 73, 159
 Regina 27, 142
 reinwardti 114
 Remedies XXV
 Rena 5, 139
 rhomboccephalus 79
 Rhabdosoma 167
 Rheoporphus XXIX
 Rhodinea 158
 Rhinocheilus 73, 159
 rhinomegas 152
 Rhinostoma, Cemophora 169
 Rhinostoma, Contia 164
 Rhizolenta VIII
 rhombatus 179
 rhombifer, Crotalus 172
 rhombifer, Nepodia 26, 141
 rhombifer, Trigonoccephalus 179
 rhombomaculatus 156
 Rhynchocephalia X
 Ribbon Snake 23, 137
 rigidus 28, 142
 rose-fusca 131
 ruber 130
 rubriceps 152
- ruficeps 140
 rufipunctatus 23, 136
- S.**
- sackenii 137
 Salamandra XXIX
 Salvadoria XXII, 38, 145
 salmii 159
 Sauria IX
 saurita 23, 137
 saxi, Ophibolus 68, 156
 saxi, Pityophis 52, 150
 Sphaeriodactylus XII
 Sphaerops XIV
 Sphargis IV-VI
 Sphenodon X
 Sphenophis 154
 Spilotes 158
 spiloides 151
 Spilotes 47, 149
 spinalis 45, 148
 splendidus 157
 Spotted Ribbon Snake 138
 Sibon 12, 15, 134
 sieboldii 96, 165
 Simotes 160
 simus, Crotalus 171
 simus, Heterodon 76, 160
 speldon 25, 140
 Siren XXVIII
 sirtalis 24, 138
 Sistrurus XXVII, 110, 118, 176
 Shedding the Skin XXVIII
 Sea Snakes XXIII, 109, 169
 Sea Turtles VI
 sebae 95, 165
 semiamulata 90, 164
 semidoliatus 102, 167
 septentrionale 17, 135
 septemvittata, Regina 142
 septemvittatus, Hydrops 145
 Serpent locomotion XVII
 Serpents swallowing young, XVIII
 Solenoxypha XXII, 104, 110, 170
 Sonora 90, 164
 sonoriensis 172
 Sloughing XXVIII
 stenophrys 123, 179
 Stenorhina 85, 162
 Stenostoma 2, 4, 130
 Stenostominae 129
 stictogenys 158
 Storeria 29, 143
 storerioides 29, 143
 straminea 99, 166
 striatula 97, 166
 Streptophorus 95, 165
 scalaris 140
 scaliger 140
 Sceloporus XIV
 Schottii 148
 Scincidae XIV
 Scalceophidia XIX, 1, 2, 129
 Scalceophis 163
 Scorpion XIV
 Scotophis 53, 151
 scutulatus 113, 172
 Sextale, Heterodon 76, 160
 Sextale, Lachesis 179
 Sextalimae XXI, 11, 18, 136
 sumichrasti 140
 surcunco 179
 Sympholis 167
- Synonymy 129
 Systematic List 129
- T.**
- Tachymenis 60, 154
 taeniatus, Bothrops 180
 taeniatus, Coluber 46, 148
 Taeniophis 61
 Tantilla 86, 162
 taxispilotus 26, 141
 Teiidae XIV
 Teius XIV
 tener 169
 tenuiculum 5, 131
 tenuis 165
 Telocraspis 126, 180
 tephropleura 143
 tergeminus 176
 terrificus 171
 tessellatus, Bothrops 180
 tessellatus, Rhinocheilus 74, 159
 testaceus 43, 148
 Testudinata IV
 Testudo V
 toxicensis 158
 Thalassochelys VI
 Thamnophis 138
 Thysanodactylus XIII
 tigrinus 159
 tigris 117, 175
 Tisiphone 177
 Tolia 82, 161
 Tomodon 154
 torquatum 18, 135
 Tortrix 131
 Toxicophidia XXII, 1, 104, 168
 Toxicophis 177
 Trachydosaurus XV
 transversus 141
 Tree Snakes XXI
 Tretanorhinus 33, 143
 triangulus 65, 155
 tricinctus 157
 tricolor 153
 Trigonoccephalus, 123, 178, 179, 180
 triseriatus 112, 117, 175
 tristis 169
 trivirgata 8, 131
 Trionycidae IV
 Tropidoclonion 143
 Tropidodipsas 12
 Tropidonotus 22, 136
 Trunkbacks VI
 Turtle's nests VII
 twiningii 138
 Typhlopidae XIX, 2, 129
 Typhlopinae 2, 129
 Typhlops 2, 129
- U.**
- undulatus 126, 180
 upsilon 134
 Uroecrotalon 170, 174
 Urodela XXVIII, XXIX
 Urotophus 170
 ustus 142
- V.**
- vagrans 139
 valeriae 98, 166
 validus 28, 143
 Varanidae XI
 variabilis, Helicops 33, 144
 variabilis, Spilotes 49, 149

varians	161	Vipera	168	Xenodon	X.	
variegata, Ficimia	81, 162	Virginia	96, 166	Xerobates		79, 161
variegata, Hydrophis	170	vitatus	158	Xerobates		xi
Venom	xviii	vulpinus	56, 153		Y.	
venustissimus	63			Yellow Bat		xvii, 132
venustus	143				Z.	
veniformis	167			Zachobus		156
vernix	101, 167	W.		zonatus		67, 156
vernalis	39, 146	wagneri	88, 163	Zoumriidae		xv
vertebratis	150	Water Snake	140	Zootoca		xiv
vetustus	147	Water Snakes	xxi			
villarsii	85	weigelti	134			
		woodhousei	141			




1. STORERIA DERAYI 2. S. OCCIPITOMACULATA 3. REGINA KITCHLANDI

4. STORERIA LINEATA 5. HYDROPS ACAECUS



1 REGINA LEBEIS. 2 DIADOPHIS PUNCTATUS

3. NERODIA SIPEDON 4. N. CYLOPIUS




1. PSEUDOPHIS AESTIVUS. 2. EUTAENIA SAURITA.

3. E. SIRTALIS. 4. CYCLOPHIS VERNALIS


L. Trouessart del.

Engraved by A. Mouss.




1. ELAPHE GUTTATA. 2. ELAPHE OBSOLETUS.

3. UROTHER CONSTRICTOR.



1. *Ophiboles triangularis*. 2. *O. dolivatus*.


3. *O. getulus*. 4. *O. sami*.



L. Trouessart d. r.


Printed by A. Mozer

1. *CEMOPHORA COCCINEA*. 2. *CONTIA EPISCOPA*. 3. *TANTILLA GRACILIS*
 4. *HETERODON SIMUS*. 5. *H. PLATYRHINUS*. 6. *H. NASICUS*.




1. CARPHOPHIS AMOENA 2. VIRGINIA STRIATULA 3. V. NUBERAE.

4. HELICOPS ALLESHI 5. COSTIA PUGAEA



1. ASCISTRODON COSTORTRIX. 2. A. PISCIVORUS VAR. PUGNAX.



1. CROTALUS HORRIDUS var. ATRICAUDATUS





QL
1
H35

v.8
Biological
& Medical
Serials

Harvard University. Museum
of Comparative Zoology
Memoirs

PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET

UNIVERSITY OF TORONTO LIBRARY

STORAGE

